

REFERENCES

- Agency for Toxic Substances and Disease Registry (ATSDR). Health effects in humans: DEET (*N,N*-diethyl-meta-toluamide) chemical technical summary for public health and public safety professionals [serial online] 2012 Dec [cited 2014 Jun 25] Available from: URL: <http://www.atsdr.cdc.gov/consultations/deet/health-effects.html>
- al-Maskri AY, Hanif MA, al-Maskari MY, et al. Essential oil from *Ocimum basilicum* (Omani Basil): a desert crop. *Nat Prod Commun* 2011;6:1487-90.
- Amer A, Mehlhorn H. Repellency effect of forty-one essential oils against *Aedes*, *Anopheles*, and *Culex* mosquitoes. *Parasitol Res* 2006;99:478-90.
- Amerasan D, Murugan K, Kovendan K, Mahesh KP, Panneerselvam C, Subramaniam J, et al. Adulticidal and repellent properties of *Cassia tora* Linn. (Family: Caesalpinaceae) against *Culex quinquefasciatus*, *Aedes aegypti*, and *Anopheles stephensi*. *Parasitol Res* 2012;111(5):1953-64.
- Andrew J, Bar A. Morphology and morphometry of *Aedes aegypti* adult mosquito. *Annu Rev & Res in Bio* 2013;3(1):52-69.
- Ansari MA, Mittal PK, Razdan RK, Sreehari U. Larvicidal and mosquito repellent activities of pine (*Pinus longifolia*, family: Pinaceae) oil. *J Vector Borne Dis* 2005;42:95-9.
- Ansari MA, Razdan RK, Tandon M, Vasudevan P. Larvicidal and repellent actions of *Dalbergia sissoo* Roxb. (*F. leguminosae*) oil against mosquitoes. *Bioresour Technol* 2000;73:207-11.
- Badolo A, Ilboudo-Sanogo E, Oue'draogo AP, Costantini C. Evaluation of the sensitivity of *Aedes aegypti* and *Anopheles gambiae* complex mosquitoes to two insect repellents: DEET and KBR 3023. *Trop Med Intl Health* 2004;9:330-4.

Barnard DR, Xue RD. Laboratory evaluation of mosquito repellents against *Aedes albopictus*, *Culex nigripalpus*, and *Ochlerotatus triseriatus* (Diptera: Culicidae). J Med Entomol 2004;41:726-30.

Beaty BJ, Marquardt WC. The Biology of Disease Vectors. Colorado: Am J Trop Med 1997;56:482.

Bell JW, Veltri JC, Page BC. Human exposures to *N,N*-diethyl-*m*-toluamide insect repellents reported to the American Association of Poison Control Centers 1993-1997. Int J Toxicol 2002;21:341-52.

Blackwell A, Stuart AE, Estambale BA. The repellent and antifeedant activity of oil of *Myrica gale* against *Aedes aegypti* mosquitoes and its enhancement by the addition of salicylic acid. J R Coll Physicians Edinb 2003;33:209-14.

Briassoulis G, Narlioglou M, Hatzis T. Toxic encephalopathy associated with use of DEET insect repellents: a case analysis of its toxicity in children. Hum Exp Toxicol 2001;20:8-14.

Carpenter SJ, LaCasse WJ. Mosquitoes of North America (North of Mexico). Berkeley, Los Angeles, London: University of California Press, 1955.

Carroll SP, Loye J. PMD, a registered botanical mosquito repellent with deet-like efficacy. J Am Mosq Control Assoc 2006;22:507-14.

Centers for Disease Control and Prevention (CDC). *Anopheles* mosquitoes [serial online] 2012a Nov [cited 2014 Jun 25]. Available from: URL: <http://www.cdc.gov/malaria/about/biology/mosquitoes/>

Centers for Disease Control and Prevention (CDC). Dengue and the *Aedes aegypti* mosquito [serial online] 2012b Jan [cited 2013 Sep 15]. Available from: URL: http://www.cdc.gov/dengue/resources/30jan2012/aegyptifact_sheet.pdf

Changbunjong T, Weluwanarak T, Taowan N, Suksai P, Chamsai T, Sedwisai P. Seasonal abundance and potential of Japanese encephalitis virus infection in mosquitoes at the nesting colony of ardeid birds, Thailand. Asian Pac J Trop Biomed 2013;3:207-10.

Choochote W, Chaithong U, Kamsuk K, Jitpakdi A, Tippawangkosol P, Tuetun B et al. Repellent activity of selected essential oils against *Aedes aegypti*. *Fitoterapia* 2007;78:359-64.

Choochote W, Kanjanapothi D, Panthong A, Taesotikul T, Jitpakdi A, Chaithong U, et al. Larvicidal, adulticidal and repellent effects of *Kaempferia galanga*. *Southeast Asian J Trop Med Public Health* 1999;30:470-6.

Christophers SR. *Aedes aegypti* (L.) The yellow fever mosquito. Its life history, bionomics and structure: Cambridge University Press, 1960.

Clements AN. The physiology of mosquitoes. Oxford: pergamon Press;1992:395.

Conti B, Benelli G, Leonardi M, Afifi FU, Cervelli C, Profeti R, et al. Repellent effect of *Salvia dorisiana*, *S. longifolia*, and *S. sclarea* (Lamiaceae) essential oils against the mosquito *Aedes albopictus* Skuse (Diptera: Culicidae). *Parasitol Res* 2012;111:291-9.

Copeland RS, Walker TW, Robert LL, Gihure JI, Wirtz RA, Klein TA. Response of wild *Anopheles funestus* to repellent-protected volunteers is unaffected by malaria infection of the vector. *J Am Mosq Control Assoc* 1995;15:342-7.

Costanzo SD, Watkinson AJ, Murby EJ, Kolpin DW, Sandstrom MW. Is there a risk associated with the insect repellent DEET (*N,N*-diethyl-*m*-toluamide) commonly found in aquatic environments?. *Sci Total Environ* 2007;384:214-20.

Curtis CF, Lines JD, Lu B, Renz A. Natural and synthetic repellent. In: Curtis CF, ed. *Control of Disease Vectors in the Community*. London: Wolf Publishers Ltd., 1991:75-92.

Curtis CF. Personal protection methods against vectors of disease. *Rev Med Vet Entomol* 1992;80:543-53.

Das NG, Nath DR, Baruah I, Talukdar PK, Das SC. Field evaluation of herbal mosquito repellents. *J Commun Dis* 1999;31:241-5.

Davis EE. Insect repellents: concepts of their mode of action relative to potential sensory mechanisms in mosquitoes (Diptera: Culicidae). J Med Entomol 1985;22:237-43.

de Garbino JP, Laborde A. Toxicity of an insect repellent: *N,N*-diethyltoluamide. Vet Hum Toxicol 1983;25:422-3.

Dethier VG. Chemical insect attractants and repellents. Philadelphia: Blakiston Co., 1947.

Dhandapani A, Kadarkarai M. HPTLC quantification of flavonoids, larvicidal and smoke repellent activities of *Cassia occidentalis* L. (Caesalpiniaceae) against malarial vector *Anopheles stephensi* Lis (Diptera: Culicidae). J Phytolo 2011;3:60-72.

Dremova VP, Markina VV, Kamennov NA. How evaporation and absorption affect the formulation of various insect repellents. Int Pest Cont 1971;13:13-6.

Elango G, Rahuman AA, Zahir AA, Kamaraj C, Bagavan A, Rajakumar G, et al. Evaluation of repellent properties of botanical extracts against *Culex tritaeniorhynchus* Giles (Diptera: Culicidae). Parasitol Res 2010;107:577-84.

Fradin MS. Protection from blood-feeding arthropods. In Auerbach PS, editor. Wilderness medicine. 4th ed. Mosby, St. Louis; 2001. p. 754-68.

Fradin MS, Day JF. Comparative efficacy of insect repellents against mosquito bites. N Engl J Med 2002;347:13-8.

Gillij YG, Gleiser RM, Zygadlo JA. Mosquito repellent activity of essential oils of aromatic plants growing in Argentina. Bioresour Technolo 2008;99:2507-15.

Goodard J. Setting up a mosquito control program [serial online]. Mississippi State Department of Health; 2003 Jun [cited 2013 Nov 8]. Available from: URL: http://msdh.ms.gov/msdhsite/_static/resources/800.pdf

Govindarajan M, Sivakumar R. Mosquito adulticidal and repellent activities of botanical extracts against malarial vector, *Anopheles stephensi* Liston (Diptera: Culicidae). Asian Pac J Trop Med 2011;4:941-7.

Govindarajan M, Rajeswary M, Sivakumar R. Repellent properties of *Delonix elata* (L.) Gamble (Family: Fabaceae) against malaria vector *Anopheles stephensi* (Liston) (Diptera: Culicidae) J Saudi Soc of Agri Sci 2014 In press.

Gratz NG. Space sprays for the control of *Aedes aegypti* in South-East Asia and the Western Pacific. Dengue Bulletin 1999;23:80-4.

Gupta RK, Rutledge LC. Role of repellents in vector control and disease prevention. Am J Trop Med Hyg 1994;50:82-6.

Gwari G, Bhandari U, Andola HC, Lohani H, Chauhan N. Volatile constituents of *Saussurea costus* roots cultivated in Uttarakhand Himalayas, India. Pharmacognosy Res 2013;5:179-82.

Harbach R. Mosquito Taxonomic Inventory (MTI) [serial online] 2013 Aug [cited 2013 Sep 1]. Available from: URL: www.mosquito-taxonomic-inventory.info/

Harborne J B. Phytochemical methods: A guide to modern technique of plant analysis. London: Chapman and Hall, 1984.

Huang YM. The subgenus *Stegomyia* of *Aedes* in the oriental region with keys to the species (Diptera: Culicidae). Contrib Amer Ent Inst 1979;15:6.

Huang YZ, Pu FD. Studies on the chemical components of the essential oil from the rhizome of *Ligusticum sinense* Oliv. cv. Chuanxiong hort. Acta Pharm Sin 1988; 23:426-9.

IBACON. Storage stability and shelf life testing [serial online] (n.d.) [cited 2014 July 2]. Available from: URL: http://www.ibacon.de/tl_files/files/test-systems/analytical/storage_stability_shelf_life_testing.pdf.pdf

Jensenus M, Pretorius AM, Clarke F, Myrvang B. Repellent efficacy of four commercial DEET lotions against *Amblyomma hebraeum* (Acari: Ixodidae), the

principal vector of *Rickettsia africae* in southern Africa. *Trans R Soc Trop Med Hyg* 2005;99:708-11.

Juengprasert W, Techadamrongsin Y, Soonthornchareonnon N, Sontanaphun U. Standard of Chinese Medical in Thailand Chuanxiong Rhizoma. Department for development of Thai traditional and alternative medicine, Ministry of Public Health, Thailand in cooperation with Chengdu University of Traditional Chinese medicine people's republic of China, 2011;136.

JunHua H, LiNa M, Chun R, HongJun L, TingShan Y, HaoQiang L, et al. Acaricidal activity of extract of *Ligusticum sinense* against *Panonychus citri*. *S Chi J of Agri Sci* 2010;23:405-9.

Kamsuk K, Choochote W, Chaithong U, Jitpakdi A, Tippawangkosol P, Riyong D, et al. Effectiveness of *Zanthoxylum piperitum*-derived essential oil as an alternative repellent under laboratory and field applications. *Parasitol Res* 2007;100:339-45.

Karunamoorthi K, Girmay A, Hayleeyesus SF. Mosquito repellent activity of essential oil of Ethiopian ethnomedicinal plant against Afro-tropical malarial vector *Anopheles arabiensis*. *J King Saud Uni-Sci*. 2014 In press.

Karunamoorthi K, Sabesan S. Laboratory evaluation of dimethyl phthalate treated wristbands against three predominant mosquito (Diptera: Culicidae) vectors of disease. *Eur Rev Med Pharmacol Sci* 2010;14:443-8.

Kengluetcha A, Sithiprasasna R, Boonsuepsakul S, Sithiprasasna R, Rodpradit P, Baimai V. Geographical distribution of *Anophelss minimus* species A and C in western Thailand. *Southeast Asian J Trop Med Public Health* 2004;35:918-26.

Khan M, Sricastava SK, Syamasundar KV, Singh M, Naqvi AA. Chemical composition of leaf and flower essential oil of *Lantana camara* from India. *Flavour Fragr J* 2002;17:75-7.

Kim JK, Kang CS, Lee JK, Kim YR, Han HY, Yun HK. Evaluation of repellency effect of two natural aroma mosquito repellent compounds, citronella and citronellal. *Entomol Res* 2005;35:117-20.

Kim SI, Yoon JS, Baeck SJ, Lee SH, Ahn YJ, Kwon HW. Toxicity and synergic repellency of plant essential oil mixtures with vanillin against *Aedes aegypti* (Diptera: Culicidae). J Med Entomol 2012;49:876-85.

Kittayapong P, Chansang U, Chansang C, Bhumiratana A. Community participation and appropriate technologies for dengue vector control at transmission foci in Thailand. J Am Mosq Control Assoc 2006;22:538-6.

Knight KL, Stone A. A catalog of the mosquitoes of the world (Diptera: Culicidae), 2nd ed. Entomol Soc Am; 1997.

Kumar R, Kumar A, Prasal CS, Dubey NK, Samant R. Insecticidal activity *Aegle marmelos* (L.) Correa essential oil against four stored grain insect pests. Internet J Food Safety 2008;10:39-49.

Leung AY, Foster S. Encyclopedia of common natural ingredients used in food, drugs, and cosmetics. 2nd ed. Wiley-Interscience; 1996.

Li HY. Screening and applicated studies in kipper process on the repellent and ovicidal essential oils for flies [serial online] 2008 Apr [cited 2014 Jun 7]. Available from: URL: <http://www.globethesis.com/?t=2121360215992357>

Limsuwan S, Rongsriyam Y, Kerdpibule V, Apiwathnasorn C, Chiang GL, Cheong WH. Rearing techniques for mosquitoes. In:Sucharit S, Supavej S Eds. Practical Entomology. Malaria and filariasis, 1st ed. Museum and Reference Center, Faculty of Tropical Medicine, Mahidol University, Thailand, 1987.

Liu ZL, HE Q, Chu SS, Wang CF, Du SS, Deng ZW. Essential oil composition and larvicidal activity of *Saussurea lappa* roots against the mosquito *Aedes albopictus* (Diptera: Culicidae). Parasitol Res 2012;110:2125-30.

Luo YM, Zhang JH, Pan JK, Yao SL, Huang HL, Zhu Y. The constituents of *Ligusticum sinense* Oliv cv. chaxiong. Chinese Phar J 1994;29:714-6.

Luo YM. Research progress of *Ligusticum sinense* Oliv cv. chaxiong. Jiangxi J Trad Chin Med 1998;10:87-9.

- Lupi E, Hatz C, Schlagenhauf P. The efficacy of repellents against *Aedes*, *Anopheles*, *Culex* and *Ixodes* spp. - A literature review. *Travel Med Infect Dis* 2013;11:374-411.
- Maia MF, Moore SJ. Plant-based insect repellents: a review of their efficacy, development and testing. *Malar J* 2011;10:S11.
- Mandal S. Repellent activity of *Eucalyptus* and *Azadirachta indica* seed oil against the filarial mosquito *Culex quinquefasciatus* Say (Diptera: Culicidae) in India. *Asian Pac J Trop Biomed* 2011;S109-12.
- Manner HI, Buker RS, Easton V, Elevitch CR. *Citrus* species (citrus) and *Fortunella* (kumquat) Rutaceae (rue family). Species profiles for pacific island agroforestry 2006.
- Mehta RM. *Pharmaceutics-I*. Delhi: Vallabh Prakashan; 2002.
- Miot HA, Batistella RF, Batista KA, Volpato DEC, Augusto LST, Madeira NG, et al. Comparative study of the topical effectiveness of the Andiroba oil (*Carapa guianensis*) and DEET 50% as repellent for *Aedes* sp. *Rev Inst Med Trop Sao Paulo* 2004;46:235-6.
- Mosquito and Vector Control Association of California (MVCAC). Mosquito Control FAQs [serial online] 2005 Jul [cited 2013 Sep 17]. Available from: URL:<http://www.cdph.ca.gov/HealthInfo/discond/Documents/MosquitoControlFAQs.pdf>
- Mourya DT, Ilkal MA, Mishra AC, Jacob PG, Pant U, Ramanujam S, et al. Isolation of Japanese encephalitis virus from mosquitoes collected in Karnataka state, India from 1985 to 1987. *Trans R Soc Trop Med Hyg* 1989;83:550-2.
- Müller GC, Junnila A, Kravchenko VD, Revay EE, Butlers J, Schlein Y. Indoor protection against mosquito and sand fly bites: a comparison between citronella, linalool and geraniol candles. *J Am Mosq Control Assoc* 2008;24:150-3.
- Murugan K, Mahesh Kumar P, Kovendan K, Amerasan D, Subrmaniam J, Hwang JS. Larvicidal, pupicidal, repellent and adulticidal activity of *Citrus sinensis* orange

peel extract against *Anopheles stephensi*, *Aedes aegypti* and *Culex quinquefasciatus* (Diptera: Culicidae). Parasitol Res 2012;111:1757-69.

Negi JS, Bisht VK, Bhandari AK, Bhatt, V.P, Sati MK, et al. Antidiarrheal activity of methanol extract and major essential oil contents of *Saussurea lappa* Clarke. Afr J Pharm Pharmacol 2013;7:474-7.

Newman M, Lhuillier A, Poulsen AD. Checklist of the Zingiberaceae of Malaysia. Blumea Suppl 2004;16.

Nitatpattana N, Dubot-Pérès A, Gouilh MA, Souris M, Barbazan P, Yoksan S, et al. Change in Japanese encephalitis virus distribution, Thailand. Emerg Infect Dis 2008;14:1762-5.

Noosidum A, Prabaripai A, Chareonviriyaphap T, Chandrapatya A. Excito-repellency properties of essential oils from *Melaleuca leucadendron* L., *Litsea cubeba* (Lour.) Persoon, and *Litsea salicifolia* (Nees) on *Aedes aegypti* (L.) mosquitoes. J Vector Ecol 2008;33:305-12.

Nurzyńska-wierdak R, Borowski B, Dzida K, Zawislak R, Kowalski R. Essential oil composition of sweet basil cultivars as affected by nitrogen and potassium fertilization. Turk J Agric For 2013;37:427-36.

O'Malley. The biology of *Anopheles quadrimaculatus* Say. Proceedings of the seventy-ninth annual meeting of the New Jersey mosquito control association, Inc. 1992;136-44.

Osimitz TG, Grothaus RH. The present safety assessment of DEET. J Am Mosq Control Assoc 1995;11:274-8.

Osimitz TG, Murphy JV. Neurological effects associated with use of the insect repellent *N,N*-diethyl-*m*-toluamide (DEET). J Toxicol Clin Toxicol 1997;35:435-41.

Oyedele A.O, Gbolade A.A, Sosan M.B, Adewoyin FB, Soyelu OL, Orafidiya OO. Formulation of an effective mosquito-repellent topical product from Lemongrass oil. Phytomedicine 2002;9:259-262.

Pålsson K, Jaenson TG. Comparison of plant products and pyrethroid-treated bed nets for protection against mosquitoes (Diptera: Culicidae) in Guinea Bissau, West Africa. *J Med Entomol* 1999;36:114-8.

Panneerselvam C, Murugan K. Adulticidal, repellent, and ovicidal properties of indigenous plant extracts against the malarial vector, *Anopheles stephensi* (Diptera: Culicidae). *Parasitol Res* 2013;112:679-92.

Patel PK, Prajapati NK, Dubey BK. *Madhuca indica*: A review of its medicinal properties. *IJPSR* 2012;3:1285-93.

Pethuan S, Jirakanjanakit N, Saengtharatip S, Chareonviriyaphap T, Kaewpa D, Rongnoparut P. Biochemical studies of insecticide resistance in *Aedes (Stegomyia) aegypti* and *Aedes (Stegomyia) albopictus* (Diptera: Culicidae) in Thailand. *Trop Biomed* 2007;24:7-15.

Petrucci N, Sardini S. Severe neurotoxic reaction associated with oral ingestion of low-dose diethyltoluamide-containing insect repellent in a child. *Pediatr Emerg Care* 2000;16:341-2.

Phanpoowong T, Lek-Uthai U, Thongrungrat S, Komalamisra N, Srisawat R, Russell B, et al. Dengue-2 virus carrying capacity of Thai *Aedes aegypti* strains with different susceptibility to deltamethrin. *Southeast Asian J Trop Med Public Health* 2012;43:3.

Phasomkusolsil S, Soonwera M. Insect repellent activity of medicinal plant oils against *Aedes aegypti* (Linn.), *Anopheles minimus* (Theobald) and *Culex quinquefasciatus* Say based on protection time and biting rate. *Southeast Asian J Trop Med Public Health* 2010;41:831-40.

PHE, Public Health England. Dengue fever in England, Wales and Northern Ireland: 2012 [serial online] 2012 May [cited 2013 Nov 8]. Available from: URL: http://www.hpa.org.uk/webc/HPAwebFile/HPAweb_C/1317138961007

- Qui H, Jun HW, John WM. Pharmacokinetics, formulation, and safety of insect repellent *N,N*-diethyl-3-methylbenzamide (DEET): a review. *J Am Mosq Control Assoc* 1998;14:12-27.
- Ratanatham S, Upatham ES, Prasittisuk C, Rojanasunan W, Theerasilp N, Tremongkol A, et al. Bionomics of *Anopheles minimus* and its role in malaria transmission in Thailand. *Southeast Asian J Trop Med Public Health* 1988;19:83-9.
- Rattanarithikul R, Panthusiri P. Illustrated keys to the medically important mosquitoes of Thailand. *Southeast Asian J Trop Med Public Health* 1994;25:1-66.
- Rehman JU, Ali A, Khan IA. Plant based products: use and development as repellents against mosquitoes: A review. *Fitoterapia* 2014;95:65-74.
- Rozendaal JA. Vector control: Methods for use by individuals and communities. Geneva: World Health Organization, 1997.
- Satyal P, Woods KE, Dosoky NS, Neupane S, Setzer WN. Biological activities and volatile constituents of *Aegle marmelos* (L.) Correa from Nepal. *J Med Active Plants* 2012;1:114-22.
- Sax NI, Lewis RJ. Dangerous properties of industrial materials. 7th ed. New York: Van Nostrand Reinhold; 1989.
- Shalan E, Canyon D, Faried MW, Abdel-Wahab H, Mansour AH. A review of botanical phytochemicals with mosquitocidal potential. *Environ Int* 2005;31:1149-66.
- Shadia E, el-Aziz A, Omer EA, Sabra AS. Chemical composition of *Ocimum americanum* essential oil and its biological effects against, *Agrotis ipsilon*, (Lepidoptera: Noctuidae). *Res J Agric & Biol Sci* 2007;3:740-7.
- Sharma VP, Ansari MA. Personal protection from mosquitoes (Diptera: Culicidae) by burning neem oil in kerosene. *J Med Entomol* 1994;31:95-102.
- Sharma VP, Ansri MA, Razdan RK. Mosquito repellent action of neem (*Azadirachta indica*) oil. *J Am Mosq Control Assoc* 1993;9:359-60.

- Singh SP, Mittal PK. Mosquito repellent and oviposition deterrent activities of *Solanum nigrum* seed extract against malaria vector *Anopheles stephensi*. OIIRJ 2013;3:326-33.
- Siriyasatien P, Tangthongchaiwiriya K, Jariyapan N, Kaewsaitiam S, Poovorawan Y, Thavara U. Analysis of salivary gland proteins of the mosquito *Armigeres subalbatus*. Southeast Asian J Trop Med Public Health 2005;36:64-7.
- Solomon B, Sahle FF, Gebre-Mariam T, Asres K, Neubert RH. Microencapsulation of citronella oil for mosquito-repellent application: formulation and *in vitro* permeation studies. Eur J Pharm Biopharm 2012;80:61-6.
- Sritabutra D, Soonwera M. Repellent activity of herbal essential oils against *Aedes aegypti* (Linn.) and *Culex quinquefasciatus* (Say.). Asian Pac J Trop Dis 2013;3:271-6.
- Sucharit S, Choochote W. Comparative studies on the morphometry of male genitalia and frequency of claspers movements during induced copulation of *Anopheles balabacensis* (Perlis form) and *Anopheles dirus* (Bangkok Colony strain). Mosq syst 1983;15:90-6.
- Sukumar K, Perich MJ, Boobar LR. Botanical derivatives in mosquito control: a review. J Am Mosq Control Assoc 1991;7:210-36.
- Tanaka K, Mizusawa K, Saugstad ES. A revision of the adult and larval mosquitoes of Japan (including the Ryukyu Archipelago and the Ogasawara Islands) and Korea (Diptera: Culicidae). Contrib Amer Ent Inst 1979;16:1-987.
- Tawatsin A, Asavadachanukorn P, Thavara U, Wongsinkongman P, Bansidhi J, Boonruad T, et al. Repellency of essential oils extracted from plants in Thailand against four mosquito vectors (Diptera: Culicidae) and oviposition deterrent effects against *Aedes aegypti* (Diptera: Culicidae). Southeast Asian J Trop Med Public Health 2006;37:915-31.

Tawatsin A, Wrattenn SD, Scott RR, Thavara U, Techadamrongsin Y. Repellency of volatile oils from plants against three mosquito vectors. *J Vector Ecol* 2001;26:76-82.

Tenenbein M. Severe toxic reactions and death following the ingestion of diethyltoluamide- containing insect repellents. *JAMA* 1987;258:1509-11.

Thorsell W, Mikiver A, Malander I, Tunon H. Efficacy of plant extracts and oils as mosquito repellents. *Phytomedicine* 1998;4:311-23.

Trongtokit Y, Rongsriyam Y, Komalamisra N, Apiwathnasorn C. Comparative repellency of 38 essential oils against mosquito bites. *Phytother Res* 2005;19:303-9.

Tuetun B, Choochote W, Kanjanopothi D, Rattanachanpichai E, Chaithong U, Chaiwong P, et al. Repellent properties of celery, *Apium graveolens* L., compared with commercial repellents, against mosquitoes under laboratory and field conditions. *Trop Med Int Health* 2005;10:1190-8.

Tuetun B, Choochote W, Pongpaibul Y, Junkum A, Kanjanapothi D, Chaithong U, et al. Celery-based topical repellents as a potential natural alternative for personal protection against mosquitoes. *Parasitol Res* 2008;104:107-15.

Tuetun B, Choochote W, Pongpaibul Y, Junkum A, Kanjanapothi D, Chaithong U, et al. Field evaluation of G10, a celery (*Apium graveolens*)-based topical repellent, against mosquitoes (Diptera: Culicidae) in Chiang Mai province, northern Thailand. *Parasitol Res* 2009;104:515-21.

Tuetun B, Choochote W, Rattanachanpichai E, Chaithong U, Jitpakdi A, Tippawangkosol P, et al. Mosquito repellency of the seeds of celery (*Apium graveolens* L.). *Ann Trop Med Parasitol* 2004;98:407-17.

Turek C, Stintzing FC. Stability of essential oils: A review. Article first published online [serial online] 2012 Jan [cited 2014 July 1]. Available from: URL: <http://onlinelibrary.wiley.com/doi/10.1111/1541-4337.12006/pdf>

Tyagi BK, Shahi AK, Kaul BL. Evaluation of repellent activities of *Cymbopogon* essential oils against mosquito vectors of malaria, filariasis and dengue fever in India. *Phytomedicine* 1998;5:324-9.

United States Environmental Protection Agency (EPA). Oil of citronella, EPA-738-F-97-002 [serial online] 1997 Feb [cited 2014 Jun 25]. Available from: URL: <http://www.epa.gov/oppsrrd1/REDS/factsheets/3105fact.pdf>

Veltri JC, Osimitz TG, Bradford DC, Page BC. Retrospective analysis of calls to poison control centers resulting from exposure to the insect repellent *N,N*-diethyl-*m*-toluamide (DEET) from 1985-1989. *J Toxicol Clin Toxicol* 1994;32:1-16.

Venkatachalam MR, Jebanesan A. Repellent activity of *Ferronia elephantum* Corr. (Rutaceae) leaf extract against *Aedes aegypti* (L.). *Bioresour Technol* 2001;76:287-8.

Verma RS, Adalia RC, Chauhan A. Essential oil composition of *Aegle marmelos* (L.) Correa: chemotypic and seasonal variations. *J Sci Food Agric* 2013;94:1904-13.

Vieira RF, Simon JE. Chemical characterization of basil (*Ocimum* spp.) found in the markets and used in traditional medicine in Barzil. *Econ Bot* 2000;54:207-16.

Vineetha AR, Murugan KA. Larvicidal and smoke repellency effect of *Toddalia asiatica* and *Aegle marmelos* against the dengue vector, *Aedes aegypti* (Insecta: Diptera: Culicidae). *Entomol Res* 2009;39:61-5.

Wandscheer CB, Duque JE, da Silva MAN, Fukuyama Y, Wohlke JL, Adelman J, et al. Larvicidal action of ethanolic extract from fruit endocarps of *Melia azedarach* and *Azadirachta indica* against the dengue mosquito *Aedes aegypti*. *Toxicon* 2004;44:829-35.

Wang J, Xu L, Yang L, Liu Z, Zhou L. Composition, antibacterial and antioxidant activities of essential oils from *Ligusticum sinense* and *L. jeholense* (Umbelliferae) from China. *Rec Nat Prod* 2011a;5:4:314-8.

- Wang J, Yang JB, Wang AG, Ji TF, Su YL. Studies on the chemical constituents of *Ligusticum sinense*. Chin J Med Mat 2011b;34:378-80.
- Wanga Z, Songb J, Chena J, Song Z, Shang S, Jiang Z, et al. QSAR study of mosquito repellents from terpenoid with a six-member-ring. Bioorg Med Chem Lett 2008;18:2854-9.
- Warrell DA, Gilles HM. Essential malariology, 4th ed. London: Hodder Arnold, MO: Mosby year book; 2002.p. 352.
- Wei Q, Yang J, Ren J, Wang A, Ji T, Su Y. Bioactive phthalides from *Ligusticum sinense* Oliv cv. Chaxiong. Fitoterapia 2014;93:226-32.
- World Health Organization. International travel and health vaccination requirement and health advice. Geneva: WHO; 1995.
- World Health Organization. Report of the WHO informal consultation on the evaluation and testing of insecticides. CTD/WHOPES/IC/96.1, Control of Tropical Diseases Division. Geneva: WHO; 1996.
- Yang P, Ma Y. Repellent effect of plant essential oils against *Aedes albopictus*. J Vector Ecol 2005;30:231-4.
- Yang YC, Lee EH, Lee HS, Lee DK, Ahn YJ. Repellency of aromatic medicinal plant extracts and a steam distillate to *Aedes aegypti*. J Am Mosq Control Assoc 2004;20:146-9.
- Yap HH, Jahangir K, Chong AS, Adanan CR, Chong NL, Malik YA. Field efficacy of a new repellent, KBR 3023, against *Aedes albopictus* (Skuse) and *Culex quinquefasciatus* (Say) in a tropical environment. J Vector Ecol 1998;23:62-8.
- Zadikoff CM. Toxic encephalopathy associated with use of insect repellent. J Pediatr 1979;95:140-2.
- Zhu Y, Luo YM. Preliminary study of polysaccharide of *L. chaxiong*. Jiangxi J Trad Chin Med 1997;9:32-3.