

Reference

- [1] Kolhaker AG, Jamison AC, Litvinov D, Willson RC, Lee TR. Tuning the Magnetic Properties of Nanoparticles. *International Journal Molecular Sciences*. 2013, 14: 15977-16009.
- [2] Krishnan, K.M.; Pakhomov, A.B.; Bao, Y.; Blomqvist, P.; Chun, Y.; Gonzales, M.; Griffin, K.; Ji, X.; Roberts, B.K. Nanomagnetism and spin electronics: Materials, microstructure and novel properties. *Journal of Materials Science*. 2006, 41, 793–815.
- [3] Sosnovik DE, Nahrendorf M, Weissleder Ralph. Magnetic nanoparticles for MRI imaging : agent, techniques and cardiovascular applications. *Basic Research in Cardiology*. 2008, 103, 122-130.
- [4] Sanjai S, Kothan K, Gonil P, Saesoo S, Sajomsang W. Chaitosan-triphosphate nanoparticles for encapsulation of super-paramagnetic iron oxide as an MRI contrast agent. *Carbohydrate Polymers*. 2014, 104, 231-237.
- [5] Hu Y, Meng L, Niu L, Lu Q. Highly Cross-Linked and Biocompatible Polyphosphazene-Coated Superparamagnetic Fe₃O₄ Nanoparticles for Magnetic Resonance Imaging. *Langmuir*. 2013, 29, 9156-9163.
- [6] Li Y, Chen Z, Jichen Wang, Zhang Z. Preparation and in vitro studies of MRI-specific superparamagnetic iron oxide antiGPC3 probe for hepatocellular carcinoma. *International Journal of Nanomedicine*. 2012, 7, 4593-4611.
- [7] Hemmingsson A, Carlsten J, Ericsson A et al. Relaxation enhancement of the dog liver and spleen by biodegradable superparamagnetic particles in proton magnetic resonance imaging. *Acta Radiol* 1987, 28, 703–705

- [8] Fukukura Y, Kamiyama T, Takumi K, Shindo T, Higashi R, Nakajo M. Comparison of ferucarbotran enhanced fluid-attenuated inversion recovery echo-planar, T2-weighted turbo spinecho, T2*-weighted gradient-echo, and diffusion-weighted echo-planar imaging for detection of malignant liver lesions. *Journal of Magnetic Resonance Imaging*. 2010, 31, 607-16.
- [9] Indhira TK, Lakshmi PK, Magnetic nanoparticles – a review, *International Journal of Pharmacy and Pharmaceutical Sciences*. Nanotech. 3. 2010, 1035–1042.
- [10] Bronstein LM, Huang X, Retrum J, Schmucker A, Pink M, Stein BD, Dragnea B. Influence of iron oleate complex structure on iron oxide nanoparticle formation. *Journal of Materials Chemistry*. 2007, 19, 3624–3632.
- [11] Briley-Saebo K, Bjornerud A, Grant D, Ahlstrom H, Berg T, Kindberg GM. Hepatic cellular distribution and degradation of iron oxide nanoparticles following single intravenous injection in rats: Implications for magnetic resonance imaging. *Cell and Tissue Research*. 2004, 316, 315–323.
- [12] Herrera AP, Barrera C, Zayas Y, Rinaldi C. Monitoring colloidal stability of polymer-coated magnetic nanoparticles using AC susceptibility measurements. *Journal of Colloid and Interface Science*. 2010, 342, 540–549.
- [13] Clement JH, Schwalbe M, Buske N, Wagner K, Schnabelrauch M, Gornert P, Kliche KO, Pachmann K, Weitschies W, Hoffken K. Differential interaction of magnetic nanoparticles with tumor cells and peripheral blood cells. *Journal of Cancer Research and Clinical Oncology*. 2006, 132, 287–292.

- [14] Khandhar AP, Ferguson RM, Simon JA, Krishnan K.M. Tailored magnetic nanoparticles for optimizing magnetic fluid hyperthermia. *Journal of Biomedical Materials Research*. 2012, 100, 728–737.
- [15] Alexiou C, Jurgons R, Seliger C, Brunke O, Iro H, Odenbach S. Delivery of superparamagnetic nanoparticles for local chemotherapy after intraarterial infusion and magnetic drug targeting. *Anticancer Research*. 2007, 27, 2019–2022.
- [16] Reimer P, Tombach B. Hepatic MRI with SPIO: Detection and characterization of focal liver lesions. *European Journal of Radiology*. 1998, 8, 1198–1204.
- [17] Grubnic S, Padhani AR, Revell PB, Husband JE. Comparative efficacy of and sequence choice for two oral contrast agents used during MR imaging. *American Journal of Roentgenology*. 1999, 173, 173–178.
- [18] Mitchell DG. MR imaging contrast agents—What’s in a name?. *Journal of Magnetic Resonance Imaging*. 1997, 7, 1–4.
- [19] Okon E, Pouliquen D, Okon P, Kovaleva ZV, Stepanova TP, Lavit SG, Kudryavtsev BN, Jallet P. Biodegradation of magnetite dextran nanoparticles in the rat. *Avhistologic and biophysical study. Lab Invest*. 1994, 71, 895-903
- [20] Thorex DLJ, Chen AK, Czupryna J, Tsourkas A. Superparamagnetic iron oxide Nanoparticle Probes for Molecular Imaging. *Annals of Biomedical Engineering*. 2006, DOI: 10.1007/s10439-005-9002-7.
- [21] Shamsi K, Balzer T, Saini S, Ros PR, Nelson RC, Carter EC, Tollerfield S, Niendorf HP. Superparamagnetic iron oxide particles (SH U 555 A): Evaluation of efficacy in three doses for hepatic MR imaging. *Radiology* 1998, 206, 365–371.

- [22] Vogl TJ, Hammerstingl R, Schwarz W, Mack MG, Muller PK, Pegios W, Keck H, Eibesfeldt AE, Hoelzl J, Woessmer B, Bergman C, Felix R. Superparamagnetic iron oxide-enhanced versus gadolinium-enhanced MR imaging for differential diagnosis of focal liver lesions. *Radiology* 1996, 198, 881– 887.
- [23] Halestrap AP, Kerr PM, Javadov S, Woodfield KY. Elucidating the molecular mechanism of the permeability transition pore and its role in reperfusion injury of the heart. *Biochemica et Biophysica Acta*. 1998, 1366, 79-94.
- [24] Crompton M. The mitochondrial permeability transition pore and its role in cell death. *Biochemical Journal*. 1999, 341, 127-132.
- [25] Kroemer G, Dallaporta B, ReschRigon M. The mitochondrial death/life regulator in apoptosis and necrosis. *Annual Review of Physiology*. 1998, 60, 619-642.
- [26] Duchen MR. Contributions of mitochondria to animal physiology: from homeostatic sensor to calcium signaling and cell death. *Journal of Physiology*. 1999, 516, 1-17.
- [27] Chang SB. A molecular understanding of ATP-dependent solute transport by multidrug resistance-associated protein MRP1. *Cancer and Metastasis Reviews*. 2007, 26, 15-37.
- [28] Broxterman HJ, Pinedo HM. Energy metabolism in multidrug resistant tumor cells: a review. *J Cell Pharmacol* 1991, 2, 239 – 247.
- [29] Mankhetkorn S, Dubru F, Hesschenbrouck J, Fiallo M, Garnier-Suillerot A. Relation among the resistance factor, kinetics of uptake, and kinetics of the P-glycoprotein-mediated efflux of doxorubicin, daunorubicin, 8-(S)-fluoroidarubicin, and idarubicin in multidrug-resistant K562 cells. *Molecular Pharmacology*. 1996, 49, 532–539.

- [30] Miccadei S, Fanciulli M, Bruno T, Paggi MG, Floridi A. Energy metabolism of adriamycin-sensitive and -resistant Ehrlich ascites tumor cells. *Oncology Research*. 1996, 8, 27– 35.
- [31] Dorward A, Sweet S, Moorehead R, Singh G. Mitochondrial contributions to cancer cell physiology: redox balance, cell cycle, and drug resistance. *Journal of Bioenergetics and Biomembranes*. 1997, 29, 385– 392.
- [32] Johnson LV, Walsh ML, Chen LB. Localization of mitochondria in living cells with rhodamine 123. *Proceedings of the National Academy of Sciences of the United States of America*. 1980, 77, 990-994.
- [33] Johnson LV, Walsh ML, Bockus BJ, Chen LB. Monitoring of Relative Mitochondrial Membrane Potential in Living Cells by Fluorescence Microscopy. *The Journal of Cell Biology*. 1981, 88, 526-535.
- [34] Reungpatthanaphong P, Deschsupa S, Meesungnoen J, Loetchutinat C, Mankhetkorn S. Rhodamine B as a mitochondrial probe for measurement and monitoring of mitochondrial membrane potential in drug-sensitive and – resistant cells. *Journal of Biochemical and Biophysical Methods*. 2003, 57, 1-16.
- [35] Saritas EU, Goodwill PW, Croft LR, Konkle JJ, Lu K, Zheng B, Conolly SM. Magnetic particle imaging (MPI) for NMR and MRI researchers. *Journal of Magnetic Resonance*. 2013, 229, 116–126.
- [36] Borgert J, Schmidt JD, Schmale I, Rahmer J, Bontus C, Gleich B, David B, Eckart R, Woywode O, Weizenecker J. Fundamentals and applications of magnetic particle imaging. *Journal of Cardiovascular Computed Tomography*. 2012, 6, 149–153.
- [37] Azzam T, Raskin A, Makovitzki A, Brem H, Vierling P, Lineal M, Domb AJ. Cationic Polysaccharides for Gene Delivery. *Macromolecules*. 2002, 35, 9947-9953.

- [38] Molday RS. Magnetic iron-dextran microspheres, US Patent 4452773 A. 1984.
- [39] Josephson L, Gaw DA. Amine functionalized superparamagnetic nanoparticles for the synthesis of bioconjugates and uses therefor, US Patent 8420055 B2. 2013.
- [40] Bikram M. Ultrasmall superparamagnetic iron oxide nanoparticles and uses thereof, WO patent 2010093420 A2. 2010.
- [41] Bulte JW, Kraitchman DL. Iron oxide MR contrast agents for molecular and cellular imaging. *NMR in Biomedicine*. 2004, 17, 484–499.
- [42] Teeter LD, Becker FF, Chisari FV, Li D, Kuo MT. Overexpression of the Multidrug Resistance Gene *mdr3* in Spontaneous and Chemically Induced Mouse Hepatocellular Carcinomas. *Molecular and Cellular Biology*. 1990, 5728-5735.
- [43] Kuo MT, Zhao JY, Teeter LD, Ikeguchi M, Chisari FV. Multidrug Resistance (P-Glycoprotein) *mdr3/mdr1a* Gene during the Development of Hepatocellular Carcinoma in Hepatitis B Virus Transgenic Mice. *Cell Growth & Differentiation*. 1992, 3, 531-540.