REFERENCES

- Asahara T, "Isolation of putative progenitor endothelial cells for angiogenesis," Science, Vol. 275 No. 5302, 1997, pp. 964-967
- [2] Shang H, "Differentiation of human peripheral blood mononuclear cells into hepatocyte-like cells in vitro," Journal of Clinical Rehabilitative Tissue Engineering Research, Vol. 13 No. 10, 2009, pp. 1859-1863
- [3] Badorff C, "Transdifferentiation of blood-derived human adult endothelial progenitor cells into functionally active cardiomyocytes," Circulation, Vol. 107 No.7, 2003, pp. 1024-1032
- [4] Simper D, "Smooth muscle progenitor cells in human blood," Circulation, Vol. 106 No. 10, 2002, pp. 1199-1204
- [5] Hu G, "Supernatant of bone marrow mesenchymal stromal cells induces peripheral blood mononuclear cells possessing mesenchymal features," International Journal of Biological Sciences, Vol. 7 No. 3, 2011, pp. 364-375
- [6] Wan C, "Allogenic peripheral blood derived mesenchymal stem cells (MSCs) enhance bone regeneration in rabbit ulna critical-sized bone defect model," Journal of Orthopaedic Research, Vol. 24 No. 4, 2006, pp. 610-618
- [7] Dellambra E, "Downregulation of 14-3-3sigma prevents clonal evolution and leads to immortalization of primary human keratinocytes," Journal of Cell Biology, Vol. 149 No. 5, 2000, pp. 1117-1130

[8] Horschitz S, "Generation of neuronal cells from human peripheral blood mononuclear cells," Neuroreport, Vol. 21 No. 3, 2010, pp. 185-190

- [9] Fahey TR, "Cytokine production in a model of wound healing: the appearance of MIP-1, MIP-2, cachectin/TNF and IL-1," Cytokine, Vol. 2 No. 2, 1990, pp. 92-99
- [10] J. Yu, "The predictive value of white cell or CD34+ cell count in the peripheral blood for timing apheresis and maximizing yield," Transplantation, Vol. 39 No.5, 2002, pp. 442-450
- [11] Anderson DJ, "Can stem cells cross lineage boundaries?," Nature Medicine, Vol.7 No. 4, 2001, pp. 393-395
- [12] Ruzanna Ab Kadir, "Molecular characterisation of human peripheral blood stem cells," South African Journal of Science, Vol. 108 No.939, 2012, pp. 1-7
- [13] Bernard Edward Tuch, "Stem cells A clinical update," Clinical Practice Update from Australian Family Physician, Vol. 35 No. 9, 2006, pp.719-21
- [14] A. J. Becker, E. A, "Cytological demonstration of the clonal nature of spleen colonies derived from transplanted mouse marrow cells," Nature, Vol. 197 No.4866, 1963, pp. 452–454
- [15] D. T. Scadden, "The stem-cell niche as an entity of action," Nature, Vol. 441 No. 7097, 2006, pp. 1075–1079
- [16] M. F. Pittenger, "Multilineage potential of adult human mesenchymal stem cells," Science, Vol. 284 No. 5411, 1999, pp. 143–147
- [17] H. Nakahara, V. M, "Cultureexpanded human periosteal-derived cells exhibit osteochondral potential in vivo," Journal of Orthopaedic Research, Vol. 9 No. 4, 1991 pp. 465–476
- [18] P. A. Zuk, "Human adipose tissue is a source of multipotent stem cells," Molecular Biology of the Cell, Vol. 13 No. 12, 2002, pp. 4279–4295

- [19] J. T. Williams, S, "Cells isolated from adult human skeletal muscle capable of differentiating into multiple mesodermal phenotypes," American Surgeon, Vol. 65 No. 1, 1999, pp. 22–26
- [20] C. De Bari, "Multipotent mesenchymal stem cells from adult human synovial membrane," Arthritis & Rheumatism, Vol. 44 No. 8, 2001, pp. 1928–1942
- [21] O.K.Lee, "Isolation of multipotent mesenchymal stem cells from umbilical cord blood," Blood, Vol. 103 No. 5, 2004, pp. 1669–1675
- [22] M. Secco, "Multipotent stem cells from umbilical cord: cord is richer than blood!," Stem Cells, Vol. 26 No. 1, 2008, pp. 146–150
- [23] Y. Fukuchi, "Human placenta-derived cells have mesenchymal stem/progenitor cell potential," Stem Cells, Vol.22 No. 5, 2004, pp. 649–658
- [24] M. Krampera, "Mesenchymal stem cells: from biology to clinical use," Blood Transfusion, Vol. 5 No. 3, 2007, pp. 120–129
- [25] Wolfgang Wagner, "Aging of hematopoietic stem cells is regulated by the stem cell niche," Experimental Gerontology, Vol. 43 No. 11, 2008, pp. 974–980
- [26] C. Zhang et al, "Mobilization of peripheral blood stem cells for autologous transplantation patients with hematological malignancies: Influence of disease, mobilization method, age and sex," Transfusion and Apheresis Science, Vol. 39 No. 1, 2008, pp.21–28
- [27] Georg Stussi, "ABO-histo blood group incompatibility in hematopoietic stem cell and solid organ transplantation," Transfusion and Apheresis Science, Vol. 35 No. 1, 2006 pp.59–69

- [28] Mats Remberger, "Major ABO Blood Group Mismatch Increases the Risk for Graft Failure after Unrelated Donor Hematopoietic Stem Cell Transplantation," Biology of Blood and Marrow Transplantation, Vol. 13 No. 6, 2007, pp.675-682
- [29] Isabel Galan, "Effect of the ABO blood group on the proliferative and clonogenic capacity of umbilical cord stem cells," Transfusion and Apheresis Science, Vol. 35 No. 2, 2006, pp.119–123
- [30] Metallo CM, "Engineering the stem cell microenvironment," Biotechnology Progress, Vol. 23 No. 1, 2007, pp.18–23
- [31] R. Peerani, "Enabling stem cell therapies through synthetic stem cell-niche engineering," The Journal of Clinical Investigation, Vol. 120 No. 1, 2010, pp.60-70
- [32] Stephen F, "The extracellular matrix as a biologic scaffold material," Biomaterials, Vol. 28 No. 25, 2007, pp. 3587-3593
- [33] Xiao hua Liu, "Porogen-induced surface modification of nano-fibrous poly(L-lactic acid) scaffolds for tissue engineering," Biomaterials, Vol.27 No.21, 2006, pp. 3980-3987
- [34] Mannello F, "Concise review: No breakthroughs for human mesenchymal and embryonic stem cell culture: conditioned medium, feeder layer, or feeder-free; medium with fetal calf serum, human serum, or enriched plasma; serum-free, serum replacement nonconditioned medium, or ad hoc formula? All that glitters is not gold!," Stem Cells, Vol.25 No. 7, 2007, pp. 1603-9
- [35] Honn KV, "Fetal bovine serum: a multivariate standard," Proceedings of the Society for Experimental Biology and Medicine, Vol. 149 No. 2, 1975, pp. 344-7

reserv

- [36] Karen Bieback, "Replicative aging and differentiation potential of human adipose tissue-derived mesenchymal stromal cells expanded in pooled human or fetal bovine serum," Cytotherapy, Vol. 14 No. 5, 2012, pp. 570-583
- [37] Doyonnas R, "Hematopoietic contribution to skeletal muscle regeneration by myelomonocytic precursors," Proceedings of the National Academy of Sciences of the united states of America, Vol. 101 No. 37, 2004, pp. 13507-13512
- [38] Orlic D, "Mobilized bone marrow cells repair the infarcted heart, improving function and survival," Proceedings of the National Academy of Sciences of the united states of America, Vol. 98 No. 18, 2001, pp. 10344-10349
- [39] Sigur jonsson O, "Adult human hematopoietic stem cells produce neurons efficiently in the regenerating chicken embryo spinal cord," Proceedings of the National Academy of Sciences of the united states of America, Vol. 102 No. 14, 2005, pp. 5227-5232
- [40] Susanne Grassel, "Influence of cellular microenvironment and paracrine signals on chondrogenic differentiation," Bioscience, Vol. 12 No. 1, 2007, pp. 4946-4956

[41] Clifford D.L Folmes, "Metabolic Plasticity in Stem Cell Homeostasis and Differentiation," Cell Stem Cell, Vol. 11 No. 5, 2012, pp. 596-606
Copyright^O by Chiang Mai University
A I i g h t s reserved