

## REFERENCES

- Achouri, A., Boye, J. I. and Belanger, D. (2005). Soybean isoflavones: efficacy of extraction conditions and effect of food type on extractability. *Food Research International*, 38(10), 1199-1204.
- Albertazzi, P., Pansini, F., Bonaccorsi, G., Zanotti, L., Forini, E. and De Aloysio, D. (1998). The effect of dietary soy supplementation on hot flushes. *Obstetrics & Gynecology*, 91(1), 129-135.
- Altschul, S. F., Madden, T. L., Schäffer, A. A., Zhang, J., Zhang, Z., Miller, W. and Lipman, D. J. (1997). Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Research*, 25(17), 3389-3402.
- American Heart Association. (2007). AHA statistical update: heart disease and stroke statistics. *Circulation*, 115, e69-e171.
- Anderson, J. J. B. and Garner, S. C. (1997). The effects of phytoestrogens on bone. *Nutrition Research*, 17(10), 1617-1632.
- Anderson, J. W., Johnstone, B. M. and Cook-Newell, M. E. (1995). Meta-analysis of the effects of soy protein intake on serum lipids. *New England Journal of Medicine*, 333(5), 276-282.
- Anthony, M. S., Clarkson, T. B., Hughes Jr, C. L., Morgan, T. M. and Burke, G. L. (1996). Soybean isoflavones improve cardiovascular risk factors without affecting the reproductive system of peripubertal rhesus monkeys. *Journal of Nutrition*, 126(1), 43-50.

Bae, S.-M., Jang, C. H., Kim, J.-H., Lim, H. A., Kim, J.-R., Kim, J. H. and Kim, J.-S. (2005). Optimization of isoflavone extraction from soy germ. *Journal of Food Science and Nutrition*, 10(3), 290-293.

Bhandari, B. R., Dumoulin, E. D., Richard, H. M. J., Noleau, I. and Lebert, A. M. (1992). Flavor encapsulation by spray drying: application to citral and linalyl acetate. *Journal of food science*, 57(1), 217-221.

Bingham, S., Atkinson, C., Liggins, J., Bluck, L. and Coward, A. (1998). Phytoestrogens: where are we now?. *British Journal of Nutrition*, 79(05), 393-406.

Brosius, J., Dull, T. J., Sleeter, D. D. and Noller, H. F. (1981). Gene organization and primary structure of a ribosomal RNA operon from *Escherichia coli*. *Journal of Molecular Biology*, 148(2), 107-127.

Cano-Chauca, M., Stringheta, P. C., Ramos, A. M. and Cal-Vidal, J. (2005). Effect of the carriers on the microstructure of mango powder obtained by spray drying and its functional characterization. *Innovative Food Science & Emerging Technologies*, 6(4), 420-428.

Cao, Q. H., Qu, W. J., Li, J. G. and Deng, Y. X. (2004). Study on purification of total flavonoids from seed residue of Hippophae rhamnoides with macroporous resin. *China Journal of Chinese Materia Medica*, 29(3), 225-228.

Chang, L.-H. and Chang, C.-M. J. (2007). Continuous hot pressurized fluids extraction of isoflavones and soyasaponins from defatted soybean flakes. *Journal of the Chinese Institute of Chemical Engineers*, 38(3-4), 313-319.

Chang, L.-H., Cheng, Y.-C. and Chang, C.-M. (2004). Extracting and purifying isoflavones from defatted soybean flakes using superheated water at elevated pressures. *Food Chemistry*, 84(2), 279-285.

- Chien, J. T., Hsieh, H. C., Kao, T. H. and Chen, B. H. (2005). Kinetic model for studying the conversion and degradation of isoflavones during heating. *Food Chemistry*, 91(3), 425-434.
- Cho, S. Y., Lee, Y. N. and Park, H. J. (2009). Optimization of ethanol extraction and further purification of isoflavones from soybean sprout cotyledon. *Food Chemistry*, 117(2), 312-317.
- Choi, Y.-B., Kim, K.-S. and Rhee, J.-S. (2002). Hydrolysis of soybean isoflavone glucosides by lactic acid bacteria. *Biotechnology Letters*, 24(24), 2113-2116.
- Chukeatirote, E., Chainun, C., Siengsubchart, A., Moukamnerd, C., Chantawannakul, P., Lumyong, S., Boontim, N. and Thakang, P. (2006). Microbiological and biochemical changes in *Thua nao* fermentation. *Research Journal of Microbiology*, 1(1), 38-44.
- Claus, D. and Berkeley, R. C. W. (1986). The genus *Bacillus*. In P. H. A. Sneath (Ed.), *Bergey's Manual of Systematic Bacteriology* (Vol. 2, pp. 1105-1139). Baltimore, USA: Williams and Wilkins.
- Couto, S. R. and Sanromán, M. A. (2006). Application of solid-state fermentation to food industry—a review. *Journal of Food Engineering*, 76(3), 291-302.
- Crouzet, J. and Chassagne, D. (1999). Glycosidically bound volatiles in plants. In R. Ikan (Ed.), *Naturally Occuring Glycosides* (pp. 225-274). Canada: John Wiely & Sons.
- Dajanta, K. (2010). *Production of High Nutritional Fermented Soybean (Thua nao) by Bacillus subtilis*. (Ph.D. Thesis in Food Science and Technology), Chiang Mai University, Chiang Mai, Thailand.

- Dajanta, K., Chukeatirote, E., Apichartsrangkoon, A. and Frazier, R. A. (2009). Enhanced aglycone production of fermented soybean products by *Bacillus* species. *Acta Biologica Szegediensis*, 53(2), 93-98.
- Dakwa, S., Sakyi-Dawson, E., Diako, C., Annan, N. T. and Amoa-Awua, W. K. (2005). Effect of boiling and roasting on the fermentation of soybeans into dawadawa (soy-dawadawa). *International Journal of Food Microbiology*, 104(1), 69-82.
- Dalais, F., Rice, G., Wahlqvist, M., Greher, M., Murkies, A., Medley, G., Ayton, R. and Strauss, B. (1998). Effects of dietary phytoestrogens in postmenopausal women. *Climacteric*, 1(2), 124-129.
- Di Mauro, A., Arena, E., Fallico, B., Passerini, A. and Maccarone, E. (2002). Recovery of anthocyanins from pulp wash of pigmented oranges by concentration on resins. *Journal of Agricultural and Food Chemistry*, 50(21), 5968-5974.
- Dike, E. and Odunfa, S. (2003). Microbiological and biochemical evaluation of a fermented soyabean product: soyadawadawa. *Journal of Food Science and Technology*, 40(6), 606-610.
- dos Santos, M. M., da Rosa, A. S., Dal'Boit, S., Mitchell, D. A. and Krieger, N. (2004). Thermal denaturation: is solid-state fermentation really a good technology for the production of enzymes?. *Bioresource Technology*, 93(3), 261-268.
- Eden, J. (1998). 3 Phytoestrogens and the menopause. *Baillière's Clinical Endocrinology and Metabolism*, 12(4), 581-587.
- Eldridge, A. C. and Kwolek, W. F. (1983). Soybean isoflavones: Effect of environment and variety on composition. *Journal of Agricultural and Food Chemistry*, 31(2), 394-396.

Ellis, M. J. (1994). The methodology of shelf life determination. In C. M. D. Man & A. A. Jones (Eds.), *Shelf Life Evaluation of Foods* (pp. 27-39): Springer US.

Felsenstein, J. (1985). Confidence limits on phylogenies: an approach using the bootstrap. *Evolution*, 783-791.

Fernández-Pérez, V., Tapiador, J., Martín, A. and de Castro, M. L. (2004). Optimization of the drying step for preparing a new commercial powdered soup. *Innovative Food Science & Emerging Technologies*, 5(3), 361-368.

Feron, G., Bonnarme, P. and Durand, A. (1996). Prospects for the microbial production of food flavours. *Trends in Food Science & Technology*, 7(9), 285-293.

Fu, B., Liu, J., Li, H., Li, L., Lee, F.S.C. and Wang, X. (2005). The application of macroporous resins in the separation of licorice flavonoids and glycyrrhizic acid. *Journal of Chromatography A*, 1089(1-2), 18-24.

Genovese, M. I. and Lajolo, F. M. (2001). Determinação de isoflavonas em derivados de soja. *Ciência e Tecnologia de Alimentos*, 21(1), 86-93.

Giannakourou, M. C. and Taoukis, P. S. (2003). Kinetic modelling of vitamin C loss in frozen green vegetables under variable storage conditions. *Food Chemistry*, 83(1), 33-41.

Gordon, R. E. (1981). One hundred and seven years of the genus *Bacillus*. In R. C. Berkeley & M. Goodfellow (Eds.), *The Aerobic Endosporeforming Bacteria*. London: Academic Press.

Griffith, A. P. and Collison, M. W. (2001). Improved methods for the extraction and analysis of isoflavones from soy-containing foods and nutritional supplements by reversed-phase high-performance liquid chromatography and liquid

chromatography–mass spectrometry. *Journal of Chromatography A*, 913(1–2), 397-413.

Grün, I. U., Adhikari, K., Li, C., Li, Y., Lin, B., Zhang, J. and Fernando, L. N. (2001). Changes in the profile of genistein, daidzein, and their conjugates during thermal processing of Tofu. *Journal of Agricultural and Food Chemistry*, 49(6), 2839-2843.

Guo, H.-D., Zhang, Q.-F., Chen, J.-G., Shangguang, X.-C. and Guo, Y.-X. (2015). Large scale purification of puerarin from Puerariae Lobatae Radix through resins adsorption and acid hydrolysis. *Journal of Chromatography B*, 980(0), 8-15.

Hall, T. A. (1999). BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series*, 41, 95-98

Han, K. K., Soares Jr, J. M., Haidar, M. A., Rodrigues de Lima, G. and Baracat, E. C. (2002). Benefits of soy isoflavone therapeutic regimen on menopausal symptoms. *Obstetrics & Gynecology*, 99(3), 389-394.

Hargreaves, D. F., Potten, C. S., Harding, C., Shaw, L. E., Morton, M. S., Roberts, S. A., Howell, A. and Bundred, N. J. (1999). Two-week dietary soy supplementation has an estrogenic effect on normal premenopausal breast 1. *The Journal of Clinical Endocrinology & Metabolism*, 84(11), 4017-4024.

Hesseltine, C. (1977). Solid-state fermentation. 1. *Process Biochemistry*, 12(6), 24-27.

Hoeck, J. A., Fehr, W. R., Murphy, P. A. and Welke, G. A. (2000). Influence of genotype and environment on isoflavone contents of soybean. *Crop Science*, 40(1), 48-51.

Hölker, U. and Lenz, J. (2005). Solid-state fermentation—are there any biotechnological advantages?. *Current Opinion in Microbiology*, 8(3), 301-306.

Hutabarat, L. S., Greenfield, H. and Mulholland, M. (2001). Isoflavones and coumestrol in soybeans and soybean products from Australia and Indonesia. *Journal of Food Composition and Analysis*, 14(1), 43-58.

Ibe, S., Kumada, K., Yoshiha, M. and Onga, T. (2001). Production of natto which contains a high level of isoflavone aglycons. *Journal-Japanese Society of Food Science and Technology*, 48(1), 27-34.

Ikeda, G., Tomizawa, A., Imayoshi, Y., Iwabuchi, H., Hinata, T. and Sagara, Y. (2006). Flavor design of sesame-flavored dressing using gas chromatography/olfactometry and food kansei model. *Food Science and Technology Research*, 12(4), 261-269.

Ishimi, Y., Yoshida, M., Wakimoto, S., Wu, J., Chiba, H., Wang, X., Takeda, K. and Miyaura, C. (2002). Genistein, a soybean isoflavone, affects bone marrow lymphopoiesis and prevents bone loss in castrated male mice. *Bone*, 31(1), 180-185.

Izumi, T., Piskula, M. K., Osawa, S., Obata, A., Tobe, K., Saito, M., Kataoka, S., Kubota, Y. and Kikuchi, M. (2000). Soy isoflavone aglycones are absorbed faster and in higher amounts than their glucosides in humans. *The Journal of Nutrition*, 130(7), 1695-1699.

Jackson, C.-J., Dini, J., Lavandier, C., Rupasinghe, H., Faulkner, H., Poysa, V., Buzzell, D. and DeGrandis, S. (2002). Effects of processing on the content and composition of isoflavones during manufacturing of soy beverage and tofu. *Process Biochemistry*, 37(10), 1117-1123.

Jankowiak, L., Kantzas, N., Boom, R. and van der Goot, A. J. (2014). Isoflavone extraction from okara using water as extractant. *Food Chemistry*, 160(0), 371-378.

Jaya, S. and Das, H. (2004). Effect of maltodextrin, glycerol monostearate and tricalcium phosphate on vacuum dried mango powder properties. *Journal of Food Engineering*, 63(2), 125-134.

Jeong, D.-W., Kim, H.-R., Jung, G., Han, S., Kim, C.-T. and Lee, J.-H. (2014). Bacterial community migration in the ripening of doenjang, a traditional korean fermented soybean food. *Journal of Microbiology and Biotechnology*, 24(5), 648-660.

Jideani, I. and Okeke, C. (1991). Comparative study of microorganisms and sensory attributes of condiments from the fermentation of different seeds. *Plant Foods for Human Nutrition*, 41(1), 27-34.

Johns, M. R. and Stuart, D. M. (1991). Production of pigments by *Monascus purpureus* in solid culture. *Journal of Industrial Microbiology*, 8(1), 23-28.

Jung, K.-O., Park, S.-Y. and Park, K.-Y. (2006). Longer aging time increases the anticancer and antimetastatic properties of doenjang. *Nutrition*, 22(5), 539-545.

Kao, T. H. and Chen, B. H. (2007). Effects of different carriers on the production of isoflavone powder from soybean cake. *Molecules*, 12(4), 917-931.

Kao, T. H., Lu, Y. F., Hsieh, H. C. and Chen, B. H. (2004). Stability of isoflavone glucosides during processing of soymilk and tofu. *Food Research International*, 37(9), 891-900.

Katsura, K., Kawasaki, H., Potacharoen, W., Saono, S., Seki, T., Yamada, Y., Uchimura, T. and Komagata, K. (2001). *Asaia siamensis* sp. nov., an acetic acid bacterium in the alpha-proteobacteria. *International Journal of Systematic and Evolutionary Microbiology*, 51(2), 559-563.

Kawasaki, H., Hoshino, Y., Hirata, A. and Yamasato, K. (1993). Is intracytoplasmic membrane structure a generic criterion? It does not coincide with phylogenetic interrelationships among phototrophic purple nonsulfur bacteria. *Archives of Microbiology*, 160(5), 358-362.

Kim, J. J., Kim, S. H., Hahn, S. J. and Chung, I. M. (2005). Changing soybean isoflavone composition and concentrations under two different storage conditions over three years. *Food Research International*, 38(4), 435-444.

Kim, O.-S., Cho, Y.-J., Lee, K., Yoon, S.-H., Kim, M., Na, H., Park, S.-C., Jeon, Y. S., Lee, J.-H., Yi, H., Won, S. and Chun, J. (2012). Introducing EzTaxon-e: a prokaryotic 16S rRNA gene sequence database with phylotypes that represent uncultured species. *International Journal of Systematic and Evolutionary Microbiology*, 62(3), 716-721.

Kiuchi, K. and Watanabe, S. (2004). Industrialization of Japanese natto. In K. H. Steinkraus (Ed.), *Industrialization of Indigenous Fermented Foods* (2<sup>nd</sup> edition ed., pp. 193-246). New York: Marcel Dekker.

Klejdus, B., Mikelová, R., Petrlová, J., Potěšil, D., Adam, V., Stiborová, M., Hodek, P., Vacek, J., Kizek, R. and Kubáň, V. (2005). Determination of isoflavones in soy bits by fast column high-performance liquid chromatography coupled with UV-visible diode-array detection. *Journal of Chromatography A*, 1084(1-2), 71-79.

Klump, S. P., Allred, M. C., MacDonald, J. L. and Ballam, J. M. (2001). Determination of isoflavones in soy and selected foods containing soy by extraction, saponification, and liquid chromatography: collaborative study. *Journal of AOAC International*, 84(6), 1865-1883.

Knight, D., Wall, P. L. and Eden, J. (1996). review of phytoestrogens and their effects in relation to menopausal symptoms. *Australian Journal of Nutrition and Dietetics*.

Krishnan, S., Bhosale, R. and Singhal, R. S. (2005). Microencapsulation of cardamom oleoresin: evaluation of blends of gum arabic, maltodextrin and a modified starch as wall materials. *Carbohydrate Polymers*, 61(1), 95-102.

Kudou, S., Fleury, Y., Welti, D., Magnolato, D., Uchida, T., Kitamura, K. and Okubo, K. (1991). Malonyl isoflavone glycosides in soybean seeds (*Glycine max* Merrill). *Agricultural and Biological Chemistry*, 55(9), 2227-2233.

Kuiper, G. G., Carlsson, B., Grandien, K., Enmark, E., Hagblad, J., Nilsson, S. and Gustafsson, J.-a. (1997). Comparison of the ligand binding specificity and transcript tissue distribution of estrogen receptors  $\alpha$  and  $\beta$ . *Endocrinology*, 138(3), 863-870.

Kuo, L.-C., Cheng, W.-Y., Wu, R.-Y., Huang, C.-J. and Lee, K.-T. (2006). Hydrolysis of black soybean isoflavone glycosides by *Bacillus subtilis* natto. *Applied Microbiology and Biotechnology*, 73(2), 314-320.

Kwak, C. S., Lee, M. S. and Park, S. C. (2007). Higher antioxidant properties of Chungkookjang, a fermented soybean paste, may be due to increased aglycone and malonylglycoside isoflavone during fermentation. *Nutrition Research*, 27(11), 719-727.

Lee, J. H., Renita, M., Fioritto, R. J., St. Martin, S. K., Schwartz, S. J. and Vodovotz, Y. (2004). Isoflavone characterization and antioxidant activity of Ohio soybeans. *Journal of Agricultural and Food Chemistry*, 52(9), 2647-2651.

Lee, S. J., Ahn, J. K., Kim, S. H., Kim, J. T., Han, S. J., Jung, M. Y. and Chung, I. M. (2003a). Variation in isoflavone of soybean cultivars with location and storage duration. *Journal of Agricultural and Food Chemistry*, 51(11), 3382-3389.

Lee, S. J., Yan, W., Ahn, J. K. and Chung, I. M. (2003b). Effects of year, site, genotype and their interactions on various soybean isoflavones. *Field Crops Research*, 81(2), 181-192.

Lee, Y.-W., Kim, J.-D., Zheng, J. and Row, K. H. (2007). Comparisons of isoflavones from Korean and Chinese soybean and processed products. *Biochemical Engineering Journal*, 36(1), 49-53.

Leejeerajumnean, A. (2003). Thua nao: alkali fermented soybean from *Bacillus subtilis*. *Silpakorn University International Journal*, 3, 277-292.

Lekhakula, P. (2010). *Process Development in Production of Thua-nao Using Selected Mixed Cultures*. (Master of Science), Chiang Mai University, Grauate School, Chiang Mai, Thailand (In Thai).

Li, A., Zhang, Q., Chen, J., Fei, Z., Long, C. and Li, W. (2001). Adsorption of phenolic compounds on Amberlite XAD-4 and its acetylated derivative MX-4. *Reactive and Functional Polymers*, 49(3), 225-233.

Li, H., Liu, J., Li, D. and Wang, H. (2011). Study on separation and purification of genistein in the soybean residue using macroporous resin adsorption. *Industrial & Engineering Chemistry Research*, 51(1), 44-49.

Lin, F. and Giusti, M. M. (2005). Effects of solvent polarity and acidity on the extraction efficiency of isoflavones from soybeans (*Glycine max*). *Journal of Agricultural and Food Chemistry*, 53(10), 3795-3800.

Liu, D., Zhen, W., Yang, Z., Carter, J. D., Si, H. and Reynolds, K. A. (2006). Genistein acutely stimulates insulin secretion in pancreatic  $\beta$ -cells through a cAMP-dependent protein kinase pathway. *Diabetes*, 55(4), 1043-1050.

Liu, K. (2004). *Soy Isoflavones: Chemistry, Processing Effects, Health Benefits, and Commercial Production*. Champaign, IL: AOCS Press.

Liu, K. (2012). *Soybeans: Chemistry, Technology, and Utilization*: Springer.

Liu, Y., Di, D., Bai, Q., Li, J., Chen, Z., Lou, S. and Ye, H. (2011). Preparative separation and purification of rebaudioside a from steviol glycosides using mixed-mode macroporous adsorption resins. *Journal of Agricultural and Food Chemistry*, 59(17), 9629-9636.

Logan, N. A. and Vos, P. D. (2009). Genus I. *Bacillus* Cohn 1872. In P. D. Vos, G. Garrity, D. Jones, N. R. Krieg, W. Ludwig, F. A. Rainey, K. H. Schleifer & W. Whitman (Eds.), *Bergey's Manual of Systematic Bacteriology: Volume 3: The Firmicutes* (pp. 21-127): Springer New York.

Mackey, R. and Eden, J. (1998). Phytoestrogens and the menopause. *Climacteric*, 1(4), 302-308.

Micke, G. A., Fujiya, N. M., Tonin, F. G., de Oliveira Costa, A. C. and Tavares, M. F. M. (2006). Method development and validation for isoflavones in soy germ pharmaceutical capsules using micellar electrokinetic chromatography. *Journal of Pharmaceutical and Biomedical Analysis*, 41(5), 1625-1632.

Murphy, P. A., Barua, K. and Hauck, C. C. (2002). Solvent extraction selection in the determination of isoflavones in soy foods. *Journal of Chromatography B*, 777(1-2), 129-138.

Murphy, P. A., Song, T., Buseman, G., Barua, K., Beecher, G. R., Trainer, D. and Holden, J. (1999). Isoflavones in retail and institutional soy foods. *Journal of Agricultural and Food Chemistry*, 47(7), 2697-2704.

Myers, E. W. and Miller, W. (1988). Optimal alignments in linear space. *Computer Applications in the Biosciences : CABIOS*, 4(1), 11-17.

Nguyenle, T., Wang, E. and Cheung, A. P. (1995). An investigation on the extraction and concentration of isoflavones in soy-based products. *Journal of Pharmaceutical and Biomedical Analysis*, 14(1–2), 221-232.

Nishio, K., Niwa, Y., Toyoshima, H., Tamakoshi, K., Kondo, T., Yatsuya, H., Yamamoto, A., Suzuki, S., Tokudome, S. and Lin, Y. (2007). Consumption of soy foods and the risk of breast cancer: findings from the Japan Collaborative Cohort (JACC) Study. *Cancer Causes & Control*, 18(8), 801-808.

Norris, J., Berkeley, R., Logan, N. and O'donnell, A. (1981). The genera *Bacillus* and *Sporolactobacillus*. In M. P. Starr, H. G. T. Stolp, A. Balows & H. G. Schlegle (Eds.), *The Prokaryotes* (Vol. 2, pp. 1711-1755). New York: Springer-Verlag.

Nout, M. J. R., Bakshi, D. and Sarkar, P. K. (1998). Microbiological safety of kinema, a fermented soya bean food. *Food Control*, 9(6), 357-362.

Nuin, M., Alfaro, B., Cruz, Z., Argarate, N., George, S., Le Marc, Y., Olley, J. and Pin, C. (2008). Modelling spoilage of fresh turbot and evaluation of a time-temperature integrator (TTI) label under fluctuating temperature. *International Journal of Food Microbiology*, 127(3), 193-199.

Ogbadu, L. and Okagbue, R. (1988). Bacterial fermentation of soya bean for 'daddawa'production. *Journal of Applied Bacteriology*, 65(5), 353-356.

Okamoto, A., Hanagata, H., Kawamura, Y. and Yanagida, F. (1995). Anti-hypertensive substances in fermented soybean, natto. *Plant Foods for Human Nutrition*, 47(1), 39-47.

Omafuvbe, B., Shonukan, O. and Abiose, S. (2000). Microbiological and biochemical changes in the traditional fermentation of soybean for ‘soy-daddawa’—Nigerian food condiment. *Food Microbiology*, 17(5), 469-474.

Pananun, T., Montalbo-Lomboy, M., Noomhorm, A., Grewell, D. and Lamsal, B. (2012). High-power ultrasonication-assisted extraction of soybean isoflavones and effect of toasting. *LWT - Food Science and Technology*, 47(1), 199-207.

Pandey, A. (2003). Solid-state fermentation. *Biochemical Engineering Journal*, 13(2), 81-84.

Pandey, A., Selvakumar, P., Soccol, C. R. and Nigam, P. (1999). Solid state fermentation for the production of industrial enzymes. *Current Science*, 77(1), 149-162.

Pandey, A., Soccol, C. R. and Mitchell, D. (2000). New developments in solid state fermentation: I-bioprocesses and products. *Process Biochemistry*, 35(10), 1153-1169.

Park, K.-Y., Jung, K.-O., Rhee, S.-H. and Choi, Y. H. (2003). Antimutagenic effects of doenjang (Korean fermented soypaste) and its active compounds. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis*, 523, 43-53.

Pongsatha, S. (2011). Phytoestrogen from  
[http://www.med.cmu.ac.th/dept/obgyn/2011/index.php?option=com\\_content&view=article&id=210:phytoestrogen&catid=37&Itemid=242](http://www.med.cmu.ac.th/dept/obgyn/2011/index.php?option=com_content&view=article&id=210:phytoestrogen&catid=37&Itemid=242)

Pongsawatmanit, R. (2009). Shelf life of product. In R. Pongsawatmanit (Ed.), *Product Development in Agro-Industry* (pp. 116-128). Bangkok, Thailand: Kasetsart University Press (*In Thai*).

- Potter, S. M., Baum, J. A., Teng, H., Stillman, R. J., Shay, N. F. and Erdman, J. (1998). Soy protein and isoflavones: their effects on blood lipids and bone density in postmenopausal women. *The American Journal of Clinical Nutrition*, 68(6), 1375S-1379S.
- Quella, S. K., Loprinzi, C. L., Barton, D. L., Knost, J. A., Sloan, J. A., LaVasseur, B. I., Swan, D., Krupp, K. R., Miller, K. D. and Novotny, P. J. (2000). Evaluation of soy phytoestrogens for the treatment of hot flashes in breast cancer survivors: a North Central Cancer Treatment Group Trial. *Journal of Clinical Oncology*, 18(5), 1068-1068.
- Raghavarao, K., Ranganathan, T. and Karanth, N. (2003). Some engineering aspects of solid-state fermentation. *Biochemical Engineering Journal*, 13(2), 127-135.
- Raimbault, M. (1998). General and microbiological aspects of solid substrate fermentation. *Electronic Journal of Biotechnology*, 1(3), 26-27.
- Rostagno, M. A., Araújo, J. M. A. and Sandi, D. (2002). Supercritical fluid extraction of isoflavones from soybean flour. *Food Chemistry*, 78(1), 111-117.
- Rostagno, M. A., Palma, M. and Barroso, C. G. (2004). Pressurized liquid extraction of isoflavones from soybeans. *Analytica Chimica Acta*, 522(2), 169-177.
- Rostagno, M. A., Villares, A., Guillamón, E., García-Lafuente, A. and Martínez, J. A. (2009). Sample preparation for the analysis of isoflavones from soybeans and soy foods. *Journal of Chromatography A*, 1216(1), 2-29.
- Roy, H., Lundy, S. and Kalicki, B. (2009). Soy Isoflavones. *Pennington Nutrition Series, number 9, healthier lives through education in nutrition and preventive medicine*. from <http://www.pbrc.edu>

Rungsardthong, V. (2002). *Food Processing Technology* (2<sup>nd</sup> ed.). Bangkok, Thailand: Text and Journal Publication [*In Thai*].

Saitou, N. and Nei, M. (1987). The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Molecular Biology and Evolution*, 4(4), 406-425.

Sandhu, A. K. and Gu, L. (2013). Adsorption/desorption characteristics and separation of anthocyanins from muscadine (*Vitis rotundifolia*) juice pomace by use of macroporous adsorbent resins. *Journal of Agricultural and Food Chemistry*, 61(7), 1441-1448.

Sarkar, P., Hasenack, B. and Nout, M. (2002). Diversity and functionality of *Bacillus* and related genera isolated from spontaneously fermented soybeans (Indian Kinema) and locust beans (African Soumbala). *International Journal of Food Microbiology*, 77(3), 175-186.

Sarkar, P., Tamang, J., Cook, P. and Owens, J. (1994). Kinema—a traditional soybean fermented food: proximate composition and microflora. *Food Microbiology*, 11(1), 47-55.

Scordino, M., Di Mauro, A., Passerini, A. and Maccarone, E. (2003). Adsorption of flavonoids on resins: hesperidin. *Journal of Agricultural and Food Chemistry*, 51(24), 6998-7004.

Scordino, M., Di Mauro, A., Passerini, A. and Maccarone, E. (2004). Adsorption of flavonoids on resins: cyanidin 3-glucoside. *Journal of Agricultural and Food Chemistry*, 52(7), 1965-1972.

Setchell, K. (1998). Phytoestrogens: the biochemistry, physiology, and implications for human health of soy isoflavones. *The American Journal of Clinical Nutrition*, 68(6), 1333S-1346S.

Setchell, K. D. (2000). Absorption and metabolism of soy isoflavones—from food to dietary supplements and adults to infants. *The Journal of Nutrition*, 130(3), 654S-655S.

Setchell, K. D. and Cassidy, A. (1999). Dietary isoflavones: biological effects and relevance to human health. *The Journal of Nutrition*, 129(3), 758S-767S.

Setchell, K. D., Brown, N. M., Zimmer-Nechemias, L., Brashears, W. T., Wolfe, B. E., Kirschner, A. S. and Heubi, J. E. (2002). Evidence for lack of absorption of soy isoflavone glycosides in humans, supporting the crucial role of intestinal metabolism for bioavailability. *The American Journal of Clinical Nutrition*, 76(2), 447-453.

Sevillano, D. M., Jankowiak, L., van Gaalen, T. L. T., van der Wielen, L. A. M., Hooshyar, N., van der Goot, A.-J. and Ottens, M. (2014). Mechanism of isoflavone adsorption from okara extracts onto food-grade resins. *Industrial & Engineering Chemistry Research*, 53(39), 15245-15252.

Shimonov, E. (2004). Stability and shelf life of bioactive compounds during food processing and storage: soy isoflavones. *Journal of Food Science*, 69(6), R160-R166.

Shuler, M. L. and Kargi, F. (2002). *Bioprocess Engineering* (2<sup>nd</sup> ed.): Prentice Hall New York.

Slepecky, R. A. and Hemphill, H. E. (2006). The genus *Bacillus*—nonmedical. *The Prokaryotes* (pp. 530-562): Springer.

Soccol, C., Marin, B., Raimbault, M. and Lebeault, J.-M. (1994). Potential of solid state fermentation for production of L (+)-lactic acid by *Rhizopus oryzae*. *Applied Microbiology and Biotechnology*, 41(3), 286-290.

- Song, J. Z., Mo, S. F., Yip, Y. K., Qiao, C. F., Han, Q. B. and Xu, H. X. (2007). Development of microwave assisted extraction for the simultaneous determination of isoflavonoids and saponins in Radix Astragalus by high performance liquid chromatography. *Journal of Separation Science*, 30(6), 819-824.
- Song, T., Barua, K., Buseman, G. and Murphy, P. A. (1998). Soy isoflavone analysis: quality control and a new internal standard. *The American Journal of Clinical Nutrition*, 68(6), 1474S-1479S.
- Steinkraus, K. (1995). *Handbook of Indigenous Fermented Foods, revised and expanded*: CRC Press.
- Sundhagul, M., Smanmathuroj, P. and Bhodacharoen, W. (1972). Thua-nao: a fermented soybean food of Northern Thailand. I. Traditional processing method. *Thai Journal of Agricultural Science*, 5, 43-56.
- Tamang, J. P. (1993). *Studies on The Microflora of Some Traditional Fermented Foods of The Darjeeling Hills and Sikkim*. (Ph.D. Thesis), University of North Bengal, India.
- Tamura, K., Nei, M. and Kumar, S. (2004). Prospects for inferring very large phylogenies by using the neighbor-joining method. *Proceedings of the National Academy of Sciences of the United States of America*, 101(30), 11030-11035.
- Tamura, K., Stecher, G., Peterson, D., Filipski, A. and Kumar, S. (2013). MEGA6: molecular evolutionary genetics analysis version 6.0. *Molecular Biology and Evolution*, 30(12), 2725-2729.
- Terigar, B. G., Balasubramanian, S., Boldor, D., Xu, Z., Lima, M. and Sabliov, C. M. (2010). Continuous microwave-assisted isoflavone extraction system: Design and performance evaluation. *Bioresource Technology*, 101(7), 2466-2471.

- Thompson, J. D., Gibson, T. J., Plewniak, F., Jeanmougin, F. and Higgins, D. G. (1997). The CLUSTAL X windows interface: flexible strategies for multiple sequence alignment aided quality analysis tools. *Nucleic Acids Research*, 25, 4876-4882.
- Toda, T., Sakamoto, A., Takayanagi, T. and Yokotsuka, K. (2001). Changes in isoflavone compositions of soybean during soaking in water. *Food Science and Technology Research*, 7(2), 171-175.
- Tsironi, T., Dermesonlouoglou, E., Giannakourou, M. and Taoukis, P. (2009). Shelf life modelling of frozen shrimp at variable temperature conditions. *LWT - Food Science and Technology*, 42(2), 664-671.
- Tsuchiya, K., Nagashima, T., Yamamoto, Y., Gomi, K., Kitamoto, K., Kumagai, C. and Tamura, G. (1994). High level secretion of calf chymosin using a glucoamylase-prochymosin fusion gene in *Aspergillus oryzae*. *Bioscience, Biotechnology, and Biochemistry*, 58(5), 895-899.
- U.S. Government. (2011). *Usda-Iowa State University Database on The Isoflavone Content of Foods*: General Books LLC.
- Wang, C. Y. and Chen, B. H. (2006). Tomato pulp as source for the production of lycopene powder containing high proportion of cis-isomers. *European Food Research and Technology*, 222(3-4), 347-353.
- Wang, H.-J. and Murphy, P. A. (1994). Isoflavone composition of American and Japanese soybeans in Iowa: effects of variety, crop year, and location. *Journal of Agricultural and Food Chemistry*, 42(8), 1674-1677.
- Wei, Q. K., Chen, T. R. and Chen, J. T. (2008). Use of *Bacillus subtilis* to enrich isoflavone aglycones in fermented natto. *Journal of the Science of Food and Agriculture*, 88(6), 1007-1011.

Wu, C. Y. and Lai, S. M. (2007). Preparative isolation of isoflavones from defatted soy flakes. *Journal of Liquid Chromatography & Related Technologies*, 30(11), 1617-1640.

Wu, Y., Wang, X. and Fan, E. (2012). Optimisation of ultrasound-assisted extraction of puerarin and total isoflavones from puerariae lobatae radix (*Pueraria lobata* (Wild.) Ohwi) with response surface methodology. *Phytochemical Analysis*, 23(5), 513-519.

Xiong, Q., Zhang, Q., Zhang, D., Shi, Y., Jiang, C. and Shi, X. (2014). Preliminary separation and purification of resveratrol from extract of peanut (*Arachis hypogaea*) sprouts by macroporous adsorption resins. *Food Chemistry*, 145(0), 1-7.

Yamada, Y., Katsura, K., Kawasaki, H., Widyastuti, Y., Saono, S., Seki, T., Uchimura, T. and Komagata, K. (2000). *Asaia bogorensis* gen. nov., sp. nov., an unusual acetic acid bacterium in the alpha-Proteobacteria. *International Journal of Systematic and Evolutionary Microbiology*, 50(2), 823-829.

Yin, L.-J., Li, L.-T., Li, Z.-G., Tatsumi, E. and Saito, M. (2004). Changes in isoflavone contents and composition of sufu (fermented tofu) during manufacturing. *Food Chemistry*, 87(4), 587-592.

Zhang, Y., Li, S., Wu, X. and Zhao, X. (2007). Macroporous resin adsorption for purification of flavonoids in *Houttuynia cordata* Thunb. *Chinese Journal of Chemical Engineering*, 15(6), 872-876.