

## REFERENCES

- [1] Cohen AR, Galanello R, Pennell DJ, Cunningham MJ, Vichinsky E. Thalassemia. Hematology Am Soc Hematol Educ Program 2004:14-34.
- [2] Fucharoen S, Winichagoon P. Haemoglobinopathies in Southeast Asia. Indian J Med Res 2011;134:498-506.
- [3] Weatherall DJ, Clegg JB. The thalassemia syndromes, ed Fourth. Oxford, Blackwell Science Ltd, 2001, ISBN 0-86542-664-3.
- [4] Bowden DK, Vickers MA, Higgs DR. A PCR-based strategy to detect the common severe determinants of alpha thalassaemia. Br J Haematol 1992;81:104-108.
- [5] Law DJ, Frossard PM, Rucknagel DL. Highly sensitive and rapid gene mapping using miniaturized blot hybridization: Application to prenatal diagnosis. Gene 1984;28:153-158.
- [6] Makonkawkeyoon L, Sanguanserm Sri T, Asato T, Nakashima Y, Takei H. Rapid detection of chain termination mutations in the alpha 2 globin gene. Blood 1993;82:3503-3504.
- [7] Smetanina NS, Huisman TH. Detection of alpha-thalassemia-2 (-3.7 kb) and its corresponding triplication (alpha)(alpha)(alpha) (anti-3.7 kb) by PCR: An improved technical change. Am J Hematol 1996;53:202-203.
- [8] Liao C, Zhou JY, Xie XM, Li R, Li DZ. Cord blood analysis for rapid prenatal confirmation of Hb Bart's disease using the Sebia Capillary electrophoresis system. Hemoglobin 2012;36:186-191.

- [9] Munkongdee T, Pichanun D, Butthep P, Klamchuen S, Chalermopolprapa V, Winichagoon P, Svasti S, Fucharoen S. Quantitative analysis of Hb Bart's in cord blood by capillary electrophoresis system. *Ann Hematol* 2011;90:741-746.
- [10] Tanphaichitr VS, Pung-amritt P, Puchaiwatananon O, Winichagoon P, Fucharoen S, Suvatte V, Wasi P. Studies of hemoglobin Bart's and deletion of alpha-globin genes from cord blood in Thailand. *Birth Defects Orig Artic Ser* 1987;23:15-21.
- [11] Lie-Injo LE, Solai A, Herrera AR, Nicolaisen L, Kan YW, Wan WP, Hasan K. Hb Bart's level in cord blood and deletions of alpha-globin genes. *Blood* 1982;59:370-376.
- [12] Makonkawkeyoon L, Ongchai S, Sanguansermisri T. Determination of hemoglobin Bart's in alpha thalassemia traits by two-site immunoradiometric assay. I. Purification of hemoglobin Bart's and preparation of specific anti-Hb Bart's. *J Med Assoc Thai* 1992;75:508-513.
- [13] Makonkawkeyoon L, Ongchai S, Sanguansermisri T. Determination of hemoglobin Bart's in alpha thalassemia traits by two-site immunoradiometric assay. II. Detection of hemoglobin Bart's in alpha thalassemia traits. *J Med Assoc Thai* 1992;75:565-569.
- [14] Wasi P, Pravatmuang P, Winichagoon P. Immunologic diagnosis of alpha-thalassemia traits. *Hemoglobin* 1979;3:21-31.
- [15] Makonkawkeyoon L, Pharephan S, Makonkawkeyoon S. Production of a mouse hybridoma secreting monoclonal antibody highly specific to hemoglobin Bart's (gamma4). *Lab Hematol* 2006;12:193-200.
- [16] Makonkawkeyoon L, Pharephan S, Tuntiwechapikul W, Makonkawkeyoon S. Application of monoclonal and polyclonal antibodies to Hb Bart's for the detection of  $\alpha$  thalassemias. *The Open Hematol J* 2009;3:11-17.

- [17] Makonkawkeyoon L, Pharephan S, Sirivatanapa P, Tuntiwechapikul W, Makonkawkeyoon S. Development of an ELISA strip for the detection of alpha thalassemias. *Haematologica* 2010;95:338-339.
- [18] Tatu T, Kiewkarnkha T, Khuntarak S, Khamrin S, Suwannasin S, Kasinrerak W. Screening for co-existence of alpha-thalassemia in beta-thalassemia and in HbE heterozygotes via an enzyme-linked immunosorbent assay for Hb Bart's and embryonic zeta-globin chain. *Int J Hematol* 2012;95:386-393.
- [19] Tayapiwatana C, Kuntaruk S, Tatu T, Chiampanichayakul S, Munkongdee T, Winichagoon P, Fuchareon S, Kasinrerak W. Simple method for screening of alpha-thalassaemia 1 carriers. *Int J Hematol* 2009;89:559-567.
- [20] Jo EC, Park HJ, Kim DI, Moon HM: Repeated fed-batch culture of hybridoma cells in nutrient-fortified high-density medium. *Biotechnol Bioeng* 1993;42:1229-1237.
- [21] Bird RE, Hardman KD, Jacobson JW, Johnson S, Kaufman BM, Lee SM, Lee T, Pope SH, Riordan GS, Whitlow M. Single-chain antigen-binding proteins. *Science* 1988;242:423-426.
- [22] Higgs DR, Weatherall DJ: The alpha thalassaemias. *Cell Mol Life Sci* 2009;66:1154-1162.
- [23] Sen R, Chakrabarti S, Sengupta B, De M, Halder A, Poddar S, Gajra B, Talukder G, Sengupta S. Alpha-thalassemia among tribal populations of Eastern India. *Hemoglobin* 2005;29:277-280.
- [24] Rudra T, Chakrabarti S, Sengupta B: Alpha thalassemia. Experience of referral cases in Kolkata, India. *Int J Hum Genet* 2008;8:357-360.
- [25] Nadkarni A, Phanasgaonkar S, Colah R, Mohanty D, Ghosh K. Prevalence and molecular characterization of alpha-thalassemia syndromes among Indians. *Genet Test* 2008;12:177-180.

- [26] Xu XM, Zhou YQ, Luo GX, Liao C, Zhou M, Chen PY, Lu JP, Jia SQ, Xiao GF, Shen X, Li J, Chen HP, Xia YY, Wen YX, Mo QH, Li WD, Li YY, Zhuo LW, Wang ZQ, Chen YJ, Qin CH, Zhong M. The prevalence and spectrum of alpha and beta thalassaemia in guangdong province: Implications for the future health burden and population screening. *J Clin Pathol* 2004;57:517-522.
- [27] Laig M, Pape M, Hundrieser J, Flatz G, Sanguansermsri T, Das BM, Dekar R, Yongvanit P, Mularlee N. The distribution of the Hb Constant Spring gene in Southeast Asian populations. *Hum Genet* 1990;84:188-190.
- [28] Tritipsombut J, Sanchaisuriya K, Phollarp P, Bouakhasith D, Sanchaisuriya P, Fucharoen G, Fucharoen S, Schelp FP. Micromapping of thalassemia and hemoglobinopathies in different regions of Northeast Thailand and Vientiane, Laos people's democratic republic. *Hemoglobin* 2012;36:47-56.
- [29] Nicholls RD, Fischel-Ghodsian N, Higgs DR. Recombination at the human alpha-globin gene cluster: Sequence features and topological constraints. *Cell* 1987;49:369-378.
- [30] Fischel-Ghodsian N, Vickers MA, Seip M, Winichagoon P, Higgs DR: Characterization of two deletions that remove the entire human zeta-alpha globin gene complex (-<sup>THAI</sup> and -<sup>FIL</sup>). *Br J Haematol* 1988;70:233-238.
- [31] Fortina P, Delgrosso K, Rappaport E, Poncz M, Ballas SK, Schwartz E, Surrey S. A large deletion encompassing the entire alpha-like globin gene cluster in a family of Northern European extraction. *Nucleic Acids Res* 1988;16:11223-11235.
- [32] Lauer J, Shen CK, Maniatis T. The chromosomal arrangement of human alpha-like globin genes: Sequence homology and alpha-globin gene deletions. *Cell* 1980;20:119-130.

- [33] Zimmer EA, Martin SL, Beverley SM, Kan YW, Wilson AC. Rapid duplication and loss of genes coding for the alpha chains of hemoglobin. *Proc Natl Acad Sci U S A* 1980;77:2158-2162.
- [34] Michelson AM, Orkin SH. The 3' untranslated regions of the duplicated human alpha-globin genes are unexpectedly divergent. *Cell* 1980;22:371-377.
- [35] Hess JF, Schmid CW, Shen CK. A gradient of sequence divergence in the human adult alpha-globin duplication units. *Science* 1984;226:67-70.
- [36] Embury SH, Miller JA, Dozy AM, Kan YW, Chan V, Todd D. Two different molecular organizations account for the single alpha-globin gene of the alpha-thalassemia-2 genotype. *J Clin Invest* 1980;66:1319-1325.
- [37] Goossens M, Dozy AM, Embury SH, Zachariades Z, Hadjiminias MG, Stamatoyannopoulos G, Kan YW. Triplicated alpha-globin loci in humans. *Proc Natl Acad Sci U S A* 1980;77:518-521.
- [38] Higgs DR, Clegg JB, Weatherall DJ, Serjeant BE, Serjeant GR. Interaction of the alpha alpha alpha globin gene haplotype and sickle haemoglobin. *Br J Haematol* 1984;57:671-678.
- [39] Kulozik AE, Kar BC, Serjeant GR, Serjeant BE, Weatherall DJ. The molecular basis of alpha thalassemia in India. Its interaction with the sickle cell gene. *Blood* 1988;71:467-472.
- [40] Lacerra G, Fioretti G, De Angioletti M, Pagano L, Guarino E, de Bonis C, Viola A, Maglione G, Scarallo A, De Rosa L, et al.. (alpha)alpha 5.3: A novel alpha(+)-thalassemia deletion with the breakpoints in the alpha 2-globin gene and in close proximity to an Alu family repeat between the psi alpha 2- and psi alpha 1-globin genes. *Blood* 1991;78:2740-2746.
- [41] Zhao JB, Zhao L, Fei YJ, Liu JC, Huisman TH. A novel alpha-thalassemia-2 (-2.7-kb) observed in a Chinese patient with Hb H disease. *Am J Hematol* 1991;38:248-249.

- [42] Orkin SH, Goff SC, Hechtman RL. Mutation in an intervening sequence splice junction in man. *Proc Natl Acad Sci USA* 1981;78:5041-5045.
- [43] Felber BK, Orkin SH, Hamer DH. Abnormal RNA splicing causes one form of alpha thalassemia. *Cell* 1982;29:895-902.
- [44] Pressley L, Higgs DR, Clegg JB, Perrine RP, Pembrey ME, Weatherall DJ. A new genetic basis for hemoglobin-H disease. *N Engl J Med* 1980;303:1383-1388.
- [45] Olivieri NF, Chang LS, Poon AO, Michelson AM, Orkin SH. An alpha-globin gene initiation codon mutation in a black family with HbH disease. *Blood* 1987;70:729-732.
- [46] Pichanun D, Munkongdee T, Klamchuen S, Butthep P, Winichagoon P, Fucharoen S, Svasti S: Molecular screening of the Hb Constant Spring (codon 142, TAA>CAA, alpha2) and Pakse' (codon 142, TAA>TAT, alpha2) mutations in Thailand. *Hemoglobin* 2010;34:582-586.
- [47] Hunt JA, Lehmann H. Haemoglobin "Bart's": A foetal haemoglobin without alpha-chains. *Nature* 1959;184:872-873.
- [48] Kekwick RA, Lehmann H: Sedimentation characteristics of the gamma-chain haemoglobin (haemoglobin Bart's). *Nature* 1960;187:158.
- [49] Fessas P, Papaspyrou A: New fast hemoglobin associated with thalassemia. *Science* 1957;126:1119.
- [50] Ager JA, Lehmann H: Observations on some fast haemoglobins: K, J, N, and Bart's. *Br Med J* 1958;1:929-931.
- [51] Pootrakul S, Wasi P, Na-Nakorn S: Studies on haemoglobin Bart's (Hb  $\gamma_4$ ) in Thailand: The incidence and the mechanism of occurrence in cord blood. *Ann Hum Genet* 1967;31:149-166.

- [52] Kontermann R, Dubel S: Antibody engineering. Heidelberg, Springer-Verlag, 2001, ISBN 3-540-41354-5.
- [53] Kohler G, Milstein C. Continuous cultures of fused cells secreting antibody of predefined specificity. *Nature* 1975;256:495-497.
- [54] Neuberger MS, Williams GT, Fox RO: Recombinant antibodies possessing novel effector functions. *Nature* 1984;312:604-608.
- [55] Neuberger MS, Williams GT, Mitchell EB, Jouhal SS, Flanagan JG, Rabbitts TH. A hapten-specific chimaeric IgE antibody with human physiological effector function. *Nature* 1985;314:268-270.
- [56] Riechmann L, Foote J, Winter G. Expression of an antibody Fv fragment in myeloma cells. *J Mol Biol* 1988;203:825-828.
- [57] Feys V, De Waele P, Van de Voorde A, Casneuf P, Fiers W. Expression of functional mouse antibodies directed against the tumour marker human placental alkaline phosphatase in non-lymphoid cells. *Int J Cancer Suppl* 1988;2:26-27.
- [58] Glockshuber R, Malia M, Pfitzinger I, Pluckthun A. A comparison of strategies to stabilize immunoglobulin Fv-fragments. *Biochemistry* 1990;29:1362-1367.
- [59] Huston JS, Levinson D, Mudgett-Hunter M, Tai MS, Novotny J, Margolies MN, Ridge RJ, Brucoleri RE, Haber E, Crea R, et al. Protein engineering of antibody binding sites: Recovery of specific activity in an anti-digoxin single-chain Fv analogue produced in *Escherichia coli*. *Proc Natl Acad Sci USA* 1988;85:5879-5883.
- [60] Peterson E, Owens SM, Henry RL. Monoclonal antibody form and function: Manufacturing the right antibodies for treating drug abuse. *AAPS J* 2006;8:E383-390.

- [61] Sandborn WJ, Feagan BG, Stoinov S, Honiball PJ, Rutgeerts P, Mason D, Bloomfield R, Schreiber S. Certolizumab pegol for the treatment of Crohn's disease. *N Engl J Med* 2007;357:228-238.
- [62] Kaushik VV, Moots RJ. CDP-870 (certolizumab) in rheumatoid arthritis. *Expert Opin Biol Ther* 2005;5:601-606.
- [63] Alvarez RD, Barnes MN, Gomez-Navarro J, Wang M, Strong TV, Arafat W, Arani RB, Johnson MR, Roberts BL, Siegal GP, Curiel DT. A cancer gene therapy approach utilizing an anti-erbB-2 single-chain antibody-encoding adenovirus (AD21): A phase I trial. *Clin Cancer Res* 2000;6:3081-3087.
- [64] Strube RW, Chen SY. Characterization of anti-cyclin E single-chain Fv antibodies and intrabodies in breast cancer cells: Enhanced intracellular stability of novel sFv-F(c) intrabodies. *J Immunol Methods* 2002;263:149-167.
- [65] Miller SA, Dykes DD, Polesky HF: A simple salting out procedure for extracting DNA from human nucleated cells. *Nucleic Acids Res* 1988;16:1215.
- [66] Baysal E, Huisman TH: Detection of common deletional alpha-thalassemia-2 determinants by PCR. *Am J Hematol* 1994;46:208-213.
- [67] Harlow E, Lane D. *Antibodies: A laboratory manual*. New York, Cold Spring Harbor Laboratory, 1988, ISBN 0-87969-314-2.
- [68] Krebber A, Bornhauser S, Burmester J, Honegger A, Willuda J, Bosshard HR, Pluckthun A. Reliable cloning of functional antibody variable domains from hybridomas and spleen cell repertoires employing a reengineered phage display system. *J Immunol Methods* 1997;201:35-55.
- [69] Sambrook J, Russell D. *Molecular cloning: A laboratory manual*, third ed. New York, Cold Spring Harbor Laboratory, 2001, ISBN 0-87969-577-3.


- [70] Laemmli UK. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 1970;227:680-685.
- [71] Beatty JD, Beatty BG, Vlahos WG. Measurement of monoclonal antibody affinity by non-competitive enzyme immunoassay. *J Immunol Methods* 1987;100:173-179.
- [72] Weatherall DJ: The thalassemias; in Stamatoyannopoulos G, Majerus PW, Perlmutter RM, Varmus H (eds): The molecular basis of blood diseases. Pennsylvania, W.B. Saunders Company, 2001, ISBN 0-72167-671-5.
- [73] Hundrieser J, Sanguansermsri T, Papp T, Flatz G: Alpha-thalassemia in northern Thailand. Frequency of deletional types characterized at the DNA level. *Hum Hered* 1988;38:211-215.
- [74] Lemmens-Zygulska M, Eigel A, Helbig B, Sanguansermsri T, Horst J, Flatz G: Prevalence of alpha-thalassemias in northern Thailand. *Hum Genet* 1996;98:345-347.
- [75] Wanapirak C, Muninthorn W, Sanguansermsri T, Dhananjayanonda P, Tongsong T: Prevalence of thalassemia in pregnant women at Maharaj Nakorn Chiang Mai Hospital. *J Med Assoc Thai* 2004;87:1415-1418.
- [76] Tongsong T, Wanapirak C, Sirivatanapa P, Sanguansermsri T, Sirichotiyakul S, Piyamongkol W, Chanprapaph P: Prenatal control of severe thalassaemia: Chiang Mai strategy. *Prenat Diagn* 2000;20:229-234.
- [77] Wu TT, Kabat EA. An analysis of the sequences of the variable regions of bence jones proteins and myeloma light chains and their implications for antibody complementarity. *J Exp Med* 1970;132:211-250.
- [78] Choi GH, Lee DH, Min WK, Cho YJ, Kweon DH, Son DH, Park K, Seo JH. Cloning, expression, and characterization of single-chain variable fragment antibody against mycotoxin deoxynivalenol in recombinant *Escherichia coli*. *Protein Expr Purif* 2004;35:84-92.

- [79] Burgess RR. Refolding solubilized inclusion body proteins; in Burgess RR, Deutscher PM (eds): Methods in enzymology guide to protein purification. Wisconsin, Elsevier Inc., 2009, vol 463, pp 259-282, ISBN 0076-6879.
- [80] Tsumoto K, Shinoki K, Kondo H, Uchikawa M, Juji T, Kumagai I. Highly efficient recovery of functional single-chain fv fragments from inclusion bodies overexpressed in *Escherichia coli* by controlled introduction of oxidizing reagent--application to a human single-chain Fv fragment. J Immunol Methods 1998;219:119-129.
- [81] Pluckthun A. Mono- and bivalent antibody fragments produced in *Escherichia coli*: Engineering, folding and antigen binding. Immunol Rev 1992;130:151-188.
- [82] Field H, Yarranton GT, Rees AR. Expression of mouse immunoglobulin light and heavy chain variable regions in *Escherichia coli* and reconstitution of antigen-binding activity. Protein Eng 1990;3:641-647.
- [83] Ahmad ZA, Yeap SK, Ali AM, Ho WY, Alitheen NB, Hamid M: Scfv antibody: Principles and clinical application. Clin Dev Immunol 2012;2012:980250.
- [84] Singh SM, Panda AK: Solubilization and refolding of bacterial inclusion body proteins. J Biosci Bioeng 2005;99:303-310.
- [85] Basu A, Li X, Leong SS: Refolding of proteins from inclusion bodies: Rational design and recipes. Appl Microbiol Biotechnol 2011;92:241-251.
- [86] Miller KD, Weaver-Feldhaus J, Gray SA, Siegel RW, Feldhaus MJ. Production, purification, and characterization of human scFv antibodies expressed in *Saccharomyces cerevisiae*, *Pichia pastoris*, and *Escherichia coli*. Protein Expr Purif 2005;42:255-267.

- [87] Menezes MA, Aires KA, Ozaki CY, Ruiz RM, Pereira MC, Abreu PA, Elias WP, Ramos OH, Piazza RM. Cloning approach and functional analysis of anti-intimin single-chain variable fragment (scFv). *BMC Res Notes* 2011;4:30.
- [88] Liu X, Wang H, Liang Y, Yang J, Zhang H, Lei H, Shen Y, Sun Y. Production and characterization of a single-chain Fv antibody-alkaline phosphatase fusion protein specific for clenbuterol. *Mol Biotechnol* 2010;45:56-64.
- [89] Cho WK, Sohn U, Kwak JW. Production and in vitro refolding of a single-chain antibody specific for human plasma apolipoprotein A-I. *J Biotechnol* 2000;77:169-178.
- [90] Intorasoot S, Tanaka H, Shoyama Y, Leelamanit W: Characterization and diagnostic use of a recombinant single-chain antibody specific for the gp116 envelop glycoprotein of Yellow head virus. *J Virol Methods* 2007;143:186-193.
- [91] Liddell JE, Cryer A. A practical guide to monoclonal antibodies. West Sussex, John Wiley and Sons Ltd., 1991, ISBN 0-471-92905-0.
- [92] Martsev SP, Kravchuk ZI, Chumanevich AA, Vlasov AP, Dubnovitsky AP, Bessalov IA, Arosio P, Deyev SM. Anti-ferritin single-chain antibody: A functional protein with incomplete folding? *FEBS Lett* 1998;441:458-462.
- [93] Marks JD, Griffiths AD, Malmqvist M, Clackson TP, Bye JM, Winter G. By-passing immunization: Building high affinity human antibodies by chain shuffling. *Biotechnology (N Y)* 1992;10:779-783.

## LIST OF PUBLICATIONS

- 1) Pharephan S, Intorasoot S, Tuntiwechapikul W, Makonkawkeyoon S, Makonkawkeyoon L. Cloning and expression of a recombinant single-chain variable fragment antibody specific to hemoglobin Bart's. Asian Pac J Health Sci 2015;2(1):28-35.
- 2) Pharephan S, Sirivatanapa P, Makonkawkeyoon S, Tuntiwechapikul W, Makonkawkeyoon L. Prevalence of  $\alpha$ -thalassemia genotypes in Thai pregnant women at Maharaj Nakorn Chiang Mai Hospital, northern Thailand. Indian J Med Res 2015;x:pp.



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่  
Copyright© by Chiang Mai University  
All rights reserved