

CONTENTS

	Page
Acknowledgement	d
Abstract in Thai	e
Abstract in English	g
List of Tables	l
List of Figures	m
List of Abbreviations	p
Statements of Originality in Thai	s
Statements of Originality in English	t
Chapter 1 Introduction	1
1.1 Statement and significant of the problems	1
1.2 Objectives	2
1.3 Literature Review	2
1.4 Principle and Rationales	13
Chapter 2 Research design and Methods	15
2.1 Research design	15
2.2 Classification of α -thalassemia genotypes by gap-PCR and PCR-RFLP	17
2.3 Growth of hybridoma and single-cell cloning	21
2.4 Detection of mouse anti-Hb Bart's by indirect ELISA	22
2.5 Isolation of RNA, synthesis of cDNA and antibody gene amplification	23
2.6 Cloning of the V _H and V _L gene	27
2.7 Analysis of the V _H and V _L gene	30

2.8	Construction of the full length scFv gene	33
2.9	Cloning of the full length scFv gene	36
2.10	Optimization of IPTG induction	39
2.11	Expression of the scFv antibody	40
2.12	Purification of the scFv antibody by affinity chromatography and refolding of the scFv antibody	41
2.13	Specificity of the refolded scFv antibody and the parent monoclonal antibody by Western blot and indirect ELISA	41
2.14	Sensitivity of the refolded scFv antibody and the parent monoclonal by dot blot ELISA	42
2.15	Affinity constant determination of the refolded scFv antibody and the parent monoclonal antibody by indirect ELISA	43
2.16	Stability of the refolded scFv antibody	43
Chapter 3 Results		44
3.1	Genotyping of α -thalassemias in primigravidae volunteers	44
3.2	Specificity of mouse anti-Hb Bart's monoclonal antibody	46
3.3	RNA isolation, cDNA synthesis and antibody gene amplification	48
3.4	V _H and V _L gene cloning	49
3.5	Analysis of V _H and V _L DNA sequence	54
3.6	Full length scFv gene construction	58
3.7	Full length scFv gene cloning	59
3.8	Full length scFv gene analysis	59
3.9	Optimization of IPTG induction and the full length scFv expression	62
3.10	ScFv antibody expression	63
3.11	ScFv antibody purification and refolding	65
3.12	Specificity of the refolded scFv antibody and the parent monoclonal antibody	66
3.13	Sensitivity of the refolded scFv antibody and the parent monoclonal antibody	69
3.14	Affinity constant of the refolded scFv antibody and the parent monoclonal antibody	70

3.15 Stability of the refolded scFv antibody	71
Chapter 4 Discussion	72
Chapter 5 Conclusion	80
References	81
List of Publications	92
Appendix	93
Appendix A List of chemicals and materials used in study	93
Appendix B List of instruments used in study	97
Appendix C List of buffers and media used in study	99
Appendix D Bacterial strains and their vectors	111
Appendix E Protocol of RNA isolation	118
Appendix F Protocol of PCR product purification from agrose gel	121
Appendix G Protocol of bacterial competent cells preparation	124
Appendix H Protocol of plasmid DNA purification	125
Appendix I Protocol of SDS-polyacrylamide and native polyacrylamide gel preparation	128
Appendix J Protocol of proteins transfer from polyacrylamide gels to membranes	130
Appendix K Protocol of protein concentration determination	132
Appendix L Protocol of scFv antibody purification by Ni-NTA affinity chromatography	135
Curriculum Vitae	136

LIST OF TABLES

	Page
Table 1.1 Phenotype of unstable α -chain variants	7
Table 1.2 Hb Bart's levels in cord blood of difference genotypes of α -thalassemia	10
Table 2.1 Sequences of primers for PCR genotyping of α -thalassemias	21
Table 2.2 List of antibody-specific DNA primers used for the V_H gene amplification	24
Table 2.3 List of antibody-specific DNA primers used for the V_L gene amplification	26
Table 2.4 Primers designed for construction of the full length scFv gene	34
Table 3.1 Classification of α -thalassemia genotypes in 638 primigravidarum volunteers from Maharaj Nakorn Chiang Mai Hospital	46
Table 3.2 Specificity of mouse monoclonal antibody to Hbs; Bart's, A ₂ , E, F, A and H by indirect ELISA	47
Table 3.3 Recovery yield of the scFv antibody from each step of purification	65
Table 3.4 Specificity of the refolded scFv antibody to Hbs; Bart's, A ₂ , E, F, A and H by indirect ELISA	68

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
 Copyright© by Chiang Mai University
 All rights reserved

LIST OF FIGURES

	Page
Figure 1.1	Deletions that cause α^0 -thalassemia. 4
Figure 1.2	The mechanism by which the common deletions underlying α^+ -thalassemia occur 5
Figure 1.3	Deletions that cause α^+ -thalassemia 6
Figure 1.4	The Fv fragment contains the antigen-binding regions of antibodies 11
Figure 1.5	Antibody model showing subunit composition 12
Figure 2.1	Schematic representation of the procedures of the scFv generation 16
Figure 2.2	The locations on chromosome 16 of the specific primers for the amplification of α^0 -thalassemia -- ^{SEA} type 18
Figure 2.3	The locations on chromosome 16 of the specific primers for the amplification of α^+ -thalassemia - $\alpha^{3.7}$ and - $\alpha^{4.2}$ type 19
Figure 2.4	The locations on chromosome 16 of the specific primers for the amplification of HbCS and <i>MseI</i> restriction sites 20
Figure 2.5	The circle map, promoter and T overhang cloning sequences of pGEM [®] -T Easy vector 29
Figure 2.6	Kabat and Chothia numbering scheme for the identification of CDRs-H 31
Figure 2.7	Kabat and Chothia numbering scheme for the identification of CDRs-L 32
Figure 2.8	The schematic diagram of the construction of the full length scFv gene by SOE-PCR 35
Figure 2.9	The circle map, promoter and multiple cloning sequences of pET28a(+) expression vector 38
Figure 2.10	Expression vector map of the pET28a(+)-scFv anti-Hb Bart's 39
Figure 3.1	PCR detection of the common genotypes of α -thalassemias 45
Figure 3.2	Reactivity of mouse monoclonal antibody to Hbs; Bart's, A ₂ , E, F, A and H by indirect ELISA 47

Figure 3.3	Agarose gel electrophoresis of amplicons of V _L and V _H gene from cDNA of mouse hybridoma	48
Figure 3.4	Agarose gel electrophoresis of amplicons of <i>E. coli</i> TOP10F white bacterial colonies for the presence of pGEM [®] -T-V _H by colony PCR	50
Figure 3.5	Agarose gel electrophoresis of amplicons of <i>E. coli</i> TOP10F white bacterial colonies for the presence of pGEM [®] -T-V _L by colony PCR	52
Figure 3.6	Agarose gel electrophoresis of <i>Eco</i> R I restriction digestion analysis of purified plasmid DNA from the pGEM [®] -T-V _H	55
Figure 3.7	Nucleotide and amino acid sequences of V _H	56
Figure 3.8	Nucleotide and amino acid sequences of V _L	57
Figure 3.9	Nucleotide and amino acid sequences of the other V _L	57
Figure 3.10	Agarose gel electrophoresis of amplicons of the V _H , V _L and full length scFv genes	58
Figure 3.11	Agarose gel electrophoresis of amplicons of <i>E. coli</i> BL21(DE3) colonies for the presence of pET28a(+)-scFv anti-Hb Bart's by colony PCR	60
Figure 3.12	Nucleotide and amino acid sequences of the N-terminal histidine fusion full length scFv	61
Figure 3.13	Polyacryamide gel electrophoresis of the expression of the scFv antibody in <i>E. coli</i> BL21(DE3) after IPTG induction at various time	62
Figure 3.14	Polyacryamide gel electrophoresis of the expression of the scFv antibody in <i>E. coli</i> BL21(DE3) for analysis of induction condition	63
Figure 3.15	Western blot analysis of the expression of N-terminal histidine fusion scFv antibody	64
Figure 3.16	Polyacryamide gel electrophoresis of the refolded scFv antibody	66
Figure 3.17	Native polyacrylamide gel electrophoresis and Western blot analysis of the specific binding of the refolded scFv antibody and the parent monoclonal antibody to Hb Bart's	67
Figure 3.18	Reactivity of the refolded scFv antibody to Hb Bart's, A ₂ , E, F, A and H by indirect ELISA	68
Figure 3.19	Sensitivity of the refolded scFv antibody and the parent monoclonal antibody to Hb Bart's by dot blot ELISA	69

Figure 3.20 Reactivity of the refolded scFv antibody and the parent monoclonal antibody to various concentrations of Hb Bart's for analysis of the affinity constant by indirect ELISA

70




ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved

LIST OF ABBREVIATIONS

AMV	Avian myeloblastosis virus
bp	Base pair
cDNA	Complementary deoxyribonucleic acid
CDRs	Complementarity determining regions
CO ₂	Carbon dioxide
CS	Constant Spring
DNA	Deoxyribonucleic acid
dNTP	Deoxynucleotide triphosphate
<i>E. coli</i>	<i>Escherichia coli</i>
EDTA	Ethylenediaminetetraacetic acid
ELISA	Enzyme linked immunosorbent assay
FBS	Fetal bovine serum
Fc	Constant fragment of immunoglobulin
Fv	Variable fragment of immunoglobulin
g	Gram
GST	Glutathione-S-transferase
h	Hour
Hb	Hemoglobin
His	Histidine
HRP	Horseradish peroxidase
IBs	Inclusion bodies
IEF	Isoelectric focusing
Igs	Immunoglobulins
IMAC	Immobilized metal affinity chromatography
IMDM	Iscoe's Modified Dulbecco's Medium
IPTG	Isopropyl-β-D-thiogalactopyranoside
K_a	Affinity constant

kb	Kilobase
kDa	Kilodalton
L	Liter
LB	Luria Bertani
M	Molar
mAb	Monoclonal antibody
MCH	Mean corpuscular hemoglobin
MCV	Mean corpuscular volume
MED	Mediterranean
mg	Milligram
min	Minute
mL	Milliliter
mM	Millimolar
μg	Microgram
μL	Microliter
μM	Micromolar
N	Normality
Ni-NTA	Nickle-nitrilotriacetic acid
nm	Nanometer
OD	Optical density
PAGE	Polyacrylamide gel electrophoresis
PBS	Phosphate buffer saline
PBS-T	Phosphate buffer saline-Tween 20
PCR	Polymerase chain reaction
PMSF	Phenylmethyl sulfonyl fluoride
PVDF	Polyvinylidene difluoride
RBC	Red blood cell
RFLP	Restriction fragment length polymorphism
RNA	Ribonucleic acid
rpm	Revolution per minute
RT-PCR	Reverse transcriptase polymerase chain reaction
s	Second

ScFv	Single-chain variable fragment
SDS	Sodium dodecyl sulfate
SEA	Southeast Asia
SOE-PCR	Splice overlapped extension-polymerase chain reaction
TAE	Tris-acetate-EDTA buffer
TMB	3, 3', 5, 5'-Tetramethylbenzidine
Trx	Thioredoxin
UV	Ultraviolet
V	Voltage
V _H	Variable domains of immunoglobulin heavy chain
V _L	Variable domains of immunoglobulin light chain



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
 Copyright© by Chiang Mai University
 All rights reserved


ข้อความแห่งการริเริ่ม

1. ศึกษารูปแบบพันธกรรมและความซุกที่เป็นปัจจุบันของโรคอัลฟาธาลัสซีเมียในหญิงตั้งครรภ์ท้องแรกที่มีภูมิลำเนาอยู่ในภาคเหนือของประเทศไทยด้วยวิธีเก็บฟีชีอาร์และฟีชีอาร์-อาร์เอฟแอลพี
2. รีคอมบีแนนท์แอนติบอดีชนิดเอสซีเอฟวีที่จำเพาะต่อฮีโมโกลบินบาร์ทส์ถูกสร้างขึ้นใหม่ด้วยเทคโนโลยีรีคอมบีแนนท์ดีเอ็นเอเพื่อใช้ทดแทน โมโนโคลนอลแอนติบอดีเป็นการแก้ปัญหาเซลล์ไฮบริดโดมาที่อาจสูญเสียความสามารถในการสร้างและหลั่งโมโนโคลนอลแอนติบอดีเมื่อเลี้ยงเซลล์เป็นเวลานาน

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved

STATEMENTS OF ORIGINALITY

1. Common genotypes and update data of the prevalence of α -thalassemias were determined in northern Thai primigravidarum by gap-PCR and PCR-RFLP.
2. A novel recombinant scFv antibody specific to Hb Bart's was generated using recombinant DNA technology as an alternative to monoclonal antibody dissolving the problems of gradual loss of the synthesis and secretion of monoclonal antibody during long-term cultivation of the hybridimas.



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved