

REFERENCES

- [Ahn, 2012] Ahn, J.H., Hong, I.P., Bok, J.I., Kim, B.Y., Song, J., Weon, H.Y., “Pyrosequencing analysis of the bacterial communities in the guts of honey bees *Apis cerana* and *Apis mellifera* in Korea,” J. Microbiol., 50, 2012, pp.735-745.
- [Allen, 1990] Allen, M.F., Ball, B.V., Underwood, B.A., “An isolate of *Melisococcus pluton* from *Apis laboriosa*,” J. Invertebr. Pathol., 55, 1990, pp. 439-440.
- [Anderson, 2014] Anderson, K.E., Carroll, M.J., Sheehan, T., Mott, B.M., Maes, P., Corby-Harris, V., “Hive-stored pollen of honey bees: many lines of evidence are consistent with pollen preservation, not nutrient conversion,” Mol. Ecol., 23, 2014, pp. 5904-5917.
- [Anderson, 2011] Anderson, K.E., Sheehan, T.H., Eckholm, B.J., Mott, B.M., DeGrandi-Hoffman, G., “An emerging paradigm of colony health: microbial balance of the honey bee and hive (*Apis mellifera*),” Insectes Soc., 58, 2011, pp. 431-444.
- [Anderson, 2013] Anderson, K.E., Sheehan, T.H., Mott, B.M., Maes, P., Snyder, L., Schwan, M.R., Walton, A., Jones, B.M., Corby-Harris, V., “Microbial ecology of the hive and pollination landscape: bacterial associates from floral nectar, the alimentary tract and stored food of the honey bees (*Apis mellifera*),” PLOS ONE, 8(12), 2013, e83125.
- [Aronstein, 2010] Aronstein, K.A., Murray, K.D., “Chalkbrood disease in honey bees,” J. Invertebr. Pathol., 103, 2010, pp. 20-29.

- [Babendreier, 2007] Babendreier, D., Joller, D., Romeis, J., Bigler, F., Widmer, F., "Bacterial community structures in honeybee intestines and their response to two insecticidal proteins," FEMS Microbiol. Ecol., 59, 2007, pp. 600-610.
- [Bailey, 1974] Bailey, L., "An unusual type of *Streptococcus pluton* from the Eastern hive bee," J. Invertebr. Pathol., 23, 1974, pp. 246-247.
- [Bailey, 1983] Bailey, L., "*Melissococcus pluton*, the cause of European foulbrood of honey bee (*Apis* spp.)," J. Appl. Bacteriol., 55, 1983, pp. 65-69.
- [Bailey, 1956] Bailey, L., "Aetiology of European foulbrood; a disease of larval honeybee," Nature, 178, 1956, pp. 1130.
- [Bailey, 1991] Bailey, L., Ball, B.V., "Honey bee pathogen," Academic Press, New York, 1991
- [Bawa, 1990] Bawa, K.S., "Plant-pollinator interactions in tropical rain forests," Annu. Rev. Ecol. Syst., 21, 1990, pp. 399-422.
- [Buchon, 2009] Buchon, N., Broderick, N.A., Chakrabarti, S., Lemaitre, B., "Invasive and indigenous microbiota impact intestinal stem cell activity through multiple pathways in *Drosophila*," Genes Dev. 23, 2009, pp. 2333-2344.
- [Chaimanee, 2010] Chaimanee, V., Pettis, J.S., Chen, Y., Evans, J.D., Khongphinitbunjong, K., Chantawannakul, P., "Susceptibility of four different honey bee species to *Nosema ceranae*," Vet. Parasitol., 193, 2010, pp. 260-265.

- [Chen, 2009] Chen, Y.P., Evans, J.D., Murphy, C., Gutell, R., Zuker, M., Gundensen-Rindal, D., Pettis, J.S., "Morphological, molecular, and phylogenetic characterization of *Nosema ceranae*, a microsporidian parasite isolated from the European honey bee, *Apis mellifera*," J. Euk. Microbiol., 56, 2009, pp. 142–147.
- [Clark, 1978] Clark, T.B., "A filamentous virus of the honey bee," J. Invertebr. Pathol., 32, 1978, pp. 332-340.
- [Clark, 1985] Clark, T.B., Whitcomb, R.F., Tully, J.G., Mouches, C., Saillard, C., Bove', J.M., Wroblewski, H., Carle, P., Rose, D.L., Henegar, R.B., Williamson, D.L., "*Spiroplasma melliferum*, a new species from the honeybee (*Apis mellifera*)," Int. J. Syst. Bacteriol., 35, 1985, pp. 296–308.
- [Corby-Harris, 2014] Corby-Harris, V., Maes, P., Anderson, K., "The bacterial communities associated with honey bee (*Apis mellifera*) foragers," PNAS, 2014, 9: e95056. doi:10.1371/journal.pone.0095056
- [Cox-Foster, 2007] Cox-Foster, D.L., Conlan, S., Holmes, E.C., Palacios, G., Evans, J.D., Moran, N.A., Quan, P.L., Brieseman, T., Hornig, M., Geiser, D.M., Martinson, V., Engelsdorp, D., Kalkstein, A.L., Drysdale, A., Hui, J., Zhai, J., Cui, L., Hutchison, S.K., Simons, F., Egholm, M., Lipkin, W. L., "A metagenomics survey of microbes in honey bee colony collapse disorder," Science, 318, 2007, pp. 283-287.
- [Crotti, 2010] Crotti, E., Rizzi, A., Chouaia, B., Ricci, I., Favia, G., Alma, A., Sacchi, L., Bourtzis, K., Mandrioli, M., Cherif, A., Bandi, C., Daffonchio, D., "Acetic acid bacteria, newly emerging symbionts of insects," Appl. Environ. Microbiol., 76, 2010, pp. 6963-6970.

- [Dade, 1977] Dade, H.A., "Anatomy and dissection of the honeybee," International Bee Research Association, London, UK., 1977.
- [Dillon, 2004] Dillon, R.J.V., Dillon, M., "The gut bacteria of insects: nonpathogenic interactions," *Annu. Rev. Entomol.*, 49, 2004, pp. 71-92.
- [Dillon, 2005] Dillon, R.J., Vennard, C.T., Buckling, A., Charnley, A.K., "Diversity of locust gut bacteria protects against pathogen invasion," *Ecol. Lett.*, 8, 2005, pp. 1291-1298.
- [Dillon, 2002] Dillon, R.J., Vennard, C.T., Charnley, A.K., "A note: gut bacteria produce components of a locust cohesion pheromone," *J. Appl. Microbiol.*, 92, 2002, pp. 759-763.
- [Disayathanoowat, 2012] Disayathanoowat, T., Young, J.P.W., Helgason, T., Chantawannakul, P., "T-RFLP analysis of bacterial communities in the midguts of *Apis mellifera* and *Apis cerana* honey bees in Thailand," *FEMS Microbiol. Ecol.*, 79, 2012, pp. 273-281.
- [Dyer, 1985] Dyer, F.C., "Nocturnal orientation by the Asian honey bee, *Apis dorsata*," *Anim. Behav.*, 33, 1985, pp. 769-774.
- [Endo, 2013] Endo, A., Salminen, S., "Honeybees and beehives are rich sources for fructophilic lactic acid bacteria," *Syst. Appl. Microbiol.*, 36, 2013, pp. 444-448.
- [Engel, 2013] Engel, P., Moran, N.A., "The gut microbiota of insects—diversity in structure and function," *FEMS Microbiol. Rev.*, 37, 2013, pp. 699-735.
- [Engel, 2012] Engel, P., Martison, V., Moran, N.A., "Functional diversity within the simple gut microbiota of the honey bee," *PNAS*, 109, 2012, pp. 11002-11007.

- [Evans, 2006] Evans, J., Armstrong, T.N., "Antagonistic interactions between honey bee bacterial symbionts and implications for disease," *BMC Ecol.*, 6, 2006, pp. 1-4.
- [Evans, 2004] Evans, J.D., Lopez, D.L., "Bacterial probiotics induce an immune response in the honey bee (hymenoptera: Apidae)," *J. Econ. Entomol.*, 97, 2004, pp. 752-756.
- [Evans, 2011] Evans, J.D., Schwarz, R.S., "Bees brought to their knees: microbes affecting honeybee health," *Trends Microbiol.*, 19, 2011, pp. 614-620.
- [Forsgren, 2010] Forsgren, E., "European foulbrood in honey bees," *J. Invertebr. Pathol.*, 103, 2010, pp. 55-59.
- [Forsgren, 2005] Forsgren, E., Lundhagen, A.C., Imdorf, A., Fries, I., "Distribution of *Melisococcus plutonius* in honeybee colonies with and without symptoms of European foulbrood," *Microb. Ecol.*, 50, 2005, pp. 369-374.
- [Fries, 1988] Fries, I., "Infectivity and multiplication of *Nosema apis* Z. in the ventriculus of the honey bee (*Apis mellifera* L.)," *Apidologie*, 19, 1988, pp. 319-328.
- [Foley, 2014] Foley, K., Fazio, G., Jensen, A., Hughes, W.O.H., "The distribution of *Aspergillus* spp. opportunistic parasites in hives and their pathogenicity to honey bees," *Vet. Microb.*, 169, 2014, pp. 203-210.
- [Fukuda, 1977] Fukuda, H., Ohtani, T., "Survival and life span of drone honeybees," *Res. Popul. Ecol.*, 19, 1977, pp. 51-68.

- [Genersch, 2006] Genersch, E., Forsgren, E., Pentikäinen, J., Ashiralieva, A., Rauch, S., Kilwinski, J., Fries, I., "Reclassification of *Paenibacillus larvae* subsp. *pulvifaciens* and *Paenibacillus larvae* subsp. *larvae* as *Paenibacillus larvae* without subspecies differentiation," Int. J. Syst. Evol. Microbiol., 56, 2006, pp. 501–511.
- [Gilliam, 1997] Gilliam, M., "Identification and roles of non-pathogenic microflora associated with honey bees," FEMS Microbiol. Lett., 155, 1997, pp. 1-10.
- [Gilliam, 1987] Gilliam, M., Prest, D.P., "Microbiology of feces of the larval honey bee, *Apis mellifera*," J. Invertebr. Pathol., 40, 1987, pp. 70–75.
- [Gilliam, 1988] Gilliam, M., Lorenz, B.J., Richardson, G.V., "Digestive enzymes and microorganisms in honeybee (*Apis mellifera*): influence of streptomycin, age, season, and pollen," Microbios, 55, 1988, pp. 95-114.
- [Hails, 2008] Hails, R.S., Ball, B.V., Genersch, E., "Infection strategies of insect viruses," Virology and the Honey Bee. European Commission, 2008, pp. 255-276.
- [Hammes, 2006] Hammes, W., Hertel, C., "The Genera *Lactobacillus* and *Carnobacterium*," The Prokaryotes, Springer-Verlag, 6, 2006, pp. 320-40.
- [Hansen, 1986] Hansen, H., Rasmussen, B., "The investigation of honey from bee colonies for *Bacillus larvae*," Dan J. Plant Soil Sci., 90, 1986, pp. 81-86.

- [Haynes, 2003] Haynes, S., Darby, A.C., Daniell, T.J., Webster, G., Veen, F.J.F., Godfray, H.C.J, Prosser, J.F., Douglas, A.E., "Diversity of bacteria associated with natural aphid populations," Appl. Environ. Microbiol., 69, 2003, pp. 7216-7223.
- [Higes, 2008] Higes, M., Martin-Hernandez, R., Garrido-Bailon, E., Garcia-Palencia, P., Meana, A., "Detection of infective *Nosema ceranae* (Microsporidia) spores in corbicular pollen of forager honeybees," J. Invertebr. Pathol., 97, 2008, pp. 76-78.
- [Hongoh, 2008] Hongoh, Y., Sharma, V.K., Prakash, T. et al., "Genome of an endosymbiont coupling N₂ fixation to cellulolysis within protist cells in termite gut," Science, 322, 2008, pp. 1108-1109.
- [Huang, 2013] Huang, S., Zhang, H., "The impact of environmental heterogeneity and life stage on the hindgut microbiota of *Holotrichia parallela* larvae (Coleoptera: Scarabaeidae)," PLOS ONE, 8, 2013, e57169.
- [Huang, 1996] Huang, Y.C., Edwards, C.G., Peterson, J.C., Haag, K.M., "Relationship between sluggish fermentations and the antagonism of yeast by lactic acid bacteria," Am. J. Enol.Vitic., 47, 1996, pp. 1-10.
- [Inglis, 1998] Inglis, G.D., Yanke, L.J., Goettel, M.S., "Anaerobic bacteria isolated from the alimentary canals of alfalfa leaf cutting bee larvae," Apidologie, 29, 1998, pp. 327-332.
- [Jeyaprakash, 2003] Jeyaprakash, A., Hoy, M.A., Allsopp, M.H., "Bacterial diversity in worker adults of *Apis mellifera* capensis and *Apis mellifera* scutellata (Insecta: Hymenoptera) assessed using 16S rRNA sequences," J. Invertebr. Pathol., 84, 2003, pp. 96-103.

- [Jojima, 2004] Jojima, Y., Mihara, Y., Suzuki, S., Yokozeki, K., Yamanaka, S., Fudou, R., “*Saccharibacter floricola* gen. nov., sp. nov., a novel osmophilic acetic acid bacterium isolated from pollen,” *Int. J. Syst. Evol. Microbiol.*, 54, 2004, pp. 2263-2267.
- [Kawasaki, 2011] Kawasaki, S., Kurosawa, K., Miyazaki, M., Sakamoto, M., Ohkuma, M., Niimura, Y., “*Lactobacillus ozensis* sp. nov., isolated from mountain flowers,” *Int. J. Syst. Evol. Microbiol.*, 61, 2011, pp. 2435-2438.
- [Kelber, 2006] Kelber, A., Warrant, E.J., Pfaff, M., Wallen, R., Theobald, J.C., Weislo, W.T., Raguso, R.A., “Light intensity limits foraging activity in nocturnal and crepuscular bees,” *Behav. Ecol.*, 17, 2006, pp. 63-72.
- [Kikuchi, 2012] Kikuchi, Y., Hayatsu, M., Hosokawa, T., Nagayama, A., Tago, K., Fukatsu, T., “Symbiont-mediated insecticide resistance,” *PNAS*, 109, 2012, pp. 8618-8622.
- [Kikuchi, 2007] Kikuchi, Y., Hosokawa, T., Fukatsu, T., “Insect-microbe mutualism without vertical transmission: a stinkbug acquires a beneficial gut symbiont from the environment every generation,” *Appl. Environ. Microbiol.*, 73, 2007, pp. 4308-4316.
- [Koch, 2011] Koch, H., Schmid-Hempel, P., “Socially transmitted gut microbiota protect bumble bees against an intestinal parasite,” *PNAS*, 108, 2011, pp. 19288-19292.
- [Koch, 2012] Koch, H., Schmid-Hempel, P., “Gut microbiota instead of host genotype drive the specificity in the interaction of a natural host-parasite system,” *Ecol. Lett.*, 15, 2012, pp. 1095-103.
- [Koch, 2013] Koch, H., Abrol, D.P., Li, J., Schmid-Hempel, P., “Diversity and evolutionary patterns of bacterial gut associates of corbiculate bees,” *Mol. Ecol.*, 22, 2013, pp. 2028-2044.

- [Kwong, 2013] Kwong, W., Moran, N., "Cultivation and characterization of the gut symbionts of honeybees and bumble bees: *Snodgrassella alvi* gen. nov., sp. nov., a member of the *Neisseriaceae* family of the *Betaproteobacteria*; and *Gilliamella apicola* gen. nov., sp. nov., a member of *Orbaceae* fam. nov., *Orbales* ord. nov., a sister taxon to the *Enterobacteriales* order of the *Gammaproteobacteria*," Int. J. Syst. Evol. Microbiol., 63, 2013, pp. 2008-2018.
- [Lane, 1991] Lane, D.J., "16S/23S rRNA sequencing," Nucleic acid techniques in bacterial systematics. John Wiley & Sons Ltd, 1991, pp. 115-147.
- [Lee, 2014] Lee, F.J., Rusch, D.B., Stewart, F.J., Mattila, H.R., Newton, I.L., "Saccharide breakdown and fermentation by the honey bee gut microbiome," Environ. Microbiol., 17, 2014, pp. 796-815.
- [Leroy, 2011] Leroy, P.D., Sabri, A., Heuskin, S., Thonart, P., Lognay, G., Verheggen, F.J., Francis, F., Brostaux, Y., Felton, G.W., Haubruge, E., "Microorganisms from aphid honeydew attract and enhance the efficacy of natural enemies," Nat. Commun., 2, 2011, pp. 348.
- [Li, 2007] Li, Y., Ge, Y., Saxena, D., Caufield, P.W., "Genetic profiling of the oral microbiota associated with severe early-childhood caries," J. Clin. Microbiol., 45, 2007, pp. 81-87.
- [Maggi, 2013] Maggi, M., Negri, P., Plischuk, S., Szawarski, N., Piano, F.D., Feudis, L.D., Eguaras, M., Audisio, C., "Effects of the organic acids produced by a lactic acid bacterium in *Apis mellifera* colony development, *Nosema ceranae* control and fumagillin efficiency," Vet. Microbiol., 176, 2013, pp. 474-483.

- [Martín-Hernández, 2011] Martín-Hernández, R., Botías, C., Barrios, L., Martínez-Salvador, A., Meana, A., Mayack, C., Higes, M., “Comparison of the energetic stress associated with experimental *Nosema ceranae* and *Nosema apis* infection of honeybees (*Apis mellifera*),” *Parasitol. Res.*, 109, 2011, pp. 605–612.
- [Maori, 2007] Maori, E., Tanne, E., Sela, I., “Reciprocal sequence exchange between non-retro viruses and hosts leading to the appearance of new host phenotypes,” *Virology*, 362, 2007, pp. 342–349.
- [Martinson, 2011] Martinson, V.G., Danforth, B.N., Minckley, R.L., “A simple and distinctive microbiota associated with honeybees and bumble bees,” *Mol. Ecol.*, 20, 2011, pp. 619-628.
- [Martinson, 2012] Martinson, V.G., Moy, J., Moran, N., “Establishment of characteristic gut bacteria during development of the honeybee worker,” *Appl. Environ. Microbiol.*, 78, 2012, pp. 2830-2840.
- [Matheson, 1993] Matheson, A., “World bee health report,” *Bee World*, 74, 1993, pp. 176-212.
- [McFrederick, 2014] McFrederick, Q., Wcislo, W.T., Hout, M.C., Mueller, U.G., “Host species and developmental stage, but not host social structure, affects bacterial community structure in socially polymorphic bees,” *FEMS Microbiol. Ecol.*, 88, 2014, pp. 1-10.
- [McFrederick, 2013] McFrederick, Q.S., Cannone, J.J., Gutell, R.R., Kellner, K., Plowes, R.M., Mueller, U.G., “Specificity between lactobacilli and hymenopteran hosts is the exception rather than the rule,” *Appl. Environ. Microbiol.*, 79, 2013, pp. 1803-1812.
- [Michener, 1974] Michener, C.D., “The Social Behavior of the Bees: A Comparative Study,” Harvard University Press, 1994.

- [Minnick, 2009] Minnick, M.F., Battisti, J.M., “Pestilence, persistence and pathogenicity: infection strategies of *Bartonella*,” *Future Microbiol.*, 4, 2009, pp. 743-758.
- [Möckel, 2011] Möckel, N., Gisder, S., Genersch, E., “Horizontal transmission of deformed wing virus: pathological consequences in adult bees (*Apis mellifera*) depend on the transmission route,” *J. Gen. Virol.*, 92, 2011, pp. 370-377.
- [Mohr, 2007] Mohr, K.I., Tebbe, C.C., “Field study results on the probability and risk of horizontal gene transfer from transgenic herbicide-resistant oil seed rape pollen to gut bacteria of bees,” *Appl. Environ. Microbiol.*, 75, 2007, pp. 573-582.
- [Moore, 1989] Moore, D., Siefried, D., Wilson, R., Rankin, M.A., “The influence of time of day on the foraging behavior on the honeybee, *Apis mellifera*,” *J. Biol. Rhythms*, 4, 1989, pp. 305-325.
- [Morales-Jiménez, 2012] Morales-Jiménez, J., Zúñiga, G., Ramírez-Saad, H.C. and Hernández Rodríguez, C., “Gut-associated bacteria throughout the life cycle of the bark beetle *Dendroctonus rhizophagus* Thomas and Bright (Curculionidae: Scolytinae) and their cellulolytic activities,” *Microb. Ecol.*, 64, 2012, pp. 268-278.
- [Moran, 2012] Moran, N.A., Hansen, A.K., Powell, J.E., Sabree, Z.L., “Distinctive gut microbiota of honeybees assessed using deep sampling from individual worker bees,” *PLOS ONE*, 7, 2012, e36393.
- [Mouches, 1983] Mouches, C., Bove, J.M., Tully, J.G., Rose, D.L., McCoy, R.E., Carlejunca, P., Garnier, M., Saillard, C., “*Spiroplasma apis*, a new species from the honeybee *Apis mellifera*,” *Annu. Microbiol.*, 134, 1983, pp. 383–397.

- [Nakano, 1990] Nakano, H., Okabe, T., Hashimoto, H., Sakaguchi, G., "Incidence of *Clostridium botulinum* honey of various origins," Jpn. J. Med. Sci. Biol., 43, 1990, pp. 183-195.
- [Nakatsu, 2000] Nakatsu, C.H., Torsvik, V., Øvreas, L., "Soil community analysis using DGGE of 16S rDNA polymerase chain reaction products," Soil. Sci. Soc. Am. J., 64, 2000, pp. 1382-1388.
- [Neveling, 2012] Neveling, D.P., Endo, A., Dicks, L.M.T., "Fructophilic *Lactobacillus kunkeei* and *Lactobacillus brevis* isolated from fresh flowers, bees and bee-hives," Curr.Microbiol., 65, 2012, pp. 507-515.
- [Newton, 2012] Newton, I., Roeselers, G., "The effect of training set on the classification of honey bee gut microbiota using the Naïve Bayesian classifier," BMC Microbial., 12, 2012, pp. 221-230.
- [Nihoh, 2011] Nihoh, N., Hosokawa, T., Oshima, K., Hattori, M., Fukatsu, T., "Reductive evolution of bacteria genome in insect gut environment," Genome Biol. Evol., 3, 2011, pp. 702-714.
- [Oldroyd, 2006] Oldroyd, B.P., Wongsiri, S., "Asian Honey Bees," Harvard University Press, 2006.
- [Olofsson, 2008] Olofsson, T., Vásquez, A., "Detection and identification of a novel lactic acid bacterial flora within the honey stomach of the honeybee *Apis mellifera*," Curr. Microbiol., 57, 2008, pp. 356-363.
- [Otis, 1991] Otis, G.W., 'Revised distribution of three recently recognized species of honey bees in Asia,' Honeybee Science, 15, 1991, pp. 167-170.

- [Potts, 2010] Potts, S.G., Biesmeijer, J.C., Kremen, C., Neumann, P., Schweiger, O., Kunin, W.E., "Global pollinator declines: trends, impacts and drivers," Trends Ecol. Evol., 25, 2010, pp. 345-353.
- [Powell, 2014] Powell, J.E., Martinson, V.G., Urban-Mead, K., Moran, N.A., "Routes of acquisition of the gut microbiota of *Apis mellifera*," Appl. Environ. Microbiol., 2014, (in press).
- [Promnuan, 2009] Promnuan, Y., Kudo, T., Chantawannakul, P., "Actinomycetes isolated from beehives in Thailand," World J. Microbiol. Biotechnol., 25, 2009, pp. 1685-1689.
- [Promnuan, 2011] Promnuan, Y., Kudo, T., Ohkuma, M., Chantawannakul, P., "*Actinomaduraapis* sp. nov., isolated from a honey bee (*Apis mellifera*) hive, and the reclassification of *Actinomadura cremea* subsp. *rifamycini* Gauze et al. 1987 as *Actinomadura rifamycini* (Gauze et al. 1987) sp. nov., comb. nov.," Int. J. Syst. Evol. Microbiol., 61, 2011, pp. 2271-2277.
- [Pumpuni, 1993] Pumpuni, C.B., Beier, M.S., Nataro, J.P., Guers, L.D., Davis, J.R., "*Plasmodium falciparum*: inhibition of sporogonic development in *Anopheles stephensi* by gram-negative bacteria," Exp. Parasitol., 77, 1993, pp. 195-199.
- [Rinderer, 1996] Rinderer, T.E., Wongsiri, S., Kuang, B.Y., Liu, J.S., Oldroyd, B.P., Sylvester, H.A., de Guzman, L.I., "Comparative nest architecture of the dwarf honey bees," J. Apicult. Res., 35, 1996, pp. 19-26.
- [Robinson, 2010] Robinson, C.J., Schloss, P., Ramos, Y., Raffa, K., Handelsman, J., "Robustness of the bacterial community in the cabbage white butterfly larval midgut," Microb. Ecol., 59, 2010, pp. 199-211.

- [Ruttner, 1988] Ruttner, F., "Biogeography and taxonomy of honeybees," Springer, Berlin, 1988.
- [Sabree, 2012] Sabree, Z.L., Hansen, A.K., Moran, N.A., "Independent studies using deep sequencing resolve the same set of core bacterial species dominating gut communities of honey bees," PLOS ONE, 7(7), 2012, e41250.
- [Sachs, 2011] Sachs, J., Skophammer, R.G., Regus, J.U., "Evolutionary transitions in bacterial symbiosis," PNAS, 108, 2011, pp. 10800-10807.
- [Schaffer, 1979] Schaffer, W.M., Jensen, D.B., Hobbs, D.E., "Competition, foraging energetics and the cost of sociality in three species of bees," Ecology, 60, 1979, pp. 976-987.
- [Seeley, 1978] Seeley, T.D., "Life history strategy of honey bee, *Apis mellifera*," Oecology, 32, 1978, pp.109-118.
- [Seeley, 1985] Seeley, T.D., "The information-center strategy of honeybee foraging," Fortschritte der Zoologie, 31, 1985, pp. 75-90.
- [Seeley, 1995] Seeley, T.D., "The wisdom of the hive: The social physiology of honey bee colonies," Harvard University Press, 1995.
- [Sharma, 2005] Sharma, A.D., Singh, J., "A nonenzymatic method to isolate genomics DNA from bacteria and actinomycetes," Annu. Rev. Biochem., 337, 2005, pp. 354-356.
- [Simpson, 1999] Simpson, J.M., McCracken, V.J., White, B.A., Gaskins, H.R., Mackie, R.I., "Application of denaturant gradient gel electrophoresis for the analysis of the porcine gastrointestinal microbiota," J. Microbiol. Methods., 36, 1999, pp. 167-179.

- [Soman, 1996] Soman, A.G. and Chawda, S.S., "A contribution to the biology and behavior of the dwarf bee, *A. florea* F. and its economic importance in Kutch, Gujarat, India," Indian Bee J., 58, 1996, pp. 81-88.
- [Somanathan, 2009] Somanathan, H., warrant, E.J., Borges, R.M., Wallen, R., Kelber, A., "Resolution and sensitivity of the eyes of the Asian honeybees *Apis florea*, *Apis cerana* and *Apis dorsata*," J. Exp. Biol., 212, 2009, pp. 2448-2453.
- [Storelli, 2011] Storelli, G., Defaye, A., Erkosar, B., Hols, p., Royet, J., Leulier, F., "*Lactobacillus plantarum* promotes *Drosophila* systemic growth by modulating hormonal signals through TOR-dependent nutrient sensing," Cell Metab., 14, 2011, pp. 403-414.
- [Tajabadi, 2011] Tajabadi, N., Mardan, M., Abdul Manap, M.Y., Shuhaimi, M., Meimandipour, A., Nateghi, L., "Detection and identification of *Lactobacillus* bacteria found in the honey stomach of the giant honeybee *Apis dorsata*," Apidologie, 42, 2011, pp. 642-649.
- [Tanaka, 2006] Tanaka, H., Aoyagi, H., Shina, S., Dodo, Y., Yoshimura, T., Nakamura, R., Uchiyama, H., "Influence of the diet components on the symbiotic microorganisms community in hindgut of *Coptotermes formosanus* Shiraki," Appl. Environ. Microbiol., 71, 2006, pp. 907-917.
- [Theantana, 2008] Theantana, T., Chantawannakul, P., "Protease and β -N-acetylglucosaminidase of honey bee chalkbrood pathogen *Ascosphaera apis*," J. Apicult. Res., 47, 2008, pp. 68-76.

- [Vasanthakuma, 2006] Vasanthakuma, A., Delalibera, I., Handelsman, J., Klepzig, K.D., Schloss, P.D., Raffa, K.F., "Characterization of gut associated bacteria in larva and adults of the Southern pine beetle, *Dendroctonus frontaliszimmermann*," *Mol. Biol. Evol.*, 35, 2006, pp. 1710-1717.
- [Vasquez, 2012] Vasquez, A., Forsgren, E., Fries, I., Paxton, R.J., Flaberg, E., Szekely, L. and Olofsson, T.C., "Symbionts as major modulators of insect health: lactic acid bacteria and honeybees," *PLOS ONE*, 7, 2012, e33188.
- [Vojvodic, 2013] Vojvodic, S., Rehan, S.M., Anderson, K.E., "Microbial gut diversity of Africanized and European honeybee larval instars," *PLOS ONE*, 8(8), 2013, e72106.
- [Warnecke, 2007] Warnecke, F., Luginbühl, P., Ivanova, N. et al., "Metagenomic and functional analysis of hindgut microbiota of a wood-feeding higher termite," *Nature*, 450, 2007, pp. 560-565.
- [Winston, 1987] Winston, M.L., "The Biology of the Honey Bee," Harvard University Press, 1987.
- [Woyke, 2003] Woyke, J., Wilde, J., Wild, M., "Flight activity reaction to temperature changes in *Apis dorsata*, *Apis laboriosa* and *Apis mellifera*," *J. Apic. Res.*, 47, 2003, pp. 73-80.
- [Wynns, 2013] Wynns, A.A., Jensen, A.B., Eilenberg, J., "Ascosphaera callicarpa, a new species of bee-loving fungus, with a key to the genus for Europe," *PLOS ONE*, 8(9), 2013, e73419.
- [Yang, 2007] Yang, X., Cox-Foster, D., "Effects of parasitization by *Varroa destructor* on survivorship and physiological traits of *Apis mellifera* in correlation with viral incidence and microbial challenge," *Parasitology*, 134, 2007, pp. 405-412.

- [Yoshiyama, 2009] Yoshiyama, M., Kimura, K., "Bacteria in the gut of Japanese honey bee, *Apis cerana japonica*, and their antagonistic effect against *Paenibacillus larvae*, the causal agent of American foulbrood," J. Invertebr. Pathol., 102, 2009, pp. 91-96.
- [Yue, 2005] Yue, C., Genersch, E., "RT-PCR analysis of deformed wing virus in honeybees (*Apis mellifera*) and mites (*Varroa destructor*)," J. Gen. Virol., 86, 2005, pp. 3419-24.
- [Yue, 2006] Yue, C., Schröder, M., Bienefeld, K., Genersch, E., "Detection of viral sequences in semen of honeybees (*Apis mellifera*): evidence for vertical transmission of viruses through drones," J. Invertebr. Pathol., 92, 2006, pp. 105-108.
- [Yue, 2007] Yue, C., Schröder, M., Gisder, S., Genersch, E., "Vertical-transmission routes for deformed wing virus of honeybees (*Apis mellifera*)," J. Gen. Virol., 88, 2007, pp. 2329-36.
- [Zhang, 2012] Zhang, X., He, S.Y., Evans, J.D., Pettis, J.S., Yin, G.F., Chen, Y.P., "New evidence that deformed wing virus and black queen cell virus are multi-host pathogens," J. Invertebr. Pathol., 109, 2012, pp. 156-159.
- [Zhang, 2008] Zhang, H., Jackson, T.A., "Autochthonous bacterial flora indicated by PCR-DGGE of 16S rRNA gene fragments from the alimentary tract of *Costelytra zealandica* (Coleoptera: Scarabaeidae)." J. Appl. Microbiol., 105, 2008, pp. 1277-1285.