

REFERENCES

- Abbagani, S., Gundimeda, S. D., Varre, S., Ponnala, D., & Mundluru, H. P. (2010). Kidney stone disease: Etiology and evaluation. *International Journal of Applied Biology and Pharmaceutical Technology*, 1(1), 175-182. Retrieved from www.ijabpt.com
- Agarwal, M. M., Singh, S. K., Mavuduru, R., & Mandal, A. K. (2011). Preventive fluid and dietary therapy for urolithiasis: An appraisal of strength, controversies and lacunae of current literature. *Indian Journal of Urology*, 27(3), 310-319. doi:10.4103/0970-1591.85423
- Agondi, Rde, F., Gallani, M. C., B. J. Rodrigues, R. C. M., & Cornelio, M. E. (2011). Relationship between beliefs regarding a low salt diet in chronic renal failure patients on dialysis. *Journal of Renal Nutrition*, 21(2), 160-168. doi:10.1053/j.jrn.2010.03.007
- Al Zahrani, H., Norman, R. W., Thompson, C., & Weerasinghe, S. (2000). The dietary habits of idiopathic calcium stone-formers and normal control subjects. *British Journal of Urology International*, 85(6), 616-620. doi:10.1046/j.1464-410x.2000.00511.x
- Al-Eisa, A. A., Al-Hunayyan, A., & Gupta R. (2002). Pediatric urolithiasis in Kuwait. *International Urology and Nephrology*, 33(1), 3-6. doi:10.1023/A:1014419830292
- Ansari, M. S., & Gupta, N. P. (2003). Impact of socioeconomic status in etiology and management of urinary stone disease. *Urologia Internationalist*, 70(4), 255-61. doi:10.1159/000070130
- Awasthi, M., Malhotra, S. R., & Modgil, R. (2011). Dietary habits of kidney stone patients of Kangra District, Himachal Pradesh, North India. *Journal of Human Ecology*, 34(3), 163-169. Retrieved from <http://www.krepublishers.com/>

- Baheiraei, A., Mirghafourvand, M., Charandabi, S. M. A., & Mohammadi, E. (2013). Facilitators and inhibitors of health-promoting behaviors: The experience of Iranian women of reproductive age. *International Journal of Preventive Medicine*, 4(8), 929-939. Retrieved from <http://www.ncbi.nlm.nih.gov/>
- Bakunts., V. (2011). *Knowledge, attitude and practice of kidney stone formers in Armenia Regarding prevention of kidney stone disease* (Master's thesis). Retrieved from <http://auachsr.com/>
- Baldwin, S. A. S. (2014). *Coronary heart disease knowledge, health promoting behaviors and perceived benefits and barriers to exercise and healthy eating in Montana females* (Master's thesis). Retrieved from <http://scholarworks.montana.edu/>
- Barnela, S. R., Soni, S. S., Saboo, S. S., & Bhansali, A. S. (2012). Medical management of renal stone. *Indian Journal of Endocrinology Metabolism*, 16(2), 236-239. doi:10.4103/2230-8210.93741
- Bartoletti, R., & Cai, T. (2008). Surgical approach to urolithiasis: The state of art. *Clinical Cases in Mineral and Bone Metabolism*, 5(2), 142-144. Retrieved from <http://www.ncbi.nlm.nih.gov/>
- Bartoletti, R., Cai, T., Mondaini, N., Melone, F., Travaglini, F., Carini, M., & Rizzo, M. (2007). Epidemiology and risk factors in urolithiasis. *Urology Internationalis*, 79(1), 3-7. doi:10.1159/000104434
- Borghi, L., Meschi, T., Amato, F., Briganti, A., Novarini, A., & Giannini, A. (1996). Urinary volume, water and recurrences in idiopathic calcium nephrolithiasis: A 5-year randomized prospective study. *Journal of Urology*, 155(3), 839-843. doi:10.1016/S0022-5347(01)66321-3
- Borghi, L., Meschi, T., Schianchi, T., Briganti, A., Guerra, A., Allegri, F., & Novarini, A. (1999). Urine volume: Stone risk factor and preventive measure. *Nephron*, 81(1), 31-37. doi:10.1159/000046296

- Borghi, L., Schianchi, T., Meschi, T., Guerra, A., Allegri, F., Maggiore, U., & Novarini, A. (2002). Comparison of two diets for the prevention of recurrent stones in idiopathic hypercalciuria. *New England Journal of Medicine*, 346(2), 77-84. doi:10.1056/NEJMoa010369
- Boyd, K. M. (2000). Disease, illness, sickness, health, healing and wholeness: Exploring some elusive concepts. *Journal of Medical Ethics: Medical Humanities*, 26, 9-17. doi:10.1136/mh.26.1.9
- Brislin, R. W., Lonner, W. J., & Thorndike, R. M. (1973). *Psychology*. Canada: John Wiley & Sons.
- Cheng, P. M., Moin, P., Dunn, M. D., Boswell, W. D., & Duddalwar, V. A. (2012). What the radiologist needs to know about urolithiasis: Part 1--Pathogenesis, types, assessment, and variant anatomy. *American Journal of Roentgenology*, 198(6), 540-547. doi:10.2214/AJR.10.7285
- Culhane-Pera, K., & Lee, M. (2006). 'Die another day': A qualitative analysis of Hmong experiences with kidney stones. *Hmong Studies Journal*, 7, 1-34. Retrieved from <http://www.hmongstudies.org/>
- Curhan, G. C. (2007). Epidemiology of stone disease. *Urologic Clinics of North America*, 34(3), 287-293. doi:10.1016/j.ucl.2007.04.003
- Curhan, G. C., Willett, W. C., Knight, E. L., & Stampfer, M. J. (2004). Dietary factors and the risk of incident kidney stones in younger women. Nurses' Health Study II. *Archives of Internal Medicine*, 164(8), 885-891. doi:10.1001/archinte.164.8.885.
- Curhan, G. C., Willett, W. C., Rimm, E. B., & Stampfer, M. J. (1993). A prospective study of dietary calcium and other nutrients and the risk of symptomatic kidney stones. *New England Journal of Medicine*, 328(12), 833-838. doi:10.1056/NEJM199303253281203

- Curhan, G. C., Willett, W. C., Rimm, E. B., & Stampfer, M. J. (1997). Family history and risk of kidney stones. *Journal of the American Society of Nephrology*, 8(10), 1568-1573. Retrieved from <http://jasn.asnjournals.org/>
- Curhan, G. C., Willett, W. C., Rimm, E. B., Spiegelman, D., & Stampfer, M. J. (1996). Prospective study of beverage use and the risk of kidney stones. *American Journal of Epidemiology*, 143(3), 240-247. Retrieved from <http://aje.oxfordjournals.org/>
- Curhan, G. C., Willett, W. C., Speizer, F. E., & Stampfer, M. J. (1998). Beverage use and risk for kidney stones in women. *Annals Internal Medicine*, 128(7), 534-540. doi:10.7326/0003-4819-128-7-199804010-00003
- Curhan, G. C., Willett, W. C., Speizer, F. E., Spiegelman, D., & Stampfer, M. J. (1997). Comparison of dietary calcium with supplemental calcium and other nutrients as factors affecting the risk for kidney stones in women. *Annals of Internal Medicine*, 126, 497-504. doi:10.7326/0003-4819-126-7-199704010-00001
- Dai, M., Zhao, A., Liu, A., You, L., & Wang, P. (2013). Dietary factors and risk of kidney stone: A case-control study in Southern China. *Journal of Renal Nutrition*, 23(2), 21-28. doi:10.1053/j.jrn.2012.04.003
- Damasio, P. C., Amaro, C. R., Padovani, C. R., Leitao, V. A., & Yamamoto, H. (2014). Influence of clinical therapy and nutritional counseling on the recurrence of urolithiasis. *400 - Acta Cirúrgica Brasileira*, 29(6). doi:<http://dx.doi.org/10.1590/S0102-86502014000600009>
- Daudon, M., Traxer, O., Conort, P., Lacour, B., & Jungers, P. (2006). Type 2 diabetes increases the risk for uric acid stones. *Journal of American Society of Nephrology*, 17(7), 2026-2033. doi:10.1681/ASN.2006030262
- Dawson, C. H. (2012). Kidney stone disease: pathophysiology, investigation and medical treatment. *Clinical Medicine*, 12(5), 467-471. doi:10.7861/clinmedicine.12-5-467

- D'Cruz, F. & Chandrasekharan, N. (1990). Renal disease in Malaysia: problem and prospects. *Medical Journal Malaysia*, 45(2). Retrieved from <http://www.emjm.org/>
- Dennison, D., Mayo, M. L., & Abraham, V. E. (2011). The development of kidney stone dietary plans for patient education. *International Electronic Journal of Health Education*, 14, 150-156. Retrieved from <http://eric.ed.gov/>
- DeVellis, R. F. (2003). *Scale development: Theory and application* (2nd ed.). Thousand Oaks, CA: Sage.
- Downey, P., & Tolley, D. (2000). Contemporary management of renal calculus disease. *Journal of the Royal College Surgical (Edinburgh)*, 47(5), 668-675. Retrieved from www.rcsed.ac.uk/
- Edvardsson, V. O., Indridason, O. S., Haraldsson, G., Kjartansson, O., & Palsson, R. (2013). Temporal trends in the incidence of kidney stone disease. *Kidney International*, 83, 146-152. doi:10.1038/ki.2012.320
- Ferraro, P. M., Taylor, E. N., Gambaro, G., & Curhan, G. C. (2013). Soda and other beverages and the risk of kidney stones. *Clinical Journal of the American Society of Nephrology*, 8(8), 1-7. doi:10.2215/CJN.11661112
- Finch W. J. G., & Irving, S. O. (2007). Role of diet in the prevention of common kidney stones. *Trends in Urology Gynaecology and Sexual Health*, 12(5), 8-9. doi:10.1002/tre.36
- Finkielstein, V. A., & Goldfarb, D. S. (2006). Strategies for preventing calcium oxalate stones. *Canadian Medical Association Journal*, 174(10), 1407-1409. doi:10.1509/cmaj.051517
- Flagg, L. R. (2007). Dietary and holistic treatment of recurrent calcium oxalate kidney stones: Review of literature to guide patient education. *Urologic Nursing*, 27(2), 113-122. Retrieved from <https://www.cbuna.org/>

- Fritsche, H. M., & Dotzer, K. (2012). Improving the compliance of the recurrence stone-former. *Arab Journal of Urology*, 10(3), 342-246.
doi:10.1016/j.aju.2012.07.003
- Furman, E. F. (2012). *The theory of compromised eating behavior* (Doctoral dissertation). Retrieved from <http://scholarworks.umass.edu>
- Gans, K. M., Risica, P. M., Wylie-Rosett, J., Ross, E. M., Strolla, L. O., McMurray, J., & Eaton, C. B. (2006). Development and evaluation of the nutrition component of the rapid eating and activity assessment for patients (REAP): A new tool for primary care providers. *Journal of Nutrition Education and Behavior*, 38(5), 286-292. doi:10.1016/j.jneb.2005.12.002
- Goel, R. (2012). Kidney stones: Diagnostic and treatment strategies. *Consultant* 360, 52(2). Retrieved from <http://www.consultant360.com/>
- Goldfarb, D. S., & Coe, F. L. (1999). Beverages, diet, and prevention of kidney stones. *American Journal of Kidney Diseases*, 33(2), 398-400. doi:10.1016/S0272-6386(99)70319-7
- Goldfarb, D., Trinchieri, A., & Curhan, G. (2007). Epidemiology of stone disease. In J. Denstedt & S. Khoury (Eds.), *The 2nd International Consultation on Stone Disease* (pp. 21-30). Retrieved from <http://www.icud.info>
- Grace, P. A., & Borley, N. R. (2013). *Surgery at a glance*. Chichester: Wiley-Blackwell.
- Gravina, G. L., Costa, A. M., Ronchi, P., Angelucci, A., Castellani, D., Narcisi, F., & Vicentini, C. (2005). Tamsulosin treatment increase clinical success rate of single extracorporeal shock wave lithotripsy of renal stones. *Urology*, 66(1), 24-28.
doi:10.1016/j.urology.2005.01.013

- Grove, S. K., Burns, N., & Gray, J. R. (2012). *The practice of nursing research: Appraisal, synthesis and generation of evidence* (7th ed.). Missouri: Elsevier Saunders.
- Gul, Z., & Monga, M. (2014). Medical and dietary therapy for kidney stone prevention. *Korean Journal of Urology*, 55(12), 775-779. doi:10.4111/kju.2014.55.12.775
- Hall, P. M. (2002). Preventing kidney stones: Calcium restriction not warranted. *Cleveland Clinic Journal of Medicine*, 69(11), 885-888. Retrieved from <http://www.ccjm.org/>
- Hamano, S., Nakatsu, H., Suzuki, N., Tomioka, S., Tanaka, M., & Murakami, S. (2005). Kidney stone disease and risk factors for coronary heart disease. *International Journal of Urology*, 12(10), 859-863. Retrieved from <http://onlinelibrary.wiley.com/>
- Hanson, K. (2005). Minimally invasive and surgical management of urinary stones. *Urologic Nursing*, 25(6), 458-464. Retrieved from <http://www.medscape.com/>
- Hassapidou, M. N., Paraskevopoulos, S. Th., Karakoltsidis, P. A., Petridis, D., & Fotiadou, E. (1999). Dietary habits of patients with renal stone disease in Greece. *Journal of Human Nutrition and Dietetics*, 12(1), 47-51. doi:10.1046/j.1365-277x.1999.00141.x
- Heilberg, I. P., & Goldfarb, D. S. (2013). Optimum nutrition for kidney stone disease. *Advances in Chronic Kidney Disease*, 20(2), 165-174. doi:10.1053/j.ackd.2012.12.001
- Heller, H. J., Sakhaee, K., Moe, O. W., & Pak, C. Y. (2002). Etiological role of estrogen status in renal stone formation. *The Journal of Urology*, 168(5), 1923-1927. doi:10.1097/00005392-200211000-00007
- Hess, B. (2012). Metabolic syndrome, obesity and kidney stones. *Arab Journal of Urology*, 10(3), 258-264. doi:10.1016/j.aju.2012.04.005

- Hesse, A., Brändle, E., Wilbert, D., Köhrmann, K. U., & Alken, P. (2003). Study on the prevalence and incidence of urolithiasis in Germany comparing the years 1979 vs. 2000. *European Urology*, *44*(6), 709-713. doi:10.1016/S0302-2838(03)00415-9
- Hesse, A., & Siener, R. (1997) Current aspects of epidemiology and nutrition in urinary stone disease. *World Journal of Urology*, *15*(3), 165-171. doi:10.1007/BF02201853
- Hiatt, R. A., Ettinger, B., Caan, B., Quesenberry, C. P., Duncan, D., & Citron, J. T. (1996). Randomized controlled trial of a low animal protein, high fiber diet in the prevention of recurrent of calcium oxalate kidney stones. *American Journal of Epidemiology*, *44*(1), 25-33. Retrieved from: <http://aje.oxfordjournals.org/>
- Holmes, R. P., Goodman, H. O., & Assimos, D. G. (2001). Contribution of dietary oxalate to urinary oxalate excretion. *Kidney International*, *59*, 270-276. doi:10.1046/j.1523-1755.2001.00488.x
- Hong, Y. H., Dublin, N., Razack, A. H., Mohd, M. A., & Husain, R. (2012). Urinary metabolic evaluation of stone formers—A Malaysian perspective. *Urology*, *80*(3), 529-534. doi:10.1016/j.urology.2012.02.053
- Hsu, T. C., Chen, J., Huang, H. S., & Wang, C. J. (2002). Association of changes in the pattern of urinary calculi in taiwanese with diet habit change between 1956 and 1999. *Journal of the Formosan Medical Association*, *101*(1), 5-10. Retrieved from <http://www.fma.org.tw/>
- Huether, S. E., & McCance K. L. (2013). *Understanding pathophysiology*. Retrieved from <https://books.google.co.th/>
- Hughes, K. E., & Arthur, J. (2013). An atypical case of large bowel obstruction. *Journal of Surgical Case Report*, *2013*(6). doi:10.1093/jscr/rjt042

- Hussein, N. S., Sadiq, S. M., Kamaliah, M. D., Norakmal, A. W., & Gohar, M. N. (2013). Twenty-four-hour urine constituents in stone formers: A study from the northeast part of Peninsular Malaysia. *Saudi Journal of Kidney Disease Transplant*, 24(3), 630-637. Retrieved from <http://www.sjkdt.org/>
- International Osteoporosis Foundation. (2015). *Calcium content of common foods*. Retrieved from <http://www.iofbonehealth.org/>
- Ismawati., S., Zainalabidin, M., & Golnaz, R. (2014). Healthy eating: the preventive factors among Malaysians. *Journal of Economic, Business and Management*, 1(4), 257-261. doi:10.7763/JOEBM.2014.V2.135
- Izamin, I., Aniza, I., Rizal, A. M., & Aljunid, S. M. (2009). Comparing extracorporeal shock wave lithotripsy and ureteroscopy for treatment of proximal ureteric calculi: a cost effectiveness study. *Medical Journal of Malaysia*, 64(1), 12-21. Retrieved from <https://www.researchgate.net/>
- Jabatan Penerangan Malaysia. (2013). *Perangkaan pendapatan dan kemiskinan isi rumah*. Retrieved from <http://www.penerangan.gov.my/>
- Janz, N. K., & Becker, M. H. (1984). The health belief model: A decade later. *Health Education Quarterly*, 11(1), 1-47. Retrieved from <http://deepblue.lib.umich.edu/>
- Jayaraman, K., Munira, H., Dababrata Chowdhury & Iranmanesh, M. (2013). The preference and consumption of chicken lovers with race as a moderator—An empirical study in Malaysia. *International Food Research Journal*, 20(1), 165-174. Retrieved from <http://www.ifrj.upm.edu.my>
- Johri, N., Jaeger, P., Robertson, W., Choong, S., & Unwin, R. (2011). Renal stone disease. *Medicine*, 39(7), 371-377. doi:10.1016/j.mpmed.2011.04.005

- Kalaitzidis, R. G., Damigos, D., & Siamopoulos, K. C. (2014). Environmental and stressful factors affecting the occurrence of kidney stones and the kidney colic. *International Journal of Urology and Nephrology*, 46(9), 1779-1784. doi:10.1007/s11255-014-0758-2
- Kambadakone, A. R., Eisner, B. H., Catalano, O. A., & Sahani, D. V. (2010). New and evolving concepts in the imaging and management of urolithiasis: Urologists' perspective. *Radiographics*, 30(3), 603-623. doi:10.1148/rg.303095146
- Karen, G. & Jay, M. (2002). *Behavior, health related*. Encyclopedia of Public Health. Retrieved from www.encyclopedia.com
- Kasl, S. V., & Cobb, S. (1966). Health behavior, illness behavior and sick role behavior. *Archives of Environmental Health: An International Journal*, 12(4), 531-541. doi:10.1080/00039896.1966.10664421
- Kirk, R. E. (2007). *Statistics: An introduction* (5th ed.). Belmont, CA: Thomson Higher Education.
- Kloeblen, A. S., & Batish, S. S. (1999). Understanding the intention to permanently follow a high folate diet among a sample of low income pregnant women according to the health belief model. *Health Education Research*, 14(3), 327-338. doi:10.1093/her/14.3.327
- Knoll, T. (2008). Epidemiology, pathogenesis and pathophysiology of urolithiasis. *European Urology Supplements*, 9(12), 802-806. doi:10.1016/j.eursup.2010.11.006
- Kok, D. J. (2012). Metaphylaxis, diet and lifestyle in stone disease. *Arab Journal of Urology*, 10(3), 240-249. doi:10.1016/j.aju.2012.03.003
- Kosar, A., Sarica, K., Aydos, K., Kupeli, S., Turkolmez, K., & Gogus, O. (1999). Comparative of long-term stone recurrence after extracorporeal shock wave lithotripsy and open stone surgery for kidney stones. *International Journal of Urology*, 6(3), 125-129. doi:10.1046/j.1442-2042.1999.06311.x

- Koyuncu, H. H., Yencilek, F., Eryildirim, B., & Sarica, K. (2010). Family history in stone disease: How important is it for the onset of the disease and the incidence of recurrence? *Urological Research*, 38(2), 105-109. doi:10.1007/s00240-009-0249-6
- Krieger, J. N., Kronmal, R. A., Coxon, V., Wortley, P., Thompson, L., & Sherrard, D. J. (1996). Dietary and behavioral risk factors for urolithiasis: potential implications for prevention. *American Journal of Kidney Diseases*, 28(2), 195-201. doi:10.1016/S0272-6386(96)90301-7
- Leonetti, F., Dussol, B., Berthezene, P., Thirion, X., & Berland, Y. (1998). Dietary and urinary risk factors for stones in idiopathic calcium stone formers compared with healthy subjects. *Nephrology Dialysis Transplantation*, 13(3), 617-622. Retrieved from <http://ndt.oxfordjournals.org/>
- Lewandowski, S., & Rodgers, A. L. (2004). Idiopathic calcium oxalate urolithiasis: Risk factors conservative treatment. *International Journal of Clinical Chemistry*, 345(1-2), 17-34. doi:10.106/j.cccn.2004.03.009
- Lopez, M., & Hoppe, B. (2010). History, epidemiology and regional diversities of urolithiasis. *Pediatric Nephrology*, 25(1), 49-59. doi:10.1007/s00467-008-09605
- Lopez-Azpiazu, I., Martinez-Gonzalez, M. A., Kearney, J., Gibney, M., & Martinez, J. A. (1999). Perceived barriers of, and benefits to, healthy eating reported by a Spanish national sample. *Public Health Nutrition*, 2(2), 209-215. doi:10.1017/S1368980099000269
- Lynn, M. R. (1986). Determination and quantification of content validity. *Nursing Research*, 35(6), 382-385. Retrieved from <http://ovidsp.tx.ovid.com/>
- Martini, L. A., & Wood, R. J. (2000). Should dietary calcium and protein be restricted in patients with nephrolithiasis? *Nutrition Reviews*, 58(4), 111-117. doi:10.1111/j.1753-4887.2000.tb07541.x

- Masarani, M., & Dinneen, M. (2007). Ureteric colic: New trends in diagnosis and treatment. *Postgraduate Medical Journal*, 83(981), 469-472.
doi:10.1136/pgmj.2006.055913
- Massey, L. K., & Wise, K. J. (1992). Impact of gender and age on urinary water and mineral excretion responses to acute caffeine doses. *Nutrition Research*, 12(4-5), 605-612. doi:10.1016/S0271-5317(05)80030-2
- McCauley, L. R., Dyer, A. J., Stern, K., Hicks, T., & Nguyen, M. M. (2012). Factors influencing fluids intake behavior among kidney stone formers. *The Journal of Urology*, 187(4), 1282-1286. doi:10.1016/j.juro.2011.11.111
- McGuire, A. M., & Anderson, D. J. (2012). Lifestyle risk factor modification in midlife women with type 2 diabetes: Theoretical modelling of perceived barriers. *Australian Journal of Advanced Nursing*, 30(1), 49-57. Retrieved from <http://eprints.qut.edu.au/54025/>
- Meschi, T., Maggiore, U., Fiaccadori, E., Scianchi, T., Bosi, S., Adorni, E. M., . . . Borghi, L. (2004). The effect of fruits and vegetables on urinary stone risk factors. *Kidney International*, 66(6), 2402-2410. doi:10.1111/j.1523-1755.2004.66029.x
- Meschi, T., Nouvenne, A., Ticinesi, A., Prati, B., Guerra, A., Allegri, F., . . . Borghi, L. (2012). Dietary habits in women with recurrent idiopathic calcium nephrolithiasis. *Journal of Translational Medicine*, 10, 63. doi:10.1186/1479-5876-10-63
- Meschi, T., Schianchi, T., Ridolo, E., Adorni, G., Allegri, F., Guerra, A., . . . Borghi, L. (1999). Body weight, diet and water Intake in preventing stone disease. *Urology Internationalis*, 72, 29-33. doi:10.1159/000076588
- Ministry of Domestic Health, Co-Operatives and Consumerism, Malaysia. (2015). *Harga barangan terkawal*. Retrieved from <http://www.kpdnkk.gov.my/>
- Ministry of Health Malaysia. (2007). *Health indicator*. Retrieved from <http://vlib.moh.gov.my/>

- Ministry of Health Malaysia. (2008). *Health indicator*. Retrieved from <http://vlib.moh.gov.my/>
- Ministry of Health Malaysia. (2009). *Health indicator*. Retrieved from <http://vlib.moh.gov.my/>
- Ministry of Health Malaysia. (2010). *Health indicator*. Retrieved from <http://vlib.moh.gov.my/>
- Ministry of Health Malaysia. (2012). *Health indicator*. Retrieved from <http://vlib.moh.gov.my/>
- Ministry of Human Resources, Malaysia. (2013). *Minimum retirement age*. Retrieved from <http://www.mohr.gov.my/>
- Moe, O. W. (2006). Kidney stones: Pathophysiology and medical management. *The Lancet*, 367(9507), 333-344. doi:10.1016/S0140-6736(06)68071-9
- Morowati Sharifabad, M. A., Pirouzeh, R., Hemayati, R., & Askarshi, M. (2015). Preventive behaviors in recurrent kidney stone and barriers to performing these behaviors. *Journal of Research and Health*, 5(2), 230-239. Retrieved from <http://jrh.gmu.ac.ir>
- Morton, A. R., Ilescu, E. A., & Wilson, J. W. L. (2002). Nephrology 1: Investigation and treatment of recurrent kidney stones. *Canadian Medical Association Journal*, 166(2), 213-218. Retrieved from <http://www.cmaj.ca/>
- Muhbes, F. J. (2012). Risk factors for renal stone formation: A field study. *Health Science Journal*, 6(4), 714. Retrieved from <http://www.hsj.gr>
- Murphy, S. P., Kaiser, L. L., Townsend, M. S., & Allen, L. H. (2001). Evaluation of validity of items for a food behavior checklist. *Journal of the American Dietetic Association*, 101(7), 751-761. doi:10.1016/S0002-8223(01)00189-4

- Nazmi, N., Zainal, D., & Hashim, M. (1997). Renal stone in Kelantan, Malaysia—Has the pattern changed? *Southeast Asian Journal of Tropical Medicine Public Health*, 28(3), 641-644. Retrieved from <http://www.ncbi.nlm.nih.gov/>
- Ngo, T. C., & Assimos, D. G. (2007). Uric acid nephrolithiasis: recent progress and future directions. *Reviews in Urology*, 9(1), 17-27. Retrieved from <http://www.ncbi.nlm.nih.gov/>
- Norimah, A. K., Safiah, M., Jamal, K., Siti, H., Zuhaida, H., Rohida, S., Fatimah, S., . . . Azmi, M. Y. (2008). Food consumption patterns: Findings from the Malaysian Adult Nutrition Survey (MANS). *Malaysian Journal of Nutrition*, 14(1), 25-39. Retrieved from <https://www.researchgate.net/>
- Norris, J. (2013). *Vegan for life: Oxalate*. Retrieved from <http://www.veganhealth.org/>
- Nouvenne, A., Meschi, T., Prati, B., Guerra, A., Allegri, F., Vezzoli, G., Soldati, L., . . . Borghi, L. (2010). Effects of a low-salt diet on idiopathic hypercalciuria in calcium-oxalate stone formers: A 3-month randomized controlled trial 1-3. *American Journal of Clinical Nutrition*, 91(3), 565-570. doi:10.3945/ajcn.2009.28614
- Pak, C. Y. (1994). Citrate and renal calculi: An update. *Mineral Electrolyte Metabolite*, 20(6), 371-377. Retrieved from <http://www.ncbi.nlm.nih.gov/>
- Pak, C. Y. (2004). Medical management of urinary stones disease. *Nephron Clinical Practice*, 98(2), c49-c53. doi:10.1159/000080252
- Pak, C. Y., Odvina, C.V., Pearle, M. S., Sakhaee, K., Peterson, R. D., Poindexter, J. R., & Brinkley, L. J. (2005). Effect on diet modification on urinary stone risk factors. *Kidney International*, 68(5), 2264-2273. doi:10.1111/j.1523-1755.2005.0068.x
- Pak, C. Y., Sakhaee, K., Moe, O., Preminger, G. M., Poindexter, J. R., Peterson, R. D., . . . Ekeruo, W. (2003). Biochemical profile of stone-forming patients with diabetes mellitus. *Adult Urology*, 61(3), 523-527. doi:10.1016/S0090-4295(02)02421-4

- Papadoukakis, S., Stolzenburg, J. U., & Trussa, M. C. (2006). Treatment strategies of ureteral stones. *European Association of urology & European Board of Urology Update Series*, 4, 184-190. doi:10.1016/j.eeus.2006.07.004
- Park, S., & Pearle, M. S. (2007). Pathophysiology and management of calcium stones. *Urologic Clinics of North America*, 34(3), 323-334. doi:10.1016/j.ucl.2007.04.009
- Parmar, M. S. (2004). Kidney stones. *British Medical Journal*, 328(7453), 1420-1424. doi:10.1136/bmj.328.7453.1420
- Patel, A. C., & Mehta, N. H. (2014). Epidemiological characteristics of renal stone patients age (21-60) and barriers in their dietary modification in Saurashtra Region. *International Research of Medical Science*, 2(1), 80-83. doi:10.5455/2320-6012.ijrms20140216
- Paterson, R., Fernandez, A., Razvi, H., & Sutton, R. (2010). Evaluation and medical management of the kidney stone patient: CUA guideline. *Canadian Urology Association Journal*, 4(6), 375-379. Retrieved from <http://www.ncbi.nlm.nih.gov/>
- Pawlak, R., & Colby, S. (2009). Benefits, barriers, self-efficacy and knowledge regarding healthy foods: Perception of African Americans living in the eastern North Carolina. *Nutrition Research and Practice*, 3(1), 56-63. doi:10.4162/nrp.2009.3.1.56
- Pearle, M. S., Goldfarb, D. S., Assimos, D. G., Curhan, G., Denu-Ciocca, C. C., Matiaga, B. R., . . . White, J. R. (2014). Medical management of kidney stones: American urological (AUA) guideline. *The Journal of Urology*, 192(2), 316-324. doi:10.1016/j.juro.2014.05.006
- Pires, C. G. S., & Mussi, F. C. (2012). Health beliefs regarding diet: A perspective of hypertensive blacks individuals. *Revista da Escola de Enfermagem da USP*, 46(3), 580-588. doi:10.1590/S0080-62342012000300008

- Pizzato, A. C., & Barros, E. J. G. (2003). Dietary calcium intake among patients with urinary calculi. *Nutrition Research*, 23(12), 1651-1660.
doi:10.1016/j.nutres.2003.09.001
- Polit, D. F., & Beck, C. T. (2006). The content validity index: Are you sure you know what's being reported? Critique and recommendations. *Research in Nursing & Health*, 29(5), 489-497. doi:10.1002/nur.20147
- Polit, D. F., & Hungler, B. P. (2004). *Nursing research: Principle and methods* (7th ed.). Philadelphia. Lippincot Williams & Wilkins.
- Prasetyo, T., Birowo, P., & Rasyid, N. (2013). The influence of increased fluid intake in the prevention of urinary stone formation: A systemic review. *Acta Medica Indonesiana - The Indonesian Journal of Internal Medicine*, 45(4), 253-258.
Retrieved from <http://www.inaactamedica.org/>
- Qaseem, A., Dallas, P., Forciea, M. A., Starkey, M., & Denberg, T. D. (2014). Dietary and pharmacologic management to prevent recurrent nephrolithiasis in a dults: A clinical practice guideline from the American College of Physicians. *Annal Internal Medicine*, 161(9), 659-667. doi:10.7326/M13-2908
- Ramello, A., Vitale, C., & Marangella, M. (2000). Epidemiology of nephrolithiasis. *Journal of Nephrology*, 3, 45-50. Retrieved from <http://www.ncbi.nlm.nih.gov/>
- Reddy, S. T., Wang, C. Y., Sakhaee, K., Brinkley, L., & Pak, C. Y. (2002). Effect of low-carbohydrate high-protein diets on acid-base balance, stone-forming propensity, and calcium metabolism. *American Journal of Kidney Diseases*, 40(2), 265-274. doi:10.1053/ajkd.2002.34504
- Remer, T. & Manz, F. (1995). Potential renal acid load of foods and its influence on urine pH. *Journal of the American Dietetic Association*, 95(7), 791-797.
doi:10.1016/S0002-8223(95)00219-7.
- Rippe, J. M. (2013). *Lifestyle medicine* (2nd ed.). Retrieved from <https://books.google.co.th>

- Romero, V., Akpinar, H., & Assimos, D. G. (2010). Kidney stones: A global picture of prevalence, incidence, and associated risk factors. *Reviews in Urology*, 12(23), e86-e96. Retrieved from <http://www.ncbi.nlm.nih.gov/pmc/>
- Rosdahl, C. B., & Kowalski, M. T. (2008). *Textbook of basic nursing*. Philadelphia: Lippincott Williams & Wilkins.
- Rosenstock, I. M. (1974). Historical origins of the Health Belief Model. *Health Education & Behavior*, 2(4), 328-335. doi:10.1177/10901981740020040
- Rosenstock, I. M., Strecher, V. J., & Becker, M. H. (1988). Social Learning Theory and the health belief model. *Health Education Quarterly*, 15, 175-183. Retrieved from <http://sjmse-library.sch.ng/>
- Roy, D. K. (2006). Role of diet in prevention of recurrent nephrolithiasis. *The ORION Medical Journal*, 20, 391-394. Retrieved from <http://www.oriongroup.net/medicaljournal/>
- Rule, A. D., Krambeck, A. E., & Lieske, J. C. (2011). Chronic kidney disease in kidney stone formers. *Clinical Journal of the American Society of Nephrology*, 6(8), 2069-2075. doi:10.2215/CJN.10651110
- Sabzmakan, L., Morowatisharifabad, M. A., Mohammadi, E., Mazloomi-Mahmoodabad, S. S., Rabiei, K., Naseri, M. H., ... Mirzaei, M. (2014). Behavioral determinants of cardiovascular diseases risk factors: A qualitative directed content analysis. *ARYA Atheroscler*, 10(2), 71-81. Retrieved from <http://www.ncbi.nlm.nih.gov/>
- Saigal, C. S., Joyce, G., & Timilsina, A. R. (2005). Direct and indirect costs of nephrolithiasis in an employed population: Opportunity for disease management? *The Urologic Diseases in America Project*, 68(4), 1808-1814. doi:10.1111/j.1523-1755.2005.00599

- Saint-Elie, D. T., Patel, P. V., Healy, K. A., Solomon, T., Pattaras, J. G., Qian, J., . . . Ogan, K. (2010). the impact of income and education on dietary habits in stone formers. *Urology*, *76*(2), 307-313. doi:10.1016/j.urology.2009.11.028
- Salahshoori, A., Sharifirad, G., Hassanzadeh, A., & Mostafavi, F. (2014). An assessment of the role of perceived benefits, barriers and self-efficacy in predicting dietary behavior in male and female high students in the City of Izeh, Iran. *Journal of Education and Health Promotion*, *3*, 8. doi:10.4103/2277-9531.127558
- Salmeh, F., Yaghoubi, T., Zakizadeh, M., Yaghoubian, M., & Shahmohammadi, S. (2012). Evaluation of health behaviours in patients with kidney stones in Sari/Iran. *International Journal of Urological Nursing*, *6*(1), 17-21. doi:10.1111/j.1749-771X.2012.01139.x
- Samplaski, M. K., Irwin, B. H., & Desai, M. (2009). Less invasive way to remove stones from the kidneys and ureters. *Cleveland Clinic Journal of Medicine*, *76*(10), 592-598. doi:10.2949/ccjm.76a.09014
- Sarica, K, Inal, Y., Erturhan, S., & Yagci, F. (2006). The effect of calcium channel blockers on stone regrowth occurrence after shock wave lithotripsy. *Urology Research*, *34*(3), 184-189. doi:10.1007/s00240-006-0040-x
- Saxena, A., & Sharma, R. K. (2010). Nutritional aspect of nephrolithiasis. *Indian Journal of Urology*, *26*(4), 523-530. doi:10.4103/0970-1591.74451
- Scales, C. D., Smith, A. C., Hanley, J. M., & Saigal, C. S. (2012). Prevalence of kidney stones in the United States. *European Urology*, *62*(1), 160-165. doi:10.1016/j.eururo.2012.03.052
- Semins, M. J., Shore, A. D., Makary, M. A., Magnuson, T., Johns, R., & Matlaga, B. R. (2010). The association of increasing body mass index and kidney stone disease. *The Journal of Urology*, *183*(2), 571-575. doi:10.1016/j.juro.2009.09.085

- Sharifirad, G., Entezari, M. H., Kamran, A., & Azadbakht, L. (2009). The effectiveness of nutritional education on the knowledge of diabetic patients using the health belief model. *Journal of Research in Medical Science*, *14*(1), 1-6. Retrieved from <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3129063/>
- Siener, R., Ebert, D., Nicolay, C., & Hesse, A. (2003). Dietary risk factors for hyperoxaluria in calcium oxalate stone formers. *Kidney International*, *63*(3), 1037-1043. doi:10.1046/j.1523-1755.2003.00807.x
- Siener, R., & Hesse, A. (2003). Fluid intake and epidemiology of urolithiasis. *European Journal of Clinical Nutrition*, *57*(Suppl. 2), S47-S51. doi:10.1038/sj.ejcn.1601901
- Skolarikos, A., Straub, M., Knoll, T., Sarica, K., Seitz, C., Petrik, A., & Turk, C. (2015). Metabolic evaluation and recurrence prevention for urinary stone patients: EAU guidelines. *European Urology*, *67*(4)750-763. doi:10.1016/j.eururo.2014.10.029
- Sorensen, M. D., Hsi, R. S., Chi, T., Shara, N., Wactawski-Wende, J., Kahn, A. J., . . . Stoller, M. L. (2014). Dietary intake of fiber, fruits and vegetables decrease the risk of incident kidney stones in women: A Women's Health Initiative (WHI) report. *The Journal of Urology*, *192*(6), 1694-1699. doi:10.1016/j.juro.2014.05.086
- Sorensen, M. D., Kahn, A. J., Reiner, A. P., Tseng, T. Y., Shikany, J. M., Wallace, R. B., . . . Stoller, M. L. (2012). Impact of nutritional factors on incident kidney stone formation: A Report from the WHI OS. *Journal of Urology*, *187*(5), 1645-1649. doi:10.1016/j.juro.2011.12.077
- Strazzullo, P., Barba, G., Vuotto, P., Farinaro, E., Siani, A., Nunziata, V., . . . Cappuccio, F. P. (2001). Past history of nephrolithiasis and incidence of hypertension in men: A reappraisal based on the results of the olivetti prospective heart study. *Nephrology Dialysis Transplantation*, *16*(11), 2232-2235. Retrieved from <http://www.ncbi.nlm.nih.gov/>

- Strecher, V. J. & Rosenstock, I. M. (1997). The health belief model. In K. Glanz, M. F., Lewis, & B. K. Rimer (Eds.), *Health behavior and health education: Theory, research, and practice* (2nd ed., pp. 41-59). San Francisco, CA: Jossey-Bass.
- Tanthanuch, M., Apiwatgaroon, A., & Pripatnanont, C. (2005). Urinary tract calculi in Southern Thailand. *Journal of Medical Association Thailand*, 88(1), 80-85. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/15960223>
- Taylor, E. N., Stampfer, M. J., & Curhan, G. C. (2004). Dietary factors and the risk of incident kidney stones in men: New insights after 14 years of follow-up. *Journal of the American Society of Nephrology*, 15(12), 3225-3232. doi:10.1097/01.ASN.0000146012.44570.20
- Timio, F., Kerry, S. M., Anson, K. M., Eastwood, J. B., & Cappuccio, F. P. (2003). Calcium urolithiasis, blood pressure and salt intake. *Blood Pressure*, 12(2), 122-127. doi:10.1080/08037050310001084
- Timmerman, G. M. (2007). Addressing barriers to health promotion in underserved women. *Family & Community Health*, 30(Suppl. 1), S34-S42. Retrieved from <http://ovidsp.tx.ovid.com/>
- Tosukhowong, P., Boonla, C., Ratchanon, S., Tanthanuch, M., Poonpirome, K., Supataravanich, P., Dissayabutra, T., & Tungsanga, K. (2007). Crystalline composition and etiologic factors of kidney stone in Thailand: Update 2007. *Asian Biomedicine*, 1(1), 89-97. Retrieved from <http://abm.digitaljournals.org/>
- Traver, M. A., Passman, C. M., LeRoy, T., Passmore, L., & Assimios, D. G. (2009). Is the internet a reliable source for dietary recommendation for stone former? *Journal of Urology*, 23(4), 715-717. doi:10.1089/end.2008.0490
- Trinchieri, A. (2008). Epidemiology of urolithiasis: An update. *Clinical Cases in Mineral and Bone Metabolism*, 5(2), 101-106. Retrieved from <http://www.ncbi.nlm.nih.gov/pmc/>

- Trinchieri, A., Ostini, F., Nespoli, R., Rovera, F., Montanari, E., & Zanetti, G. (1999). A prospective study of recurrence rate and risk factors for recurrence after a first renal stone. *The Journal of Urology*, *162*(1), 27-30. doi:10.1097/00005392-199907000-00007
- Türk, C., Knoll, T., Petrik, A., Sarica, K., Skolarikos, A., Straub, M., . . . Seitz, C. (2014). *Guidelines on urolithiasis*. European Association of Urology. Retrieved from <http://www.uroweb.org/>
- Walker, S. N., Pullen, C. H., Hertzog, M., Boeckner, L., & Hageman, P. (2006). Determinants of older rural women's activity and eating. *Western Journal of Nursing Research*, *28*(4), 449-468. doi:10.1177/0193945906286613
- Wan Hafiz, W. Z. S. (2005). *Consumer eating habits and nutritional perception of hawker foods* (Master's thesis). Retrieved from <http://ir.uitm.edu.my/>
- Wilkinson, H. (2001). Clinical investigation and management of patients with renal stones. *Annals of Clinical Biochemistry*, *38*, 180-187. Retrieved from <http://oclc.lib.cmu.ac.th>
- Wolf, J. S., Jr., Howes, D. S., & Craig, S. (2014). *Nephrolithiasis*. Retrieved from <http://emedicine.medscape.com/>
- Wood, R. G., Goesling, B. & Avellar., S. (2007). *The effect of marriage on Health: A synthesis of recent research evidence*. Retrieved from <http://aspe.hhs.gov/sites/default/files/pdf/75106/report.pdf>
- Xu, H., Zisman, A. L., Coe, F. L., & Worcester, E. M. (2013). Kidney stones: An update pharmacological management and future directions. *Expert Opinion Pharmacotherapy*, *14*(4), 435-447. doi:10.1217/14656566.2103.775250

Yanagawa, M., Kawamura, J., Onishi, T., Soga, N., Kameda, K., Sriboonlue, P., . . .
Borwornpadungkitti, S. (1997). Incidence of urolithiasis in Northeast Thailand.
International Journal of Urology, 4(6), 537-540. Retrieved from
<http://oclc.lib.cmu.ac.th>

Yasui, T., Iguchi, M., Suzuki, S., & Kohri, K. (2008). Prevalence and epidemiological
characteristics of urolithiasis in Japan: National trends between 1965 and 2005.
Urology, 71(2), 209-213. doi:10.1016/j.urology.2007.09.034

Zargooshi, J. (2001). Open stone surgery in children: is it justified in the era of
minimally invasive therapies? *British Journal of Urology International*, 88(9),
928-931. doi:10.1046/j.1464-4096.2001.01544.x



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved