

REFERENCES

- Aldrich & John. (2005). Fisher and Regression. *Statistical Science*, 20 (4); 401–417.
- Alfonso, V., Toulson, A., Bermbach, N., Erskine, Y. & Montaner, J. (2009). Psychosocial issues influencing treatment adherence in patients on multidrug rescue therapy: perspectives from patients and their health care providers. *AIDS Patient Care STDS*, 23, 119–126.
- Ajzen, I. (1988). *Attitudes, personality, and behavior*. Milton-Keynes, England: Open University Press & Chicago, IL: Dorsey Press.
- Ajzen, I. (1991). The theory of Planned Behavior. *Organizational Behavior and Human Decision Processes*, 50, 179-211.
- Alemnesh, H.M., Mitike, M.S., Karen, M.M. & Anne N.A. (2011). Applying the theory of planned behaviour to explain HIV testing in antenatal settings in Addis Ababa - a cohort study. *BMC Health Services Research*, 11, 196.
- Amberbir, A., Woldemichael, K., Getachew, S.,Girma, B. & Deribe, K. (2008). Predictors of adherence to antiretroviral therapy among HIV-infected person: a prospective study in Southwest Ethiopia. *BMC Public Health*, 8, 265.
- American Academy of Pediatrics, (1999). Committee on Pediatric AIDS. Disclosure of illness status to children and adolescents with HIV infection. *Pediatrics*, 103, 164-166.
- Amico, K.R., Fisher, W.A., Cornman, D.H., Shuper, P.A., Redding, C.G. & Konkle-Parker, D.J., et al. (2006). Visual analog scale of ART adherence: association with 3-day self-report and adherence barriers. *J Acquir Immune Defect Syndr*, 42(4); 455-9.
- Ammassari, A., Trotta, M.P., Murri, R., Castelli, F., Narciso, P. & Noto, P., et al. (2002). Correlates and predictors of adherence to highly active antiretroviral therapy: overview of published literature. *J Acquir Immune Defic Syndr*, 31(Suppl 3); S123–127.

- Ammassari, A., Antinori, A., Aloisi, M.S., Trotta, M.P., Murri, R. & Bartoli, L., et al. (2004). Depressive symptoms, neurocognitive impairment, and adherence to highly active antiretroviral therapy among HIV-infected persons. *Psychosomatics*, 45, 395-405.
- Ananworanich, J., Kosalaraksa, P., Siangphoe, U., Engchanil, C., Pancharoen, C. & Lumbiganon, P., et al. (2008). A feasibility study of immediate versus deferred antiretroviral therapy in children with HIV infection. *AIDS Research and Therapy*, 5, 24.
- Andrea, S., Carol, G., Jane, M., Luborsky, M. & Pearson, C. (2006). How qualitative methods contribute to understanding combination antiretroviral therapy adherence. *J Acquir Immune Defic Syndr*, 43(1); S54-S68.
- Anthony, G., Greenwald, Richard Gonzalez, Harris, R.J. & Guthrie, D. (1996). Effect sizes and p values: What should be reported and what should be replicated ?. *Psychophysiology*, 33, 175-83.
- Arsalan, M.G. (2010). AIDS/HIV Knowledge, Attitude and Beliefs of Adolescents of Pakistan. *European Journal of Social Sciences*, 16(2); 275-285.
- Azmeraw, D. & Wasie, B. (2012). Factors associated with adherence to highly active antiretroviral therapy among children in two referral hospitals, northwest Ethiopia. *Ethiop Med J*, 50(2); 115-24.
- Baghianmoghaddam, M.H., Forghani, H., Zolghadr, R., Rahaii, Z. & Khani P. (2010). Health Belief Model and HIV/AIDS among high school female students in Yazd, Iran. *J Res Med Sci*, 15(3); 189-190.
- Baker, J.V., Peng, G., Rapkin, J., Abrams, D.I., Silverberg, M.J. & MacArthur, R.D., et al. (2008). CD4+ count and risk of non-AIDS diseases following initial treatment for HIV infection. *AIDS*, 22, 841-8.
- Balfour, L., Lowal, J., Silverman, A., Tasca, G.A., Angel, J.B., & Macpherson, P.A., et al. (2006). A randomized controlled psycho-education intervention trial: improving psychological readiness for successful HIV medication adherence and reducing depression before initiating HARRT. *AIDS Care*, 18, 830–838.
- Bandura, A. (1977). “Self-Efficacy: Toward a Unifying Theory of Behavioral Change” *.Psychology Review*, 84, 191-215.

- Bandura, A. (1986). *Social foundations of thought and action: A social cognitive theory*, Prentice-Hall, Englewood Cliffs, NJ.
- Bandura, A. Human Agency in Social Cognitive Theory. (1989). *American Psychologist*, 44, 1175–1184.
- Bandura, A. (1988). Organizational Application of Social Cognitive Theory. *Australian Journal of Management*, 13(2); 275–302.
- Bandura, A. (1994). Self-efficacy. In V. S. Ramachaudran (ed.), *Encyclopedia of human behavior*. New York: Academic Press, (4); 71-81.
- Bangsberg, D. (2006). Less than 95% adherence to nonnucleoside reverse-transcriptase inhibitor therapy can lead to viralsuppression. *Clin Infec Dis*, 43(7); 939-41.
- Bangsberg, D.R., Perry, S., Charlebois, E.D., Clark, R.A., Roberston, M. & Zolopa, A.R., et al. (2001). Nonadherence to highly active antiretroviral therapy predicts progression to AIDS. *AIDS*, 15, 1181-3.
- Barclay, Terry, R., Hinkin, Mason, K.I., Reinhard, M.J. & Marion, S.D., et al. (2007). Age-associated predictors of medication adherence in HIV-positive adults: Health beliefs, self-efficacy, and neurocognitive status. *Health Psychology*, 26(1); 40-49.
- Belzer, M.E., Fuchs, D.N., Luftman, G.S. & Tucker, D.J. (1999). Antiretroviral adherence issues among HIV-positive adolescents and young adults. *J Adolesc Health*, 25, 316-9.
- Berg, K.M., Demas, P.A., Howard, A.A., Schoenbaum, E.E., Gourevitch, M.N. & Arnsten, J.H. (2004). Gender differences in factors associated with adherence to antiretroviral therapy. *J Gen Intern Med*, 19(11); 1111-1117.
- Bernstein, K., D'Angelo, L.J. & Lyon, M.E. (2012). An exploratory study of HIV+ adolescents' spirituality: Will you pray with me?. *J Relig Health*, 19[Epub ahead of print].

- Bhakeecheep, S. (2011). *Pediatric HIV treatment and care in universal health coverage*. Slide presentation in international comprehensive HIV/AIDS care management, 1-5 Aug 2011, Thailand.
- Biering, P. (2007). Adapting the concept of explanatory models of illness to the study of youth violence. *Journal of Interpersonal Violence*, 22(7); 791-811.
- Birbeck, G., Chomba, E., Kvalsund, M., Bradbury, R., Mang'ombe, C. & Malama, K., et al. (2009). Antiretroviral adherence in rural Zambia: the first year of treatment availability. *Am J Trop Med Hyg*, 80(4); 669-674.
- Blumer, H. (1969). *Symbolic interactionism: perspective and method*. Englewood Cliffs, NJ: Prentice Hall.
- Bouhnik, A.D., Preau, M., Vincent, E., Carrieri, M.P., Gallais, H. & Lepeu, G., et al. (2005). Depression and clinical progression in HIV-infected drug users treated with highly active antiretroviral therapy. *Antivir Ther*, 10, 53-61.
- Bruce, K. E., & Walker, L. J. (2001). College students' attitudes about AIDS 1986 to 2000. *AIDS Education and Prevention*, 13(5); 428-437.
- Buchanan, A.L., Montepiedra, G., Sirois, P.A., Kammerer, B., Garvie, P.A. & Storm, D.S., et al. (2012). Barriers to medication adherence in HIV-infected children and youth based on self and caregiver report. *Pediatrics*, 129(5); 1244-51.
- Busza, J., Dauya, E., Bandason, T., Mujuru, H. & Ferrand, R.A. (2014). "I don't want financial support but verbal support." How do caregivers manage children's access to and retention in HIV care in urban Zimbabwe?. *J Int AIDS Soc*. Available from URL : <http://www.jiasociety.org/index.php/jias/article/view/18839> <http://dx.doi.org/10.7448/IAS.17.1.18839>. Accessed: 18 Jan 2015.
- Camelia, P., Francois, R., Perrine, R., Reynes, J., Dellamonica, P. & Spire, B., et al. (2009). Factors associated with non-adherence to long-term highly active antiretroviral therapy: a 10 year follow-up analysis with correction for the bias induced by missing data. *Journal of Antimicrobial Chemotherapy*, 64, 599-606.
- Castro, M., González, I. & Pérez, J. (2015). Factors Related to Antiretroviral Therapy Adherence in Children and Adolescents with HIV/AIDS in Cuba. *MEDICC Rev*, 17(1); 35-40.

- Catz, S.L., Kelly, J.A., Bogart, L.M., Benotsch, E.G. & McAuliffe, T.L. (2000). Patterns, correlates, and barriers to medication adherence among persons prescribed new treatments for HIV disease. *Health Psychology*, 19(2); 124–133.
- Center of Disease Control and Prevention (CDC). (2011). *Basic information about HIV and AIDS*. Available from URL :<http://www.cdc.gov/hiv/topics/basic/#origin>. Accessed: 10 Nov 2011.
- C. Dollfus, J. Le Chenadec, A. Faye, Blanche, S., Briand, N. & Rouzioux, C., et al. (2010). Long-term outcome in adolescents perinatally infected with HIV-1 and followed up since birth in the France Perinatal Cohort (EPF/ANRS CO10). *CID*, 51(15); 214-224.
- Chasombat, S., Lertpiriyasuwat, C., Thanprasertsuk, S., Suebsaeng, L. & Lo, Y.R. (2006). The National Access to Antiretroviral Program for PHA in Thailand. *Southeast Asian J Trop Med Public Health*, 37(4); 704-15.
- Chesney, M.A., Ickovics, J.R., Chambers, D.B., Gifford, A.L., Neidig, J. & Zwickl, B., et al. (2000). Self-reported adherence to antiretroviral medications among participants in HIV clinical trials: the AACTG adherence instruments. Patient Care Committee & Adherence Working Group of the Outcomes Committee of the Adult AIDS Clinical Trials Group (AACTG). *AIDS Care*, 12(3); 255-266.
- Chesney, M.A. (2003). Adherence to HAART regimens. *AIDS Patient Care STDS*, 17, 169-177.
- Cohen, J. (1988). *Statistical power analysis for the behavioral sciences* (2nd ed.). Hillsdale, NJ: Lawrence Earlbaum Associates.
- Crum, N.F., Riffenburgh, R.H., Wegner, S., Agan, B.K., Tasker, S.A. & Spooner, K.M., et al. (2006). Comparison of causes of death and mortality rates among HIV-infected persons: analysis of the pre-, early, and late HAART (highly active antiretroviral therapy) eras. *J Acquir Immune Defic Syndr*, 41, 194-200.
- Cynthia, R., Pearson, Mark M., Matediana, E., Martin, D.P. & Gloyd, S. (2006). Modified directly observed therapy to facilitate highly active antiretroviral therapy adherence in Beira, Mozambique. *J Acquir Immune Defic Syndr*, 1(43); S134-S141.
- Dahab, M., Charalambous, S., Hamilton, R., Fielding, K., Kielmann, K. & Churchyard, G.J., et al. (2008). “That is why I stopped the ART”: patients’ & providers’

- perspectives on barriers to and enablers of HIV treatment adherence in a South African workplace programme. *BMC Public Health*, 8, 63.
- DeLaMora, P., Aledort, N. & Stavola, J. (2006). Caring for adolescents with HIV. *Curr HIV/AIDS Rep*, 3(2); 74-8.
- Denison, J.A., Banda, H., Dennis, A.C., Packer, C., Nyambe, N. & Stalter, R.M., et al. (2015). "The sky is the limit": adhering to antiretroviral therapy and HIV self-management from the perspectives of adolescents living with HIV and their adult caregivers. *J Int AIDS Soc*, 18(1):19358. doi: 10.7448/IAS.18.1.19358. eCollection 2015.
- DiClemente, C., & Prochaska, J. (1985). *Processes and stages of self-change: coping and competence in smoking behavior change*. In: Shiffman S, Willis T, editors. *Coping and substance abuse*. Orlando, FL: Academic Press, 319–343.
- Dube' M.P., Sprecher, D., Henry, W.K., Aberg, J.A., Torriani, F.J. & Hodis, H.N., et al. (2000). Preliminary guidelines for the evaluation and management of dyslipidemia in adults infected with human immunodeficiency virus and receiving antiretroviral therapy: Recommendations of the Adult AIDS Clinical Trials Group Cardiovascular Disease Focus Group. *Clin Infect Dis*, 31, 1216-1224.
- Elizabeth, H., Bradley & Leslie, A. (2006). Qualitative data analysis for health services research: Developing taxonomy, themes and theory. *Health Research and Educational Trust*, 1758-72.
- Emery, S., Neuhaus, J.A., Phillips, A.N., Babiker, A., Cohen, C.J. & Gatell, J.M., et al. (2008). Major clinical outcomes in antiretroviral therapy (ART)-naive participants and in those not receiving ART at baseline in the SMART study. *J Infect Dis*, 197, 1133-44.
- Enriquez, M., Cheng, A.L., McKinsey, D.S. & Stanford, J. (2009). Development and efficacy of an intervention to enhance readiness for adherence among adults who had previously failed HIV treatment. *AIDS Patient Care STDS*, 23, 177–184.
- Enriquez, M., Gore, P.A., O'Connor, M.C. & McKinsey, D.S. (2004). Assessment of readiness for adherence by HIV-positive males who had previously failed treatment. *J Assoc Nurses AIDS Care*, 15, 42–49.

- Filho, L.F.B., Nogueira, S.A., Machado, E.S., Abreu, T.F., de Oliveira, R.H. & Evangelista, L., et al. (2008). Factors associated with lack of antiretroviral adherence among adolescents in a reference centre in Rio de Janeiro, Brazil. *Int J STD AIDS*, 19(10); 685-688.
- Fisher, E.B., Strunk, R.C., Highstein, G.R., Kelley-Sykes, R., Tarr, K.L. & Trinkaus, K., et al. (2009). A randomized controlled evaluation of the effect of community health workers on hospitalization for asthma: the asthma coach. *Arch Pediatr Adolesc Med*, 163, 225-232.
- Fleury, J. (1994). The index of readiness: development and psychometric analysis. *J Nurs Manag*, 31, 470-477.
- Fogarty, L., Roter, D., Larson, S., Burke, J., Gillespie, J. & Levy, R. (2002). Patient adherence to HIV medication regimens: a review of published and abstract reports [Review]. *Patient Education & Counseling*, 46(2); 93-108.
- Fowler, M.E. (1998). Recognizing the phenomenon of readiness: concept analysis and case study. *J Assoc Nurses AIDS Care*, 9, 72-76.
- Gardner, E.M., Sharma, S., Peng, G., Hullsiek, K.H., Burman, W.J. & Macarthur, R.D., et al. (2008). Differential adherence to combination antiretroviral therapy is associated with virologic failure with resistance. *AIDS*, 22, 75-82.
- Gebrekristos, H.T., Mlisana, K.P., Krim, Q.A. (2005). Patients' readiness to start highly active antiretroviral treatment for HIV. *Br Med J*, 331, 772-775.
- Gifford, A.L., Bormann, J.E., Shively, M.J., Wright, B.C., Richman, D.D. & Bozzette, S.A. (2000). Predictors of self-reported adherence and plasma HIV concentrations in patients on multidrug antiretroviral regimens. *Journal of Acquired Immune Deficiency Syndromes*, 23(5); 386-395.
- Gossell, W.M., Williams, J.J., Williams, E.W. & Levy, P. (2014). A Case for a Holistic Approach to the Improvement of Compliance among Hypertensive Patients: A Hospital Review. *West Indian Med J*, 63(3); 275-277.
- Green, S.B. (1991). How many subjects does it take to do a regression analysis? *Multivariate Behavioral Research*, 26, 499-510.
- Gross, R., Bandason, T., Langhaug, L., Mujuru, H., Lowenthal, E. & Ferrand, R. (2015). Factors associated with self-reported adherence among adolescents on antiretroviral therapy in Zimbabwe. *AIDS Care*, 22;1-5.

- Hammami, N., Nöstlinger, C., Hoérée, T., Lefèvre, P., Jonckheer, T. & Kolsteren, P. (2004). Integrating adherence to highly active antiretroviral therapy into children's daily lives: a qualitative study. *Pediatrics*, 114(5):e591-7.
- Hammer, S.M., Saag, M.S., Schechter, M., Schooley, R.T., Thompson, M.A. & Walmsley, S., et al. (2008). Antiretroviral treatment of adult HIV infection: 2008 recommendations of the International AIDS Society-USA Panel. *JAMA*, 300, 555-70.
- Harris, R.J. (1985). *A primer of multivariate statistics* (2nd ed.). New York: Academic Press.
- Hartzell, J.D., Spooner, K., Howard, R., Wegner, S. & Wortmann, G. (2007). Race and mental health diagnosis are risk factors for highly active antiretroviral therapy failure in a military cohort despite equal access to care. *J Acquir Immune Defic Syndr*, 44, 411-416.
- Helestine, K., Howells, K. & Day, A. (2010). Brief anger interventions with offenders may be ineffective: a replication and extension. *Behav Res Therapy*, 48, 246–250.
- HIV guideline 2011. (2011). *Disclosure of HIV to Perinatally Infected Children and Adolescents*. Available from: URL: <http://www.hivguidelines.org/wp-content/uploads/disclosure-posted-08-02-2010.pdf>. Accessed: 18 June 2011.
- Hodgson, I. (2000). *Culture, meaning and perceptions: explanatory models and the delivery of HIV care*. Abstract MoPeD2772, XIIIth International AIDS Conference, Durban, South Africa, July14th-19th. Available from:URL: <http://www.brad.ac.uk/staff/ijhodgson/summaries/Publications/durban2000.htm>. Accessed: 10 Nov 2011.
- Holzemer, W.L. & Uys, L. (2004). Managing AIDS Stigma. *SAHARAJ*, 1(3); 165-174.
- Horne, R., Buick, D., Fisher, M., Leake, H., Cooper, V. & Weinman, J. (2004). Doubts about necessity and concerns about adverse effects: Identifying the types of beliefs that are associated with non-adherence to HAART. *International Journal of STD and AIDS*, 15, 38-44.
- Hudelson, C. & Cluver, L. (2015). Factors associated with adherence to antiretroviral therapy among adolescents living with HIV/AIDS in low- and middle-income countries: a systematic review. *AIDS Care*, 23;1-12.

- James, P., Anderson, J., Prior, J., White, J.P., Henry, J.A. & Cochrane, G.M. (1985). Patterns of drug taking in patients with chronic airflow obstruction. *Postgrad Med J*, 14, 7–10.
- Jeffrey, L.J., Kent, D., Kevin, D. & William, S. (2005). Introduction to structural equation modeling (path analysis). *SGIM Precourse PA 08*, 1-17.
- Jessica, E.H., Cook, A., Walker, A.S., Ngambi, M., Ferrier, A. & Mulenga, V., et al. (2011). Excellent adherence to antiretrovirals in HIV+ Zambian children is compromised by disrupted routine, HIV nondisclosure and paradoxical income effects. *PLoS ONE*, 6(4); 1-8.
- Johnson, M.O., Catz, S.L., Remien, R.H., Rotheram-Borus, M.J., Morin, S.F. & Charlebois, E., et al. (2003). Theory guided, empirically supported avenues for intervention on HIV medication non-adherence: Findings from the Healthy Living Project. *AIDS Patient Care STDS*, 17(12); 645–656.
- Joint United Nations Program on HIV/AIDS (UNAIDS). (2010). *Global report: UNAIDS report on the global AIDS epidemic* [internet]. 2010 [cited on 2011 June 17]. Available from URL : http://www.unaids.org/documents/20101123_GlobalReport_em.pdf
- Judith, A.E., EunSeok, C., Kevin, H., Caruthers, D. & Sereika, S.M. (2010). The HIV Medication Taking Self-Efficacy Scale: Psychometric Evaluation. *J Adv Nurs*, 66(11); 2560-2572.
- Judy, E. & Mill R.N. (2000). *Describing an Explanatory Model of HIV Illness among Aboriginal Women. Holistic Nursing Practice*, 15(1); 42. Available from URL : http://www.nursingcenter.com/library/JournalArticle.asp?Article_ID=100290. Accessed: 10 Nov 2011.
- Kalichman, S.C., Ramachandran, B. & Catz, S. (1999). Adherence to the combination antiretroviral therapies in HIV patients of low health literacy. *J Gen Intern Med*, 14, 267-273.
- Kalichman, S.C., Rompa, D., DiFonzo, K., Simpson, D., Austin, J. & Luke, W., et al. (2001). HIV treatment adherence in women living with HIV/AIDS: research based on the Information-Motivation-Behavioral Skills model of health behavior. *J Assoc Nurses AIDS Care*, 12(4); 58–67.

- Kalyango, J.N., Owino, E. & Nambuya, A.P. (2008). Non-adherence to diabetes treatment at Mulago Hospital in Uganda: Prevalence and associated factors. *Afr Health Sci*, 8, 67–73.
- Karl Peltzer, Natalie, F.P., Shandir R & Anderson, J. (2010). Antiretroviral treatment adherence among HIV patients in KwaZulu-Natal, South Africa. *BMC Public Health*, 10, 111.
- Kathleen Malee, Paige, L.W., Grace, M., Nichols, S., Sirois, P.A. & Storm, D., et al. (2009). The role of cognitive functioning in medication adherence of children and adolescents with HIV infection. *Journal of Pediatric Psychology*, 34(2); 164-175.
- Katz, I.T., Ryu, A.E., Onuegbu, A.G., Psaros, C., Weiser, S.D., Bangsberg, D.R. & et al. (2013). Impact of HIV-related stigma on treatment adherence: systematic review and meta-synthesis. *J Int AIDS Soc*, 16(3 Suppl 2):18640. doi: 10.7448/IAS.16.3.18640.
- Kaufmann, G.R., Perrin, L., Pantaleo, G., Opravil, M., Furrer, H. & Telenti, A., et al. (2003). CD4 T-lymphocyte recovery in individual with advanced HIV-1 infection receiving potent antiretroviral therapy for 4 years: the Swiss HIV cohort study. *Arch Intern Med*, 163, 2187-95.
- Kempf, M., Pisu, M., Dumcheva, A., Westfall, A.O., Kilby, J.M. & Saag, M.S. (2009). Gender differences in discontinuation of antiretroviral treatment regimens. *J Acquir Immune Defic Syndr*, 52(3); 336-41.
- Kenu, E., Obo-Akwa, A., Nuamah, G.B., Brefo, A., Sam, M. & Lartey, M. (2014). Knowledge and disclosure of HIV status among adolescents and young adults attending an adolescent HIV clinic in Accra, Ghana. *BMC Res Notes*, 26, 7(1); 844.
- Kip, E., Ehlers, V. & Wal, D. (2009). Patients' adherence to antiretroviral therapy in Botswana. *J Nurs Scholarsh*, 41(2); 149-157.
- Kleinman, A. (1976). Culture, illness and care: Clinical lessons from anthropologic and cross-cultural research. *Annals of Internal Medicine*, 88, 251-258.
- Kleinman, A. (1978). Culture, illness and cure: Clinical lesions from anthropologic and cross-cultural research. *Annals Int Med*, 88, 251-258.
- Knafl, G., Bova, C., Fennie, K., O'Malley, J.P., Dieckhaus, K.D. & Williams, A.B.

- (2008). An analysis of electronically monitored adherence to antiretroviral medications. *AIDS Behav*, 14(4); 755-68.
- Koopman, C, Rotherman-Borus, M.J., Henderson, R., Bradley, J.S. & Hunter, J. (1990). Assessment of knowledge of AIDS and beliefs about AIDS prevention among adolescents. *AIDS Educ Prev*, 2(1); 58-69.
- Kuhn, L., Semrau, K., Ramachandran, S., Sinkala, M., Scott, N. & Kasonde, P., et al. (2009). Mortality and virologic outcomes after access to antiretroviral therapy among a cohort of HIV-infected women who received single-dose nevirapine in Lusaka, Zambia. *J Acquir Immune Defic Syndr*, 52(1); 132-136.
- Lallemant, M., Jourdain, G. (2010). Preventing mother-to-child transmission of HIV-protecting this generation and the next. *N Eng J Med*, 363, 1570-2.
- Laurence, A., Gunar, G., Loretxu, Pinoges, L., Anguzu, P. & Taburet, A.M., et al. (2011). Immunological response to combined antiretroviral therapy and drug resistance pattern in children: 1- and 2-year outcomes in rural Uganda. *BMC Pediatrics*, 11, 67.
- Lee, A. Becker. (2000). *Effect size (ES)*. Available from URL : <http://web.uccs.edu/lbecker/Psy590/es.htm>. Accessed: 10 Feb 2012.
- Lima, V., Harrigan, R., Murray, M., Moore, D.M., Wood, E. & Hogg, R.S., et al. (2008). Differential impact of adherence on long-term treatment response among naive HIV-infected individuals. *AIDS*, 22, 2371-2380.
- Li, X., Huang, L., Wang, H., Fennie, K.P., He, G. & Williams, A.B. (2011). Stigma Mediates the Relationship Between Self-Efficacy, Medication Adherence, and Quality of Life Among People Living with HIV/AIDS in China. *AIDS Patient Care STDS*, 25(11); 665-71.
- Lohse, N., Hansen, A., Gerstoft, J. & Obel, N. (2007). Improved survival in HIV-infected persons: consequences and perspectives. *J Antimicrob Chemother*, 60(3); 461-463.
- Lowenthal, E.D., Marukutira, T.C., Chapman, J., Mokete, K., Riva, K. & Tshume, O., et al. (2014). Psychosocial assessments for HIV+ African adolescents: establishing construct validity and exploring under-appreciated correlates of adherence. *PLoS One*, 9(10):e109302. doi: 10.1371/journal.pone.0109302. eCollection 2014.

- Lynch, E., & Medin, D. (2006). Explanatory models of illness: A study within-culture variation. *Cognitive Psychology*, 53(4); 285.
- Mahat, G. & Scoloveno, M.A. (2006). HIV/AIDS knowledge, attitudes and beliefs among Nepalese adolescents. *J Adv Nurs*, 53(5); 583-90.
- Maithe Enriquez & David S McKinsey. (2011). Strategies to improve HIV treatment adherence in developed countries: clinical management at the individual level. *HIV/AIDS* (Auckl), 3, 45–51.
- Malow, R.W., Baker, S.M., Klimas, N., Antoni, M.H., Schneiderman, N. & Penedo, F.J., et al. (1998). Adherence to complex combination antiretroviral therapies by HIV-positive drug abusers. *Psychiatric services*, 49(8); 1021-1024.
- Maneesriwongul, W.L., Tulathong, S., Fennie, K.P. & Williams, A.B. (2006). Adherence to antiretroviral medication among HIV-positive patients in Thailand. *J Acquir Immune Defic Syndr*, 43(1); S119-S122.
- Manosuthi, W., Chottanapand, S., Thongyen, S., Chaovavanich, A. & Sungkanuparph, S. (2006). Survival rate and risk factors of mortality among HIV/tuberculosis-coinfected patients with and without antiretroviral therapy. *J Acquir Immune Defic Syndr*, 43, 42-6.
- Marhefka, S.L., Koenig, L.J., Allison, S., Bachanas, P., Bulterys, M. & Bettica, L., et al. (2008). Family experiences with pediatric antiretroviral therapy: responsibilities, barriers, and strategies for remembering medications. *AIDS Patient Care STDS*, 22(8); 637-47.
- Maria, L.S. Cruz, Claudete, A.A. Cardoso, Mariana, Q. & Darmont. (2014). Taking care of the caretakers to enhance antiretroviral adherence in HIV-infected children and adolescents. *Jornal de Pediatria*, 90 (6); 563-71.
- Martinez, J., Bell, D., Camacho, R., Henry-Reid, L.M., Bell, M. & Watson, C., et al. (2000). Adherence to antiviral drug regimens in HIV-infected adolescent patients engaged in care in a comprehensive adolescent and young adult clinic. *J Natl Med Assoc*, 92, 55-61.
- Massimo, M.S.L., Riccardo, C., Paola, G., Guidotti, G., Mancinelli, S. & Magid, N.A., et al. (2008). Evaluating Adherence to Highly Active Antiretroviral Therapy with Used of Pill Counts and Viral Load Measurement in the Drug Resources Enhancement

- against AIDS and Malnutrition Program in Mozambique. *Clinical Infectious Diseases*, 46, 1609-16.
- McAlister, A.L., Perry, C.L. & Parcel, G.S. (2008). *How Individuals, Environments, and Health Behaviors Interact: Social Cognitive Theory*. In: *Health Behavior and Health Education: Theory, Research, and Practice* 4th Edition. San Francisco, CA: John Wiley & Sons, Inc, 169–188.
- McInerney, P.A., Ncama, B.P., Wantland, D., Bhengu, B.R., McGibbon, C. & Davis, S.M., et al. (2008). Quality of life and physical functioning in HIV-infected individuals receiving antiretroviral therapy in KwaZulu-Natal, *South Africa. Nursing and Health Sciences*, 4, 266-272.
- Mellins, C.A., Brackis-Cott, E. & Dolezal, C., (2004). The role of psychosocial and family factors in adherence to antiretroviral treatment in human immunodeficiency virus-infected children. *Pediatr Infect Dis J*, 23(11); 1035-41.
- Merzel, C., VanDevanter, N. & Irvine, M. (2008). Adherence to antiretroviral therapy among older children and adolescents with HIV: a qualitative study of psychosocial contexts. *AIDS Patient Care STDS*, 22(12); 977-87.
- Miller & Katherine. (2005). *Communication Theories: Perspectives, Processes, and Contexts* (2nd ed.). New York, New York: McGraw-Hill.
- Miller, N.E. & Dollard, J. (1941). *Social Learning and Imitation*. New Haven: Yale University Press.
- Mills, E., Nachega, J., Bangsberg, D., Singh, S., Rachlis, B. & Wu, P., et al. (2006). Adherence to HAART: a systematic review of developed and developing nation patient-reported barriers and facilitators. *PLoS Med*, 3(11); e438.
- Minzi, O., Mugoyela, V. & Gustafsson, L. (2011). Correlation between lamivudine plasma concentrations and patient self-reported adherence to antiretroviral treatment in experienced HIV patients. *Ther Clin Risk Manag*, 7, 441-6.
- Murphy, D.A., Belzer, M., Durako, S.J., Sarr, M., Wilson, C.M. & Muenz, L.R., et al. (2005). Adolescent Medicine HIV/AIDS Research Network. Longitudinal antiretroviral adherence among adolescents infected with human immunodeficiency virus. *Arch Pediatr Adolesc Med*, 159, 764-770.
- Murphy, D.A., Wilson, C.M., Durako, S.J., Muenz, L.R., Belzer, M. & Adolescent Medicine HIV/AIDS Research Network. (2001). Antiretroviral

- medication adherence among the REACH HIV-infected adolescent cohort in the USA. *AIDS Care*, 13, 27-40.
- Murphy, D.A., Greenwell, L., Hoffman, D. (2002). Factors associated with antiretroviral adherence among HIV-infected women with children. *Women Health*, 36(1); 97–111.
- Murray, L., Semrau, K., McCurley, E., Thea, D.M., Scott, N. & Mwiya, M., et al. (2009). Barriers to acceptance and adherence of antiretroviral therapy in urban Zambian women: a qualitative study. *AIDS Care*, 21(1); 78-86.
- Mutwa, P.R., Van Nuil, J.I., Asiimwe-Kateera, B., Kestelyn, E., Vyankandondera, J. & Pool, R., et al. (2013). Living situation affects adherence to combination antiretroviral therapy in HIV-infected adolescents in Rwanda: a qualitative study. *PLoS One*, 8(4); e60073. Published online 2013 April 3. doi: 10.1371/journal.pone.0060073.
- Naar-King, S., Montepiedra, G., Garvie, P., Kammerer, B., Malee, K. & Sirois, P.A., et al. (2013). Social ecological predictors of longitudinal HIV treatment adherence in youth with perinatally acquired HIV. *J Pediatr Psychol*, 38(6); 664-74.
- Nachega, J.B., Knowlton, A.R., Deluca, A., Schoeman, J.H., Watkinson, L. & Efron, A., et al. (2006). Treatment supporter to improve adherence to antiretroviral therapy in HIV-infected South African Adults: A qualitative study. *JAIDS*, 1(43); S127-S133.
- National Health Security Office (NHSO). (2011). *Country wide Sealing up Quality Improvement in HIV Care and Treatment Report 2011*. Available from URL : <http://www.cqihiiv.com>. Accessed: 28 June 2012.
- Ncama, B.P., McInerney, P.A., Behngu, B.R., Corless, I.B., Wantland, D.J. & Nicholas, P.K., et al. (2008). Social support and medication adherence in HIV disease in KwaZulu-Natal, South Africa. *Intl J Nur Studies*, 45, 1757-1763.
- Nicholson, O., Mellin, C., Dolezal, C., Brackis-Cott, E. & Abrams, E.J.. (2006). HIV treatment-related knowledge and self-efficacy among caregivers of HIV-infected children. *Patient Educ Couns*, 61(3); 405-10.

- Nordqvist, O., Sodergard, B., Tully, M.P., Sönnernborg, A. & Lindblad, A.K. (2006). Assessing and achieving readiness to initiate HIV medication. *Patient Educ Couns*, 62, 21–30.
- Oberdorfer, P. (2006). *Disclosure of HIV/AIDS status: Update on Pediatric Infectious Diseases 2006*. Bangkok; Rungsin Press, Thailand, 133-9. [Thai Language]
- Ogden, J. (2000). *Health Psychology: A textbook*. Groot Brittanje: Biddles.
- Otis Dudley Duncan. (1966). Path analysis: Sociological examples. *The American Journal of Sociology*, 72(1); 1-16.
- Overby, K.J. & Kegeles, S.M. (1994). The impact of AIDS on an urban population of high-risk female minority adolescents: implications for intervention. *J Adolesc Health*, 15(3); 216-27.
- Pallela, F., Delaney, K. & Moorman, A. (1998). Declining morbidity and mortality among patients with advanced human immunodeficiency virus infection. *N Engl J Med*, 338(13); 853-860.
- Panupak, P., Leechawengwong, M. & Siraprapasiri, T. (2010). *National Guidelines on HIV/AIDS Diagnosis and treatment, Thailand 2010*. Bangkok; Agricultural Credit Cooperatives of Thailand Printing Company Limited, Thailand. [Thai Language]
- Paterson, D.L., Swindells, S., Mohr, J., Brester, M., Vergis, E.N. & Squier, C., et al. (2002). Adherence to protease inhibitor therapy and outcomes in patients with HIV infection. *Ann Intern Med*, 133, 21-30.
- Peltzer, K. & Pengpid, S. (2013). Socioeconomic factors in adherence to HIV therapy in low- and middle-income countries. *J Health Popul Nutr*, 31(2);150-70.
- Pequegnat, W., Page, B., Strauss, A., Amaro, H., Goldstein, P. & Fritz, R.B. et al. (1995). *Qualitative Inquiry: An underutilized strategy in AIDS Research*. New York:Plenum Press.
- Peter, J.C., Tuba, M. & Fylkesnes, K. (2011). Conceptual models for Mental Distress among HIV-infected and uninfected individuals: A contribution to clinical practice and research in primary-health-care centers in Zambia. *BMC Health Services Research*, 11, 7. Available from URL : <http://www.biomedcentral.com/1472-6963/11/7>. Accessed: 11 Nov 2011.

- Petosa, R. & Wessinger, J. (1990). The AIDS education needs of adolescents: a theory-based approach. *AIDS Educ Prev*, 2(2); 127-36.
- PLATO II ;The Pursuing Later Treatment Options II, project team for the Collaborations of Observational HIV Epidemiological Research Europe (COHERE). (2011). Risk of triple-class virological failure in children with HIV: a retrospective cohort study. *Lancet*, 7(377); 1580-1587.
- Raboud, J.M., Rae, S., Hogg, R.S., Yip, B., Sherlock, C.H. & Harrigan, P.R., et al. (1999). Suppression of the plasma viral load below the detection limit of a human immunodeficiency virus kit is associated with longer virologic response than suppression below the limit of quantitation. *J Infect Dis*, 180, 1347-50.
- Rachlis, B.S., Mills, E.J. & Cole, D.C. (2011). Livelihood security and adherence to antiretroviral therapy in low and middle income settings: a systematic review. *PLoS One*, 6(5):e18948. doi: 10.1371/journal.pone.0018948.
- Rathanarat, P. & Deewong, S. (2009). *The Epidemiology of AIDS Institute ;Thailand. Disease Surveillance Report 2009*. Available from URL: http://epid.moph.go.th/Annual/Annual%202552/AESR52_Part1/B_Part1_52/2952_AIDS.doc. Accessed: 17 June 2011.
- Reisner, S.L., Mimiaga, M.J., Skeer, M., Perovich, B., Johnson, C.V. & Safren, S.A. (2009). A review of HIV antiretroviral adherence and intervention studies among HIV-infected youth. *Top HIV Med*, 17(1); 14-25.
- Reynolds, N.R., Testa, M.A., Marc, L.G., Chesney, M.A., Neidig, J.L. & Smith, S.R., et al. (2004). Factors influencing medication adherence beliefs and self-efficacy in persons naive to antiretroviral therapy: a multicenter, cross-sectional study. *AIDS Behav*, 8(2); 141–150.
- Richard, T.L., Paul, R., Edward, J.M., Kariri, A., Karanja, S. & Chung, M.H., et al. (2010). Effects of a mobile phone short message service on antiretroviral treatment adherence in Kenya (WelTel Kenya1): a randomised trial. *The Lancet*, 376, 1838-1845.
- Rochon, D., Ross, M.W., Looney, C., Nepal, V.P., Price, A.J. & Giordano, T.P. (2011). Communication strategies to improve HIV treatment adherence. *Health Commun*, 26(5); 461-7.

- Rosenstock, I., Strecher, V., & Becker, M. (1994). *The Health Belief Model and HIV risk behavior change*. In R.J. DiClemente, and J.L. Peterson (Eds.), *Preventing AIDS: Theories and Methods of Behavioral Interventions* (pp. 5-24). New York : Plenum Press.
- Roux, P., Kouanfack, C., Cohen, J., Marcellin, F., Boyer, S. & Delaporte, E., et al. (2011). Adherence to antiretroviral treatment in HIV-positive patients in the Cameroon context: promoting the use of medication reminder methods. *J Acquir Immune Defic Syndr*, 57, Suppl 1, S40-3.
- Sanjobo, N., Frich, J. & Fretheim, A. (2008). Barriers and facilitators to patients' adherence to antiretroviral treatment in Zambia: a qualitative study. *SAHARA J*, 5(3); 136-143.
- Santer, M., Ring, N., Yardley, L., Geraghty, A.W. & Wyke, S. (2014). Treatment non-adherence in pediatric long-term medical conditions: systematic review and synthesis of qualitative studies of caregivers' views. *BMC Pediatr*, 4 (14); 14-63.
- Sarna, A., Luchters, S., Geibel, S., Munyao, P. & Kaai, S. (2005). *Promoting adherence to antiretroviral therapy through a directly administered antiretroviral therapy (DAART) strategy in Mombasa, Kenya*. Nairobi: Population Council.
- Sarna, A. & Weiss, E. (2007). *Current research and good practice in HIV and AIDS treatment education (UNESCO and WHO, 2007)*. Available from: URL:<http://unesdoc.unesco.org/images/0014/001497/149722E.pdf>. Accessed: 1 Feb 2012.
- Sayles, J., Wong, M., Kinsler, J., Martins, D. & Cunningham, W.E. (2009). The association of stigma with self-reported access to medical care and antiretroviral therapy adherence in persons living with HIV/AIDS. *J Gen Intern Med*. in press.
- Scheper-Hughes & Nancy. (1990). "Three Propositions for a Critically Applies Medical Anthropology." *Social Science Medicine*, 30, 189-197.
- Shegog, R., Maekham, C.M., Leonard, A.D., Bui, T.C. & Paul, M.E. (2012). +CLICK: pilot of web-based training program to enhance ART adherence among HIV-positive youth. *AIDS Care*, 24(3); 310-8.
- Shuter, J., Sarlo, J., Kanmaz, T., Rode, R.A. & Zingman, B.S. (2007). HIV-infected patients receiving lopinavir/ritonavir-based antiretroviral therapy achieve high

- rates of virologic suppression despite adherence rates less than 95%. *J Acquir Immune Defic Syndr*, 45(1); 4-8.
- Sibhabu, B., Amare, D., Alemayehu, A. & Deribe, K. (2008). Adherence to highly active antiretroviral therapy and its correlates among HIV infected pediatric patients in Ethiopia. *BMC Pediatric*, 8, 53.
- Sivapalasingam, S., Mendillo, M., Ahmed, A., Mwamzuka, M., Said, S. & Marshed, F., et al. (2014). The importance of caregivers in the outcome of pediatric HIV management, Mombasa, Kenya. *AIDS Care*, 26(4); 425-33.
- Skinner, D. & Mfecane, S. (2004). Stigma, discrimination and the implications for people living with HIV/AIDS in South Africa. *SAHARA J*, 1(3); 157-164.
- Stein, L.A., Minugh, P.A., Longabaugh, R., Wirtz, P., Baird, J. & Nirenberg, Ted D., et al. (2009). Readiness to change as a mediator of the effect of a brief motivational intervention on post treatment alcohol-related consequences of injured emergency department hazardous drinkers. *Psychol Addict Behav*, 23, 185–195.
- Sterne, J.A., May, M., Costagliola, D., de Wolf, F., Phillips, A. & Harris, R., et al. (2009). Timing of initiation of antiretroviral therapy in AIDS-free HIV-1-infected patients: a collaborative analysis of 18 HIV cohort studies. *Lancet*, 373, 1352-63.
- Steven, A.S., Otto, M.W., Worth, J.L., Salomon, E., Johnson, W. & Mayer, K., et al. (2001). Two strategies to increase adherence to HIV antiretroviral medication: Life-Steps and self-medication monitoring. *Behaviour Research and Therapy*, 39, 1151-1162.
- Sudjarittrak, T. & Oberdorfer, P. (2011). *Adolescent HIV Care Management from mother to child transmission: Update on Pediatric Infectious Diseases 2011*. Bangkok; Beyond Enterprise Co.,Ltd., 212-228. [Thai Language]
- Sungkanuparph, S., Techasathit, W. & Utaipiboon, C. (2010). Thai national guidelines for antiretroviral therapy in HIV-1 infected adults and adolescents 2010. *Asian Biomedicine*, 4(4); 515-528.
- Surin Hospital. (2011). *Immunology clinic report : Surin Hospital Report*. 21.
- Taddese, A., Sibhabu, B., Kebede, D. & Escudero, H.R. (2013). Experience of stigma and discrimination and the implications for healthcare seeking behavior among people living with HIV/AIDS in resource-limited setting. *SAHARA J*, 10(1); 1–7.

- Trocme, N., Vaudre, G., Dollfus, C. & Leverger, G. (2002). Factors impacting on antiretroviral therapy compliance in HIV positive adolescents. *Archives de Pediatrie*, 9, 1241-1247.
- Tucker, J.S., Burnam, M.A., Sherbourne, C.D., Kung, F.Y. & Gifford, A.L. (2003). Substance use and mental health correlates of nonadherence to antiretroviral medications in a sample of patients with human immunodeficiency virus infection. *Am J Med*, 114, 573-580.
- Tuldra, A., Fumaz, C.R., Ferrer, M.J., Bayés, R., Arnó, A. & Balagué, M., et al. (2000). Prospective randomized two-arm controlled study to determine the efficacy of a specific intervention to improve long-term adherence to highly active antiretroviral therapy. *J Acquir Immune Defic Syndr Hum Retroviral*, 25, 221-8.
- Udompanich, T., Kosalaraksa, P. & Ratanamanee, S. (2008). Duration of Antiretroviral Therapy and Adherence in HIV-Infected Children at Srinagarind Hospital. *Srinagarind Med J*, 23(3).
- UNGASS. (2008). Geneva, Switzerland: *UNAIDS; 2008. Country Progress Report, Thailand*, Reporting period January 2006-December 2007.
- Usitalo, A., Leister, E., Tassiopoulos, K., Allison, S., Malee, K. & Paul, M.E., et al. (2014). Relationship between viral load and self-report measures of medication adherence among youth with perinatal HIV infection. *AIDS Care*, 26(1); 107-15.
- Ugwu, R. & Eneh, A. (2013). Factors influencing adherence to paediatric antiretroviral therapy in Portharcourt, South- South Nigeria. *Pan Afr Med J*, 2013 Sep 29;16:30. doi: 10.11604/pamj.2013.16.30.1877. eCollection 2013.
- Venanzio, V., Thiloshini, G., Sicelo, D., Taylor, M., Moodley, I. & David, V., et al. (2010). Retrospective study on the critical factors for retaining patients on antiretroviral therapy in KwaZulu-Natal, South Africa. *J Acquir Immune Defic Syndr*, 55(1); 109-116.
- Vergidis, P., Falaga, M. & Hamer, D. (2009). Meta-analytical studies on the epidemiology, prevention, and treatment of human immunodeficiency virus infection. *Infect Dis Clin North Am*, 23(2); 295-308.
- Vissman, A.T., Hergenrather, K.C., Rojas, G., Langdon, S.E., Wilkin, A.M. & Rhodes, S.D. (2011). Applying the theory of planned behavior to explore HAART

- adherence among HIV-positive immigrant latinos: Elicitation interview results. *Patient Educ Couns*, 85(3); 454-60.
- Viswanathan, H., Anderson, R. & Thomas, J. (2005). Evaluation of an antiretroviral medication attitude scale and relationships between medication attitudes and medication non-adherence. *AIDS patient care and STDs*, 19, 306-316.
- Vlassoff, C., & Ali, F. (2011). HIV-related stigma among South Asians in Toronto. *Ethnicity Health*, 16(1); 25-42.
- Vreeman, R.C., Nyandiko, W.M., Ayaya, S.O., Walumbe, E.G., Marrero, D.G. & Inui, T.S. (2009). Factors sustaining pediatric adherence to antiretroviral therapy in western Kenya. *Qual Health Res*, 19(12); 1716-29.
- Vreeman, R.C., Wiehe, S.E., Pearce, E.C. & Nyandiko, W.M. (2008). A systematic review of pediatric adherence to antiretroviral therapy in low- and middle-income countries. *Pediatr Infect Dis J*, 27(8); 686-91.
- Walsh, J.C., Mandalia, S. & Gazzard, B.G. (2002). Responses to a 1 month self-report on adherence to antiretroviral therapy are consistent with electronic data and virological treatment outcome. *AIDS*, 16, 269-277.
- Washing, C.H. & Oberdorfer, P. (2009). *Stigma in perinatally HIV-infected adolescents* [Abstract No.WEPEC102]. 5th IAS Conference on HIV Pathogenesis, Treatment and Prevention. Cape Town, South Africa: International AIDS Society.
- Weinberg, Robert S. & Gould. (2007). *Foundation of Sport and Exercise Psychology* 4th Edition, 422.
- Weiss, L., French, T., Finkelstein, R., Waters, M., Mukherjee, R. & Agins, B. (2003). HIV- related knowledge and adherence to HAART. *AIDS Care*, 15, 673-679.
- World Health Organization. (2003). *Adherence to long-term therapies: Evidence for action*. Geneva.
- World Health Organization. (2009). *Rapid advice: antiretroviral therapy for HIV infection in adults and adolescents*, 1-24. Available at http://www.who.int/HIV/pub/arv/rapid_advice_art.pdf. Accessed 20 Dec 2011.
- World Health Organization. (2010). *Antiretroviral Therapy for HIV infection in adults and adolescents ; Recommendations for a public health approach*, 2010 revision.

- Wouters, E., Van Damme, W. & Van Loon, F. (2009). Public-sector ART in the Free State Province, South Africa: Community support as an important determinant of outcome. *Soc Sci Med*, 20, 415-421.
- Zackie, A., David, A., Drew, A. & (Global HIV Prevention Working Group). (2008). *Behavior change and HIV prevention*. Available from URL : http://www.globalhivprevention.org/pdfs/PWG_behavior%20report_FINAL.pdf. Accessed: 19 Dec 2011.
- Zolopa, A., Andersen, J., Powderly, W., Sanchez, A., Sanne, I. & Suckow, C., et al. (2009). Early antiretroviral therapy reduces AIDS progression/death in individuals with acute opportunistic infections: a multicenter randomized strategy trial. *PLoS One*, 4, e5575.
- Zulkifli, S.N. & Wong, Y.L. (2002). Knowledge, attitudes and beliefs related to HIV/AIDS among adolescents in Malaysia. *Med J Malaysia*. 57(1); 3-23.