

REFERENCES

- [1] Visscher C.F., Klein G., Verspohl J., Beyerbach M., Stratmann-Selke J., Kamphues J., “Serodiversity and serological as well as cultural distribution of *Salmonella* on farms and in abattoirs in Lower Saxony, Germany,” International Journal of Food Microbiology, Vol. 146, No. 1. 15 Mar 2011, pp. 44–51.
- [2] Giovannini A., Prencipe V., Conte A., Marino L., Petrini A., Pomilio F., Rizzi V., Migliorati G., “Quantitative risk assessment of *Salmonella* spp. infection for the consumer of pork products in an Italian region,” Food Control, Vol. 15, No. 2. Mar 2004, pp. 139–144.
- [3] Rostagno M.H., Callaway T.R., “Pre-harvest risk factors for *Salmonella enterica* in pork production,” Food Reserch International, Vol. 45, No.2. Mar 2012, pp. 634–640.
- [4] Bureau of Epidemiology, “Annual epidemiological surveillance report,” Available at: http://www.boe.moph.go.th/boedb/d506_1/index.php ,accessed on 24 Jul 2013.
- [5] Berends B.R., Van Knapen F., Mossel D.A., Burt S., Snijders J.M., “Impact on human health of *Salmonella* spp. on pork in The Netherlands and the anticipated effects of some currently proposed control strategies,” International Journal of Food Microbiology, Vol.44, No.3. 10 Nov 1998, pp. 219–229.
- [6] D’Aoust J.Y., Maurer J., Bailey J.S., “Food Microbiology Fundamentals and Frontiers.” 2001, ASM Press, Washington DC, USA.
- [7] Jawetz E., Melnick J.L., Adelberg E.A. “Review of Medical Microbiology,” Appleton & Lange, Norwalk, Conn, 1954, pp. 244-248.
- [8] WHO., “Risk assessment of *Salmonella* in eggs and broiler chickens.” 2002, Food and Agriculture Organization of the United Nations and World Health Organization, Geneva, Switzerland.
- [9] Patrick A.D., Weill F.X. “Antigenic formula of *Salmonella* serovars,” WHO Collaborating Center for Reference and Research on *Salmonella*. 2007, pp1-66.

- [10] Brenner F.W., Villar R.G., Angulo R., Swaminathan B., “*Salmonella* nomenclature,” *Journal of Clinical Microbiology*, Vol.38, No. 7. July 2000, pp. 2465-2467.
- [11] Popoff M.Y., Bockemuhl J., Brenner F.W., Gheesling L.L., “Supplement 2000 (no. 44) to the Kauffmann-White scheme,” *Research Microbiology*. Vol. 152, No. 10. Dec 2011, pp. 907-909.
- [12] Liu W., Liu B., Zhu X., Yu S., Shi X., “Diversity of *Salmonella* isolates using serotyping and multilocus sequence typing,” *Food Microbiology*. Vol. 28, No. 6. Sep 2011, pp. 1182–1189.
- [13] Baggesen D.L., Sørensen G., Nielsen E.M., Wegener H.C., “Phage typing of *Salmonella* Typhimurium – is it still a useful tool for surveillance and outbreak investigation?,” *Eurosurveillance*, Vol. 15, No. 4. 28 Jan 2010, pp. 1-3
- [14] Rabsch W., “*Salmonella* Typhimurium phage typing for pathogens,” *Methods in Molecular Biology*, Vol. 394, 2007, pp. 177-211.
- [15] Chmielewski R., Wieliczko A., Kuczkowski M., Mazurkiewicz M., Ugorski M., “Comparison of ITS profiling, REP- and ERIC-PCR of *Salmonella* Enteritidis isolates from Poland,” *Zoonoses and Public health*, Vol. 49, No.4. May 2002. pp. 163-168.
- [16] López – Molina N., Laconcha I., Rementería A., Audicana A., Perales I., Garaizar J., “Typing of *Salmonella* Enteritidis of different phage types of PCR fingerprinting,” *Journal of Applied Microbiology*, Vol. 84, No. 5. May 1998, pp. 877-882.
- [17] Torpdahl M., Skov M.N., Sandvang D., Baggesen D.L., “Genotypic characterization of *Salmonella* by multilocus sequence typing, pulsed-field gel electrophoresis and amplified fragment length polymorphism,” *Journal of Microbiological Methods*, Vol.63, No. 2005, pp. Nov 2005, pp. 173–184.
- [18] Campioni F., Bergamini A.M.M., Falcão J.P., “Genetic diversity, virulence genes and antimicrobial resistance of *Salmonella* Enteritidis isolated from food and humans over a 24-year period in Brazil,” *Food Microbiology*, Vol. 32, No.2 . Dec 2012, pp. 254–264.
- [19] Foxman B., “*Molecular Tools and Infectious Disease Epidemiology*,” 1st ed, Academic Press, United States of America, 2012, pp. 23-40.

- [20] Prendergast D.M., Duggan S.J., Gonzales-Barron U., Fanning S., Butler F., Cormican M., Duffy G., "Prevalence, numbers and characteristics of *Salmonella* spp. on Irish retail pork," International Journal of Food Microbiology, Vol. 131, No. 2. 31 May 2009, pp. 233–239.
- [21] Hauser E, Hebner F, Tietze E, Helmuth R, Junker E, Prager R, Schroeter A., Rabsch W., Fruth A., Malorny B., "Diversity of *Salmonella* enterica serovar Derby isolated from pig, pork and humans in Germany," International Journal of Food Microbiology, Vol. 151, No.2, pp. 141–149.
- [22] Dorn-in S., Fries R., Padungtod P., Kyule M.N., Baumann M.P., Srikitjakarn, L., Chantong W., Sanguangiat A., Zessin K.H. 2009, "A cross-sectional study of *Salmonella* in pre-slaughter pigs in a production compartment of northern Thailand," Preventive Veterinary Medicine, Vol. 88, No.1. 1 Jan 2009, pp. 15–23.
- [23] Farzan A., Friendship R.M., Dewey C.E., Warriner K., Poppe C., Klotins K., "Prevalence of *Salmonella* spp. on Canadian pig farms using liquid or dry-feeding," Preventive Veterinary Medicine, Vol. 73, No. 4. 16 Mar 2006, pp. 241–254.
- [24] Swanenburg M., Urlings H.A.P., Snijders J.M.A., Keuzenkamp D.A., van Knapen F., "*Salmonella* in slaughter pigs: prevalences, serotypes and critical control points during slaughter in two slaughterhouses," International Journal of Food Microbiology, Vol. 70, No.3. 8 Nov 2001, pp. 243–254.
- [25] Delhalle L., Saegerman C., Farnir F., Korsak N., Maes D., Messens W., De Sadeleer L., De Zutter, L., "*Salmonella* surveillance and control at post-harvest in the Belgian pork meat chain," Food Microbiology, Vol. 26, No. 3. May 2009, pp.265–271.
- [26] Gomes-Neves E., Antunes P., Tavares A., Themudo P., Cardoso M.F., Gärtner F., Costa J.M. and Peixe L., "*Salmonella* cross-contamination in swine abattoirs in Portugal: Carcasses, meat and meat handlers," International Journal of Food Microbiology, Vol. 157, No,1. Apr 2012, pp. 82–87.

- [27] Van Hoek A.H.A.M., de Jonge R., van Overbeek W.M., Bouw E., Pielaat A., Smid J.H., Malorny B., Junker E., Löfström C., Pedersen K., Aarts H.J., Heres, L., "A quantitative approach towards a better understanding of the dynamics of *Salmonella* spp. in a pork slaughter-line," International Journal of Food Microbiology, Vol. 153, No.1. 1 Feb 2012, pp. 45–52.
- [28] Mürmann L., Dos Santos M.C., Cardoso M., "Prevalence, genetic characterization and antimicrobial resistance of *Salmonella* isolated from fresh pork sausages in Porto Alegre, Brazil," Food Control, Vol. 20, No. 1. Jan 2009, pp. 191–195.
- [29] Padungtod P., Kaneene J.B., "*Salmonella* in food animals and humans in northern Thailand," International Journal of Food Microbiology, Vol. 108, No. 3. 1 May 2006, pp. 346–354.
- [30] Lo Fo Wong D.M., Hald T., Van der Wolf P., Swanenburg M., "Epidemiology and control measures for *Salmonella* in pigs and pork. Livestock Production Science, Vol.76, No. 3. Sep 2002, pp. 215–222.
- [31] Berends B.R., Van Knapen F., Snijders J.M., Mossel D.A., "Identification and quantification of risk factors regarding *Salmonella* spp. on pork carcasses," International Journal of Food Microbiology, Vol. 36, No. 2. May 1997, pp. 199–206.
- [32] Hotes S., Kemper N., Traulsen I., Rave G., Krieter J., "Risk factors for *Salmonella* infection in fattening pigs-An evaluation of blood and meat juices samples," Zoonoses and Public health, Vol. 57, No. 1. Nov 20120, pp. 30-38
- [33] Mollar K., Jensen T.K., Jorsal S.E., Leser T.D., Carstensen B., "Detection of *Lawsonia intracellularis*, *Serpulina hyodysenteriae*, weakly beta-haemolytic intestinal spirochaets, *Salmonella enterica*, and haemolytic *Escherichia coli* from swine herds with and without diarrhea among growing pigs," Veterinary Microbiology, Vol. , No. 1. 30 Apr 1998, pp.59-62.
- [34] Wills R.W. Gray J.T., Fedorka-Cray P.J., Yoon K.J., Ladely S., Zimmerman J.J., "Synergism between porcine reproductive and respiratory syndrome virus (PRRSV) and *Salmonella* Cholerasuis in swine," Veterinary Microbiology, Vol. 71, No. 3. Feb 2000, pp.81-99.

- [35] Steenhard N.R., Jensen T.K., Baggesen D.L., Roepstorff A., Moller K., "Excretion in feces and mucosal persistence of *Salmonella* ser. Typhimurium in pigs subclinically infected with *Oesophagostomum* spp.," American Journal of Veterinary Research, Vol.63, No. 1. Jan 2002, pp. 130-136.
- [36] Lo Fo Wong D.M.A., Dahl J., Stege H., van der Wolf P., Leontides L., von Altrock A., Herd-Level risk factors for subclinical *Salmonella* infection in European finishing-pig herds," Livestock Production Science, Vol. 62, No.4. 16 Apr 2004, pp. 253-266.
- [37] Schwaiger K., Huther S., Hölzel C., Kämpf P., Bauer, J., "Prevalence of antibiotic-resistant enterobacteriaceae isolated from chicken and pork meat purchased at the slaughterhouse and at retail in Bavaria, Germany," International Journal of Food Microbiology, Vol. 154, No. 3. 15 Mar 2012, pp. 206–211.
- [38] Clayton N.C., "The efficacy of various *Salmonella* intervention methods applied to pork carcasses during slaughter," Lexington, University of Kentucky Libraries, 2002.
- [39] Fernandez-Escartin E., Saldana L.J., Rodriguez O., Martinez O.N., Antonio, T.J., " Incidence and level of *Salmonella* serovars in raw pork obtained from Mexican butcher shops," Food Microbiology, Vol. 12. Feb 1995, pp. 435-439.
- [40] Berends B.R., Van Knapen F., Mossel D.A., Burt S., Snijders J.M, "*Salmonella* spp. on pork at cutting plants and at the retail level and the influence of particular risk factors," International Journal of Food Microbiology, Vol. 44, No. 3. 10 Nov 1998, pp. 207–217.
- [41] Cassin M.H., Lammerding A.M., Todd A.C.D., Ross W., McColl R.S., "Quantitative risk assessment for *Escherichia coli* O157:H7 inground beef hamburgers," International Journal of Food Microbiology, Vol. 41. 21 Jan 1998, pp. 21-44.
- [42] Kuan C.H., Wong W.C., Pui C.F., Mahyudin N.A., Tang J.Y.H., Nishibuchi M, Radu S, "Prevalence and quantification of *Listeria monocytogenes* in beef offal at retail level in Selangor, Malaysia," Brazilian Journal of Microbiology. Vol. 44, No. 4. 24 Jan 2012, pp. 1169–1172.

- [43] Gonzales-Barron U.A., Redmond G., Butler F., “A risk characterization model of *Salmonella* Typhimurium in Irish fresh pork sausages,” Food Research International, Vol. 45, No. 2. March 2012, pp. 1184–1193.
- [44] García-Feliz C., Collazos J.A., Carvajal A., Vidal A.B., Aladueña A., Ramiro R., de la Fuente M., Echeita M.A., Rubio P., “*Salmonella* enterica infections in Spanish swine fattening units,” Zoonoses and Public health, Vol. 54, No. 8. Feb 2007, pp. 294–300.
- [45] ISO 6579. Horizontal method for the detection of *Salmonella* spp. International Standard Organization, Geneva, 2002.
- [46] MPN calculator. Available at: <http://www.i2workout.com/mcuriale/mpn> (accessed on 26 May 2012).
- [47] Barber D.A., Bahnson P.B., Isaacson R., Jones C.J., Weigel R.M., “Distribution of *Salmonella* in swine production ecosystems,” Journal of Food Protection, Vol. 65, No. 12. Dec 2002, pp. 1861-1868.
- [48] Stephens P.J., Joynson J.A., Davies K.W., Holbrook R., Lappin-Scott H.M., Humphrey T.J., “The use of an automated growth analyser to measure recovery times of single heat-injured *Salmonella* cells,” Journal Applied Microbiology, Vol. 83, No. 4. 30 Oct 2003, pp.445–455.
- [49] Quintavalla S., Larini S., Mutti P., Barbuti, S., “Evaluation of the thermal resistance of different *Salmonella* serotypes in pork meat containing curing additives,” International Journal of Food Microbiology, Vol. 67, No.1. 20 Jul 2001, pp.107–114.
- [50] Patchanee P., Zessin, K.H., Staak, C., Srikijsarn, L., Taravijitkul, P., Tesaprateep T., “Pre-slaughter infection of *Salmonella* spp. and consideration of using the DANISH MIX-ELISA for monitoring *Salmonella* in pigs,” Chiang Mai Veterinary Journal, Vol. 1, No. 1. Oct 2002, pp. 33-38.
- [51] Sandberg M., Hopp P., Jarp J., Skjerve E., “An evaluation of the Norwegian *Salmonella* surveillance and control program in live pig and pork,” International Journal of Food Microbiology, Vol. 72, No. 1. 30 Jan 2002, pp. 1–11.

- [52] Sutton S., The Most Probable Number method and it's use in enumeration, quantification and validation," Journal of Validation Technology, Vol.16, No. 3. 2010, pp.35-38.
- [53] Vose D., " Risk Analysis: A Quantitative Guide," 2nd ed., Wiley John & Sons, Incorporated, New York.
- [54] Majowicz S.E., Musto J., Scallan E., Angulo F.J., Kirk M., O'Brien S.J., Jones T.F., Fazil A., Hoekstra R.M., International Collaboration on Enteric Disease "Burden of Illness" Studies, "The global burden of nontyphoidal *Salmonella* gastroenteritis," Clinical Infectious Disease: An Official Publication of the Infectious Diseases Society of America, Vol. 50, No. .15 Mar 2012, pp. 882–889.
- [55] Sirichote P., Bangtrakulnonth A., Tianmanee K., Unahalekhaka A., Oulai A, Chittaphithakchai P., Kheowrod W., Hendriksen R.S., "Serotype and antimicrobial resistance of *Salmonella enterica* spp. in central Thailand, 2001-2006," Southeast Asian Journal Tropical Public Health, Vol. 41, No. 6. November 2010, pp. 1405-1415.
- [56] Pulsrikarn C., Chaichana P., Pornruangwong S., Morita Y., Yamamoto S., Boonmar S., "Serotype, antimicrobial susceptibility, and genotype of *Salmonella* Isolates from swine and pork in Sa Kaew province, Thailand," The Thai Journal of Veterinary Medicine, Vol. 42, No. 1. 2012, pp. 21–28.
- [57] Tadee P., Kumpapong K., Sinthuya D., Yamsakul .P, Chokesajjawatee N., Nuanualsuwan S., Pornsukarom S., Molla B.Z., Gebreyes W.A., Patchanee P., "Distribution, quantitative load and characterization of *Salmonella* associated with swine farms in upper-northern Thailand. Journal of Veterinary Science, Vol. 15, No. 2. Jun 2012, pp. 327–334.
- [58] Van Boxstael S., Dierick K., Van Huffel X., Uyttendaele M., Berkvens D., Herman L., Bertrand S., Wildemauwe C., Catry B., Butaye P., Imberechts H., "Comparison of antimicrobial resistance patterns and phage types of *Salmonella* Typhimurium isolated from pigs, pork and humans in Belgium between 2001 and 2006," Food Reserch International, Vol. 45, No.2. March 2012, pp. 913–918.

- [59] Boonmar S., Bangtrakulnonth A., Pornrunangwong S., Terajima J. Watanabe H., Kaneko K.I., Ogawa M., “Epidemiological analysis of *Salmonella* enteritidis isolates from humans and broiler chickens in Thailand by phage typing and pulsed-field gel electrophoresis,” *Journal of Clinical Microbiology*, Vol. 36, No. 4. Apr 1998, pp. 971–974.
- [60] Ridley A.M., Threlfall E.J., Rowe B., “Genotypic characterization of *Salmonella* Enteritidis phage types by plasmid analysis, ribotyping, and pulsed-field gel electrophoresis. *Journal of Clinical Microbiology*, Vol. 36, No. 8. Aug 1998, pp. 2314–2321.
- [61] Aktas Z., Day M., Kayacan C.B., Diren S., Threlfall E.J., “Molecular characterization of *Salmonella* Typhimurium and *Salmonella* Enteritidis by plasmid analysis and pulsed-field gel electrophoresis,” *International Journal of Antimicrobial Agents*, Vol. 30, No. 6. Dec 2007, pp.541–545.
- [62] Maslow J.N., Mulligan M.E., Arbeit R.D., “Molecular epidemiology: application of contemporary techniques to the typing of microorganisms,” *Clinical Infectious Disease: An Official Publication of the Infectious Diseases Society of America*, Vol. 17, No. 2, Aug 1993, pp. 153–162.
- [63] Clinical and Laboratory Standards Institute (CLSI), Performance standards for antimicrobial disc susceptibility test; Approved standard-Tenth Edition. CLSI document M02-A10. Wayne, PA: Clinical and Laboratory Standards Institute, 2011
- [64] Ribot E.M., Fair M.A., Gautam R., Cameron D.N., Hunter S.B., Swaminathan B., Barrett T.J., “Standardization of pulsed field gel electrophoresis protocols for the subtyping of *Escherichia coli* O157:H7, *Salmonella* and *Shigella* for PulseNet,” *Foodborne Pathogen and Disease*, Vol. 3, No. 1. 2006, pp. 59-67.
- [65] Hunter P.R., Gaston M.A., “Numerical index of the discriminatory ability of typing system: an application of Simpson’s index of diversity,” *Journal of Clinical Microbiology*, Vol. 26, No. 11. Nov 1998, pp. 2465-2466.
- [66] Phillips I., Casewell M., Cox T., De Groot B., Friis C., Jones R., Nightingale C., Preston R., Waddell J. “Does the use of antibiotics in food animals pose a risk to human health? A critical review of published data,” *Journal of Antimicrobial Chemotherapy*, Vol. 53, No. 1. Jan 2004, pp. 28–52.

- [67] Lu Y, Zhao H, Sun J, Liu Y, Zhou X, Beier R.C., Wu G., Hou X., “Characterization of multidrug-resistant *Salmonella enterica* Serovars Indiana and Enteritidis from chickens in eastern China.” PLOS ONE, Vol. 9, No. 5. 2 May 2014 : e(96050).
- [68] Mc Mahon M.S., Blair I.S., Moore J.E., Mc Dowell D.A., “The rate of horizontal transmission of antibiotic resistance plasmids is increased in food preservation-stressed bacteria,” Journal Applied Microbiology Vol. 103, No. 5. Nov 2007, pp. 1883–1888.
- [69] Soyer Y., Switt A.M., Davis M.A., Maurer J., McDonough P.L., Schoonmaker-Bopp D.J., Dumas N.B., Root T., Warnick L.D., Gröhn Y.T., Wiedmann M., “*Salmonella enterica* serotype 4,5,12:i:-, an emerging *Salmonella* serotype that represents multiple distinct clones,” Journal of Clinical Microbiology, Vol. 47, No. 11. Nov 2009, pp. 3546-3556.
- [70] Kuo H.C., Lauderdale T.L., Lo D.Y., Chen C.L., Chen P.C., Liang S.Y., Kuo J.C., Liao Y.S., Liao C.H., Tsao C.S., Chiou C.S., “An association of genotypes and antimicrobial resistance patterns among *Salmonella* Isolates from pigs and humans in Taiwan,” PLOS ONE, Vol. 9, No.4. 23 Apr 2014, e(95772).

LIST OF PUBLICATION

- [1]. Tadee P, Kumpapong K, Sinthuya D, Yamsakul P, Chokesajjawatee N, Nuanualsuwan S, Pornsukarom S, Molla BZ, Gebreyes WA, Patchanee P, “Distribution, quantitative load and characterization of *Salmonella* associated with swine farms in upper-northern Thailand,” Journal of Vet Sciences, Vol. 15 No. 2. Jun2014, Page No. 327–334.
- [2]. Kumpapong K, Sinthuya D, Tadee P, Boonkhot P, Patchanee P, “Risk factor for the herd level of *Salmonella* in swine production, northern Thailand,” 52nd Kasetsart university annual conference, 2014, Page No.193-201.
- [3]. Rungrudeesombatkit R, Sukroongruang P, Choksajjawatee N, Pornaem S, Santiyanont P, Boonkhot P, Tadee P, Patchanee P. Repetitive Sequence-Based PCR (rep-PCR) technique in differentiation of *Salmonella* serotype Rissen and Typhimurium from swine farm of northern Thailand. 52nd Kasetsart university annual conference, 2014, Page No. 202-210.
- [4]. Tadee P, Boonkhot P, Kumpapong K, Patchanee P, “Prevalence and contamination level of *Salmonella* in slaughtering processes in Chiang Mai and Lamphun,” 52nd Kasetsart university annual conference, 2014, Page No. 211-221.
- [5]. Kumpapong K, Sinthuya D, Tadee P, Boonkhot P, Patchanee P, “Serotypes and Antimicrobial Resistance of *Salmonella* spp.from Swine Breeder in Chiang Mai-Lamphun Infectious disease,” Thai Journal of Animal Science, Vol.1 No.2. 2014, Page No. 139-142.
- [6]. Patchanee P, Tadee P, Ingkaninan P, Tankaew P, Hoet AE, Chupia V, “Distribution and characterization of methicillin-resistant *Staphylococcus aureus* (MRSA) at the small animal hospital, faculty of veterinary medicine, Chiang Mai university, Chiang Mai, Thailand,” The South East Asian journal of tropical medicine and public health. Vol. 45 No. 2. Mar 2014, Page No. 413-420.

- [7]. Patchanee P, Tadee P, Arjkumpa O, Love D, Chanachai K, Alter T, Hinjoy S, Tharavichitkul P, "Occurrence and Characterization of Livestock Associated-Methicillin Resistant *Staphylococcus aureus* in Pig Industries of Northern Thailand," Journal of Vet Sciences, 20 Jun 2014.
- [8]. Tadee P, Boonkhot P, Patchanee P, "Quantification and expected numbers of *Salmonella* spp. in Slaughter pigs in Chiang Mai and Lamphun, Thailand," Japanese journal of Veterinary Reserch.
- [9]. Chotinun S, Rojanasthien S, Unger F, Tadee P, Patchanee P, "Prevalence and antimicrobial resistance of *Salmonella* isolated from carcasses, processing and environment of small scale poultry slauterhouses in Thailand," The South East Asian journal of tropical medicine and public health. Vol. 45 No. 6. Jan 2014.
- [10]. Chotinun S, Rojanasthien S, Unger F, Suwan M, Tadee P, Patchanee P, "An integrative approach to enhancing small-scale poultry slaughterhouses by addressing regulations and food safety in northern-Thailand," Infectious Disease of Poverty.
- [11]. Tadee P, Boonkhot P, Patchanee P, "Comparative phenotypic and genotypic characterization of *Salmonella* spp. in pig production and processing in two provinces in Thailand," PLOS ONE.