

# CONTENTS


	Page
Acknowledgements	c
Abstract in Thai	d
Abstract in English	f
List of Tables	k
List of Figure	l
Chapter 1 Introduction	1
1.1 Introduction	1
1.2 Literature review	2
1.2.1. Cold Atmospheric Pressure Plasma Jets	2
1.2.2. Plasma interaction on DNA	3
1.3 Research objectives	4
1.4 Usefulness of research	4
1.5 Methodology and scope	5
Chapter 2 Theory	6
2.1 Plasma definition	6
2.2 Plasma Sheath (Debye shielding)	8
2.3 Cold Atmospheric Pressure Plasma Jets	10

2.4 Measurement of Plasma energy by using V-Q lissajous method	11
2.5 Measurement of Plasma species by using Optical Emission Spectroscopy (OES)	13
2.6 Interaction between energetic ions with DNA	16
2.6.1. Plasmid DNA used	16
2.6.2. Production of free radicals	18
2.6.3. DNA damage and mutation	19
2.6.4. Measurement of Naked DNA concentration	23
2.6.5. Gel electrophoresis	24
2.6.6. Electroporation	27
Chapter 3 Experiment	30
3.1. Cold atmospheric pressure plasma systems	30
3.2. Measurement of the Cold Atmospheric Pressure Plasma	33
Generating energy and generating power	
3.3. Optical Emission Spectroscopy	35
3.4. Biological preparation	36
3.4.1. Naked DNA extraction	36
3.4.2. Measurement of naked DNA concentration	37
3.4.3. Competent cell preparation	38
3.5. Plasma treated on Naked DNA	40

3.6. Gel electrophoresis for analyzing plasmid DNA form	42
3.7. Electroporation for transferring plasmid DNA to competent cell	43
3.8. Observation and Selection of DNA mutation	44
Chapter 4 Result & Discussion	46
4.1. Plasma power	46
4.2. Plasma species result (OES)	53
4.3. Gel electrophoresis (Changed plasmid form)	56
4.4. Mutation	60
Chapter 5 Conclusion	63
5.1 Cold atmospheric pressure plasma jet power	62
5.2. Plasma species (optical emission spectroscopy: OES)	63
5.3 Changed plasmid DNA form	63
5.4 Bacterial Mutation	63
References	65
Appendices	69
Appendix A Plasmid purification	69
Appendix B Plasmid DNA concentration	71
Appendix C Electrophoresis	73
Appendix D Electroporation	78
Curriculum Vitae	80

## LIST OF TABLES

	Page
Table 4.1. Summary of plasma energy measurement	51
Table 4.2. Summary of plasma power measurement	52
Table.4.3. Percentage of light intensity is result of DNA change form	57
Table.4.4. Percentage of light intensity is result of DNA change form, the dry-wet condition and electric field condition	59



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่  
Copyright© by Chiang Mai University  
All rights reserved

## LIST OF FIGURE

	Page
Figure 2.1. Plasma behaviors when plasma is in close system, electron will move to surface chamber and collecting on surface become.	8
Figure 2.2. Plasma sheath occur at contract area near floating target, electrons move to folding target faster than ion. Electrons collect continuously until electron potential equal floating potential, electric field is established.	9
Figure 2.3. Plasma boundary sheath, typical potential change in dependence on the distance from the surface.	9
Figure.2.4. Schematic diagram of a plasma jet output consists of a plastic nozzle with a needle electrode inside.	10
Figure 2.5. The plasma generator can adjust plasma power (radiofrequency – power) and frequency (plasma pluses)	11
Figure 2.6. Schematic diagram of lissajous pattern from the same sinusoids frequency by varying phase relationships.	12
Figure 2.7. Mechanism of plasma light emission. Excess energy, $E$ , is released as the electron decays from a higher energy level to a lower one.	14
Figure.2.8. Wavelength of light emission spectra of Ne and Hg.	15
Figure 2.9. Schematic diagram of optical emission spectroscopy	15

Figure 2.10. The overall shape of the protein and its association into dimers. Eleven strands of $\beta$ -sheet (green) form the walls of a cylinder. Short segments of helices (blue) cap the top and bottom of the ' $\beta$ -can' and also provide a scaffold for the fluorophore which is near geometric center of the can. This folding motif, with $\beta$ -sheet outside and helix inside, represents a new class of proteins. Two monomers are associated into a dimer in the crystal and in solution at low ionic strengths. This view is directly down the two-fold axis of the non-crystallographic symmetry	17
Figure 2.11. There are three forms of the plasmid DNA: supercoiled, relaxed circular, full-length linear supercoiled	18
Figure 2.12. There are breaking the phosphate	20
Figure 2.13. Point Mutation: this is a single change in one of the four nucleotide bases. An "A" might change to a "C" for example	21
Figure 2.14. Deletion: part of the DNA sequence is missing	22
Figure 2.15. Insertion: extra nucleotides are added to the sequence	22
Figure 2.16. Inversion: a portion of the DNA sequence is reversed	23
Figure 2.17. Schematic of spectrophotometers	24
Figure 2.18. Schematic views of gel electrophoresis, separate proteins and nucleic acids.	25
Figure 2.19. Schematic diagram of gel electrophoresis side view shown equipment for gel electrophoresis method.	26
Figure 2.20. The molecules will move through the matrix at different rates	26

Figure 2.21. Picture has shown mechanism of transformation. DNA fragment has taken into bacterial cell by electroporation, then it has 2 possible ways are stable transformation and unsuccessful transformation (a) and right side, Plasmid DNA has taken into bacterial cell (b).	27
Figure 2.22. Schematic diagram of electroporation	28
Figure 3.1. A photograph of the whole system (a) the standing column is the compact power supply, about 1.2 m high with wheels for easy move. (b) A photograph of the hand-held D-1 type jet, about 15 cm long. Now it was in the experimental fixture.	31
Figure 3.2. The CAPPJ apparatus used in the experiment. A photograph of the control panel's interface is at the top systems. It allows adjusting intensity and frequency of the plasma output.	32
Figure 3.3 Photograph of experimental set up consist of CAPPJ, spectrum receiver, DC- voltage power supply and plastic sample holder	32
Figure 3.4. Photograph (front view) of the plasma jet charge separation setup using a traversal electrical field.	33
Figure.3.5. Schematic of aluminum sample holder, which can load sample by inserting into between gaps of electrodes.	33
Figure 3.6. Photograph of probes (Tektrinox P 6015A) which reduced the voltage drop in the ratio of 1/1000.	34
Figure 3.7. Schematic circuits of the V-Q Lissajous figure method for measurement of the plasma power.	34
Figure.3.8. Picture of oscilloscope screen, x-axis represented applied voltage and y-axis represented charge.	35

Figure 3.9. A photograph of spectrum analyzer (a) and Input spectrum receiver set up near parallel aluminium plate (b)	36
Figure 3.10. Plasmid DNA prepared steps.	37
Figure 3.11. Photograph of BioDrop TOUCH, the gray circle is a direct micro-volume sample port	38
Figure.3.12. Dropping area for measuring naked DNA concentration is stand on middle of Bio-drop spectrophotometer, which is a small shallow pit.	38
Figure 3.13. Step of competent bacterial cell preparation.	39
Figure 3.14. Photograph of centrifuge mechanic, which is MCF-1350 LMS mini-centrifuge from LMS, Laboratory & Medical Supply, Inc.	40
Figure 3.15. Photograph of slides, they were marked treat condition by vary intensity level and frequency of plasma machine and dry (D) and wet (W) condition.	41
Figure 3.16. Photograph of electrophoresis system.	42
Figure 3.17. Photograph of a UV trans-illuminator	43
Figure 3.18. Electroporation systems in Left: Device and Right is Cuvette which use for fill sample.	43
Figure 4.1. Lissajous graph was measured by using condition is intensity level 1 and frequency 10 Hz	47
Figure 4.2. Lissajous graph was measured by using condition is intensity level 1 and frequency 50 Hz	47
Figure 4.3. Lissajous graph was measured by using condition is intensity level 1 and frequency 110 Hz	48
Figure 4.4. Lissajous graph was measured by using condition is intensity level 5 and frequency 10 Hz	48

Figure 4.5. Lissajous graph was measured by using condition is intensity level 5 and frequency 50 Hz	49
Figure 4.6. Lissajous graph was measured by using condition is intensity level 5 and frequency 110 Hz	49
Figure 4.7. Lissajous graph was measured by using condition is intensity level 10 and frequency 10 Hz	50
Figure 4.8. Lissajous graph was measured by using condition is intensity level 10 and frequency 50 Hz	50
Figure 4.9. Lissajous graph was measured by using condition is intensity level 10 and frequency 110 Hz	51
Figure 4.10. Graph of measured plasma generating energy, as a function of control parameter combinations of intensity in I number and frequency in Hz, i.e. I/Hz.	52
Figure 4.11. Graph of measured plasma generating power, as a function of control parameter combinations of intensity in I number and frequency in Hz, i.e. I/Hz.	53
Figure 4.12. Photograph of the plasma jet when it traverses a pair of parallel electrodes when the electrical field is (a) off and (b) on	54
Figure 4.13. Optical emission spectra (OES) of the CAPPJ, OES detect from the original plasma jet.	54
Figure 4.14. Optical emission spectra (OES) of the CAPPJ, OES detect from the area directly under the plasma jet exit, where the main part of the original plasma jet was bent away after it traversed the electrical field	55

Figure.4.15. Optical emission spectra (OES) of the CAPPJ, OES detect from the bent plasma jet after it traversed the electrical field	55
Figure.4.16. Photograph of gel electrophoresis result, it has 11 lanes.	57
Figure.4.17. Percentages of the DNA forms under varied CCAPPJ treatment conditions, quantified from the gel electrophoresis band light intensity	58
Figure.4.18. Electrophoresis result for comparison. M: Marker. A and E: I1/50Hz, dry. B and F: I1/50Hz, wet. C and G: I5/50, dry. D and H: I5/50Hz, wet. A-D: without electrical field. E-H: with electrical field. R: relaxed form. S: supercoiled form. L: linear form.	58
Figure.4.19. DNA form concentrations quantified from the electrophoresis result in Figure 4.18. I/Hz: plasma intensity level/frequency. Dry: dry sample. Wet: wet sample. O: original treatment without the electricla field. E: with the electrical field.	59
Figure.4.20. Picture of bacteria plate, which is a control. In petri - dish has shown a lot of green spot by using UV trans-illuminator.	61
Figure.4.21. Picture of bacterial petri dish, which is intensity 1 with frequency 50 Hz. In petri - dish has shown a small white spot follow the arrow.	61