

REFERENCES

- (1) Siegel RL, Miller KD, Jemal A. Cancer statistics, 2015. *CA: a cancer journal for clinicians*. 2015;65(1):5-29.
- (2) Imsamran W, Chaiwerawattana A, Wiangnon S, Pongnikorn D, Suwanrungruang K, Sangrajang S, *et al*. Cancer incidence in Thailand. *Cancer in Thailand*. 2015;8:59.
- (3) Hong Gee Sim, Cheng CWS. Changing demography of prostate cancer in Asia. *European Journal of Cancer*. 2005;44(6):834-54.
- (4) Surh YJ. Cancer chemoprevention with dietary phytochemicals. *Nature Reviews Cancer*. 2003;3(10):768-80.
- (5) Phannasorn W, Chewonarin T. Chemoprotective Effect of Purple Rice (*Oryza sativa* L. *indica*) Extract on the Promotion of Rat Colon Carcinogenesis. *Proceedings of International Graduate Research Conference*. 2014:64-8.
- (6) Naiki- Ito A, Chewonarin T, Tang M, Pitchakarn P, Kuno T, Ogawa K, *et al*. Ellagic acid, a component of pomegranate fruit juice, suppresses androgen dependent prostate carcinogenesis via induction of apoptosis. *The Prostate*. 2015;75(2):151-60.
- (7) Jang H, Ha US, Kim SJ, Yoon BI, Han DS, Yuk SM, *et al*. Anthocyanin extracted from black soybean reduces prostate weight and promotes apoptosis in the prostatic hyperplasia-induced rat model. *J Agric Food Chem*. 2010;58(24):12686-91.

- (8) Ha U, Bae WJ, Kim SJ, Yoon BI, Hong SH, Lee JY, *et al.* Anthocyanin Induces Apoptosis of DU-145 Cells In Vitro and Inhibits Xenograft Growth of Prostate Cancer. *Yonsei Med J.* 2015;56(1):16-23.
- (9) Wongjaikam S, Summart R, Chewonarin T. Apoptosis induction in colon cancer cell lines and alteration of aberrant crypt foci in rat colon by purple rice (*Oryza sativa* L. var. *glutinosa*) extracts. *Nutrition and cancer.* 2014;66(4):690-9.
- (10) Summart R, Chewonarin T. Purple rice extract supplemented diet reduces DMH-induced aberrant crypt foci in the rat colon by inhibition of bacterial β -glucuronidase. *Asian Pacific journal of cancer prevention: APJCP.* 2013;15(2):749-55.
- (11) Banjerdpongchai R, Wudtiwai B, Sringarm K. Cytotoxic and apoptotic-inducing effects of purple rice extracts and chemotherapeutic drugs on human cancer cell lines. *Asian Pac J Cancer Prev.* 2013;14(11):6541-8.
- (12) Bechis SK, Carroll PR, Cooperberg MR. Impact of Age at Diagnosis on Prostate Cancer Treatment and Survival. *J Clin Oncol.* 2011;29.
- (13) Pignot G, Salomon L, Lebacle C, Neuzillet Y, Lunardi P, Rischmann P, *et al.* Prostate cancer incidence on cystoprostatectomy specimens is directly linked to age: results from a multicentre study. *BJU Int.* 2015;115(1):87-93.
- (14) Chen R, Ren S, Yiu MK, Fai NC, Cheng WS, Ian LH, *et al.* Prostate cancer in Asia: a collaborative report. *Asian Journal of Urology.* 2014;1(1):15-29.
- (15) Baade PD, Youlden DR, Cramb SM, Dunn J, Gardiner RA. Epidemiology of prostate cancer in the Asia-Pacific region. *Prostate international.* 2013;1(2):47-58.

- (16) Pollard M, Luckert PH. Promotional effects of testosterone and high fat diet on the development of autochthonous prostate cancer in rats. *Cancer letters*. 1986;32(2):223-7.
- (17) Cho HJ, Kwon GT, Park H, Song H, Lee KW, Kim JI, et al. A high-fat diet containing lard accelerates prostate cancer progression and reduces survival rate in mice: possible contribution of adipose tissue-derived cytokines. *Nutrients*. 2015;7(4):2539-61.
- (18) Tan ME, Li J, Xu HE, Melcher K, Yong EL. Androgen receptor: structure, role in prostate cancer and drug discovery. *Acta Pharmacol Sin*. 2015;36(1):3-23.
- (19) Duskova M, Pospisilova H. The role of non-aromatizable testosterone metabolite in metabolic pathways. *Physiol Res*. 2011;60(2):253.
- (20) Carson C, Rittmaster R. The role of dihydrotestosterone in benign prostatic hyperplasia. *Urology*. 2003;61(4):2-7.
- (21) Suzuki H, Ueda T, Ichikawa T, Ito H. Androgen receptor involvement in the progression of prostate cancer. *Endocr Relat Cancer*. 2003;10(2):209-16.
- (22) Sharifi N, Gulley JL, Dahut WL. Androgen deprivation therapy for prostate cancer. *JAMA*. 2005;294(2):238-44.
- (23) Leitzmann MF, Stampfer MJ, Michaud DS, Augustsson K, Colditz GC, Willett WC, et al. Dietary intake of n-3 and n-6 fatty acids and the risk of prostate cancer. *The American journal of clinical nutrition*. 2004;80(1):204-16.
- (24) Sesso HD, Paffenbarger RS, Lee IM. Alcohol consumption and risk of prostate cancer: The Harvard Alumni Health Study. *Int J Epidemiol*. 2001;30(4):749-55.

- (25) Kenfield SA, Stampfer MJ, Giovannucci E, Chan JM. Physical activity and survival after prostate cancer diagnosis in the health professionals follow-up study. *J Clin Oncol*. 2011;29(6):726-32.
- (26) Wang LS, Stoner GD. Anthocyanins and their role in cancer prevention. *Cancer letters*. 2008;269(2):281-90.
- (27) Karna P, Chagani S, Gundala SR, Rida PC, Asif G, Sharma V, *et al*. Benefits of whole ginger extract in prostate cancer. *Br J Nutr*. 2012;107(04):473-84.
- (28) Agarwal C, Singh RP, Agarwal R. Grape seed extract induces apoptotic death of human prostate carcinoma DU145 cells via caspases activation accompanied by dissipation of mitochondrial membrane potential and cytochrome c release. *Carcinogenesis*. 2002;23(11):1869-76.
- (29) Johnson JJ, Bailey H, Mukhtar H. Green tea polyphenols for prostate cancer chemoprevention: a translational perspective. *Phytomedicine*. 2010;17(1):3-13.
- (30) Bemis DL, Capodice JL, Desai M, Katz AE, Buttyan R. β -Carboline alkaloid-enriched extract from the Amazonian rain forest tree *pao pereira* suppresses prostate cancer cells. *J Soc Integr Oncol*. 2009;7(2):59.
- (31) Shukla S, Gupta S. Dietary agents in the chemoprevention of prostate cancer. *Nutrition and cancer*. 2005;53(1):18-32.
- (32) Goufo P, Trindade H. Rice antioxidants: phenolic acids, flavonoids, anthocyanins, proanthocyanidins, tocopherols, tocotrienols, γ -oryzanol, and phytic acid. *Food science & nutrition*. 2014;2(2):75-104.
- (33) Oryza oil & fat chemical CO. Components of purple rice. Purple rice extract. 2012;4:3.

- (34) Lietti A, Cristoni A, Picci M. Studies on *Vaccinium myrtillus* anthocyanosides. I. Vasoprotective and antiinflammatory activity. *Arzneimittelforschung*. 1975;26(5):829-32.
- (35) Chen PN, Chu SC, Chiou HL, Kuo WH, Chiang CL, Hsieh YS. Mulberry anthocyanins, cyanidin 3-rutinoside and cyanidin 3-glucoside, exhibited an inhibitory effect on the migration and invasion of a human lung cancer cell line. *Cancer letters*. 2006;235(2):248-59.
- (36) Jang H, Bae WJ, Yuk SM, Han DS, Ha U, Hwang SY, *et al.* Seoritae extract reduces prostate weight and suppresses prostate cell proliferation in a rat model of benign prostate hyperplasia. *Evid Based Complement Alternat Med*. 2014;2014.
- (37) Zeng L, Gao J, Zhang R. [Study on anti-tumor effect of cyanidin-3-glucoside on ovarian cancer]. *Zhongguo Zhong yao za zhi= Zhongguo zhongyao zazhi= China journal of Chinese materia medica*. 2012;37(11):1651-4.
- (38) Cooke D, Schwarz M, Boocock D, Winterhalter P, Steward WP, Gescher AJ, *et al.* Effect of cyanidin- 3- glucoside and an anthocyanin mixture from bilberry on adenoma development in the ApcMin mouse model of intestinal carcinogenesis—Relationship with tissue anthocyanin levels. *Int J Cancer*. 2006;119(9):2213-20.
- (39) Klongpityapong P, Supabphol R, Supabphol A. Antioxidant effects of gamma-oryzanol on human prostate cancer cells. *Asian Pac J Cancer Prev*. 2013;14(9):5421-5.
- (40) Rungwattanapong T, Sringarm K, Karladee D. Total Phenolic Compound in Glutinous Purple Rice Grains and Relation to Phenotype Characteristics. *Agricultural Sci J*. 2011;42:417-20.
- (41) Jomduang S. Bioactive Compound Contents in Germinated Unpolished Purple Glutinous Rice from Kum Doi Saket and Kum Phayao Varieties.

CMUJ NS Special Issue on Food and Applied Bioscience.
2014;13(1):449-457.

- (42) Boonsit P, Pongpiachan P, Julsrigival S, Karladee D. Gamma oryzanol content in glutinous purple rice landrace varieties. CMUJ Nat Sci. 2010;9(1):151-7.
- (43) Karladee D, Boonsit P, Suriyong S, Sringarm K. Antioxidant capacities of vitamin E (tocopherols) in purple rice (*Oryza sativa* L. *indica*), perilla (*Perilla frutescens* L.) and sesame (*Sesamum indicum*). J Agri Technol. 2013;9(1):177-91.
- (44) Punyatong M, Pongpiachan P, Pongpiachan P, Karladee D, Mankhetkorn S. Cytotoxicity of crude proanthocyanidin extract from purple glutinous rice bran (*Oryza sativa* L.)(Kum Doi Saket) compared with cyanidin 3-glucoside on X63 myeloma cancer cell lines. Kasetsart J (Nat Sci). 2008;42:676-81.
- (45) Punvittayagul C, Sankam P, Taya S, Wongpoomchai R. Anticlastogenicity and Anticarcinogenicity of Purple Rice Extract in Rats. Nutrition and cancer. 2016;68(4):646-53.
- (46) Lee V, De Kretser D, Hudson B, Wang C. Variations in serum FSH, LH and testosterone levels in male rats from birth to sexual maturity. J Reprod Fertil. 1975;42(1):121-6.
- (47) Yawadio R, Tanimori S, Morita N. Identification of phenolic compounds isolated from pigmented rices and their aldose reductase inhibitory activities. Food Chem. 2007;101(4):1616-25.
- (48) Ghasemzadeh A, Jaafar HZ, Juraimi AS, Tayebi-Meigooni A. Comparative evaluation of different extraction techniques and solvents for the assay of phytochemicals and antioxidant activity of Hashemi Rice Bran. Molecules. 2015;20(6):10822-38.

- (49) Ling WH, Cheng QX, Ma J, Wang T. Red and black rice decrease atherosclerotic plaque formation and increase antioxidant status in rabbits. *The Journal of nutrition*. 2001;131(5):1421-6.
- (50) Xia M, Ling WH, Ma J, Kitts DD, Zawistowski J. Supplementation of diets with the black rice pigment fraction attenuates atherosclerotic plaque formation in apolipoprotein e deficient mice. *The Journal of nutrition*. 2003;133(3):744-51.
- (51) Rice-Evans C, Miller N, Paganga G. Antioxidant properties of phenolic compounds. *Trends in plant science*. 1997;2(4):152-9.
- (52) Hudson EA, Dinh PA, Kokubun T, Simmonds MS, Gescher A. Characterization of potentially chemopreventive phenols in extracts of brown rice that inhibit the growth of human breast and colon cancer cells. *Cancer Epidemiology Biomarkers & Prevention*. 2000;9(11):1163-70.
- (53) Chen PN, Chu SC, Chiou HL, Chiang CL, Yang SF, Hsieh YS. Cyanidin 3-glucoside and peonidin 3-glucoside inhibit tumor cell growth and induce apoptosis in vitro and suppress tumor growth in vivo. *Nutrition and cancer*. 2005;53(2):232-43.
- (54) Ding M, Feng R, Wang SY, Bowman L, Lu Y, Qian Y, et al. Cyanidin-3-glucoside, a natural product derived from blackberry, exhibits chemopreventive and chemotherapeutic activity. *J Biol Chem*. 2006;281(25):17359-68.
- (55) Chen PN, Kuo WH, Chiang CL, Chiou HL, Hsieh YS, Chu SC. Black rice anthocyanins inhibit cancer cells invasion via repressions of MMPs and u-PA expression. *Chem Biol Interact*. 2006;163(3):218-29.
- (56) Lowe FC, McConnell JD, Hudson PB, Romas NA, Boake R, Lieber M, et al. Long-term 6-year experience with finasteride in patients with benign prostatic hyperplasia. *Urology*. 2003;61(4):791-6.

- (57) Rolf C, Nieschlag E. Potential adverse effects of long-term testosterone therapy. *Baillieres Clin Endocrinol Metab.* 1998;12(3):521-34.
- (58) Soderfan AJ, Arlotto MP, Parkinson A. Identification of the Cytochrome P-450 Isozymes Responsible for Testosterone Oxidation in Rat Lung, Kidney, and Testis: Evidence that Cytochrome P-450a (P450IIA1) Is the Physiologically Important Testosterone 7 α -Hydroxylase in Rat Testis*. *Endocrinology.* 1989;125(2):857-66.
- (59) Lacroix E, Eechaute W, Leusen I. Influence of age on the formation of 5 α -androstenediol and 7 α -hydroxy-testosterone by incubated rat testes. *Steroids.* 1975;25(5):649-61.
- (60) Gao W, Kearbey JD, Nair VA, Chung K, Parlow AF, Miller DD, et al. Comparison of the pharmacological effects of a novel selective androgen receptor modulator, the 5 α -reductase inhibitor finasteride, and the antiandrogen hydroxyflutamide in intact rats: new approach for benign prostate hyperplasia. *Endocrinology.* 2004;145(12):5420-8.
- (61) Lee MY, Shin IS, Seo CS, *et al.* Effects of *Melandrium firmum* methanolic extract on testosterone-induced benign prostatic hyperplasia in Wistar rats. *Asian Journal of Andrology.* 2012;14(2):320-324.
- (62) Tisell LE and Angervall. The growth of the ventral prostate, the dorsolateral prostate, the coagulating glands and the seminal vesicles in castrated meal rats injected with ACTH and/or insulin. *Acta Endocrinologica.* 1969; 62: 694-710.
- (63) Yadav N, Heemers H. Androgen action in the prostate gland. *Minerva urologica e nefrologica= The Italian journal of urology and nephrology.* 2012;64(1):35-49.
- (64) Banerjee PP, Banerjee S, Brown TR. Increased androgen receptor expression correlates with development of age-dependent, lobe-specific

spontaneous hyperplasia of the brown Norway rat prostate.
Endocrinology. 2001;142(9):4066-75.

- (65) Choi HM, Jung Y, Park J, Kim HL, Youn DH, Kang J, *et al.* Cinnamomi Cortex (*Cinnamomum verum*) Suppresses Testosterone-induced Benign Prostatic Hyperplasia by Regulating 5 α -reductase. Sci Rep. 2016;6.
- (66) Song L, Shen W, Zhang H, Wang Q, Wang Y, Zhou Z. Differential expression of androgen, estrogen, and progesterone receptors in benign prostatic hyperplasia. Bosnian Journal of Basic Medical Sciences. 2016;16(3):201.
- (67) Brolin J, Ekman P, Skoog L. Immunohistochemistry and biochemistry in detection of androgen, progesterone, and estrogen receptors in benign and malignant human prostatic tissue. The Prostate. 1992;20(4):281-95.
- (68) Kim YN, Kim MS, Chun SS, Choi JH. Effect of Phellius linteus water extract on benign prostatic hyperplasia. Nutrition research and practice. 2013;7(3):172-7.
- (69) Long N, Suzuki S, Sato S, Naiki-Ito A, Sakatani K, Shirai T, *et al.* Purple corn color inhibition of prostate carcinogenesis by targeting cell growth pathways. Cancer Sci. 2013;104(3):298-303.
- (70) Balk SP, Ko YJ, Bubley GJ. Biology of prostate-specific antigen. J Clin Oncol. 2003;21(2):383-91.
- (71) Montgomery BT, Young CYF, Bilhartz DL, Andrews PE, Thompson NF, Tindall DJ, *et al.* Hormonal regulation of prostate- specific antigen (PSA) glycoprotein in the human prostatic adenocarcinoma cell line, LNCaP. The Prostate. 1992;21(1):63-73.
- (72) Lee C, Sutkowski DM, Sensibar JA, Zelner D, Kim I, Amsel I, *et al.* Regulation of proliferation and production of prostate-specific antigen in androgen-sensitive prostatic cancer cells, LNCaP, by dihydrotestosterone. Endocrinology. 1995;136(2):796-803.

- (73) Lee MS, Lee SO, Kim SH, Lee EO, Lee HJ. Anti-Cancer Effect of Lambertianic Acid by Inhibiting the AR in LNCaP Cells. *International Journal of Molecular Sciences*. 2016;17(7):1066.
- (74) Das K, Lorena PD, Ng LK, Lim D, Shen L, Siow WY, *et al*. Differential expression of steroid 5 α -reductase isozymes and association with disease severity and angiogenic genes predict their biological role in prostate cancer. *Endocr Relat Cancer*. 2010;17(3):757-70.
- (75) Kumar T, Chaiyavat C, Rungseevijitprapa W, Suttajit M. Screening of steroid 5 α -reductase inhibitory activity and total phenolic content of Thai plants. *Journal of Medicinal Plants Research*. 2011; 5(7): 1265-1271.