

REFERENCES

- [1] World Health Organization. Obesity and overweight. 2006. [Cited on 14 October 2013]. Available from: <http://www.who.int/mediacentre/factsheets/fs311/en/index.html>.
- [2] Disease Control Priorities Project (DCPP). Risk factors: Millions of deaths could be prevented by reducing risk factor exposure. 2006.
- [3] Pruenglampoo S, Pruenglampoo S, Kingkeow C, Mangklabruks A. Simple index for screening overweight and obesity. *Public Health Nutr.* 2003;6:225-226.
- [4] Gibson R. Principles of nutrition assessment. 2nd ed. Oxford: Oxford University Press. 2005.
- [5] Benn R. Some mathematical properties of weight-for-height indices used as measures of adiposity. *Br J Prev Soc Med.* 1971;25:42-50.
- [6] Pruenglampoo S, Pruenglampoo S, Kingkeow C, Mangklabruks A. A new simple healthy index for nutritional status assessment. Department of Internal Medicine, Faculty of Medicine, Chiang Mai University, Thailand, December. 2005;49.
- [7] Chittawatanarat K, Pruenglampoo S, Kongsawasdi S, et al. The variations of body mass index and body fat in adult Thai people across the age spectrum measured by bioelectrical impedance analysis. *Clin Interv Aging.* 2011;6:285-294.
- [8] World Health Organization. Obesity: preventing and managing the global epidemic: report of a WHO consultation. Geneva:World Health Organization. 2000.

- [9] Chittawatanarat K. Anthropometric measurement in clinical applications: Height, body weight and body composition prediction. PhD Thesis, Doctor of philosophy in clinical epidemiology, Chiangmai University, Thailand. 2012.
- [10] Mandal A. What is a Biomarker? 2014. [Cited on 9 July 2015]. Available from: <http://www.news-medical.net/health/What-is-a-Biomarker.aspx>.
- [11] Robert CW. Body mass index as an indicator of obesity. Asia Pacific J Clin Nutr. 2002;11:S681–S684.
- [12] Worldobesity. Developmental origin of fat. [Cited on 9 July 2015] Available from: <http://www.worldobesity.org/aboutobesity>.
- [13] Nguyen T, Kamsrichan W, Chompikul J. Obesity and related factors among students grade 7-12 in Phuttha Monthon district, Nakhon Pathom province, Thailand. A thesis for the degree of master of primary health care management faculty of graduate studies, Mahidol University. 2008.
- [14] Kumcheen P. Thailand under obesity crisis. 2009. [Cited on 9 July 2015] Available from: http://www.siam.edu/siamedu_thai_mainpage/images/stories/article/54/siam_article_7_54.pdf.
- [15] Somnuek P. Childhood Obesity: A Weighty Problem. 2013. [Cited on 9 July 2015] Available from: <http://www.nationmultimedia.com/life/Childhood-Obesity-A-Weighty-Problem-30198882.html>.
- [16] Porapakkham Y, Pattaraarchachai J, Aekplakorn W. Prevalence, awareness, treatment and control of hypertension and diabetes mellitus among the elderly: the 2004 National Health Examination Survey III, Thailand. Singapore Med J. 2008;49:868-873.

- [17] Aekplakorn W, Hogan MC, Chongsuvivatwong V, et al. Trends in obesity and associations with education and urban or rural residence in Thailand. *Obesity*. 2007;15:3113–3121.
- [18] World Health Organization/International Association for the Study of Obesity/International Obesity Task Force. *The Asia Pacific Perspective: Redefining Obesity and Its Treatment*. Health Communications Australia Pty Ltd.: Melbourne, 2000.
- [19] World Health Organization. *Obesity: Preventing and managing the Global Epidemic: WHO Technical Report series*. World Health Organization: Geneva. 2000;894.
- [20] Gallagher D, Heymsfield SB, Heo M, Jebb SA, Murgatroyd PR, Sakamoto Y. Healthy percentage body fat ranges: an approach for developing guidelines based on body mass index. *Am J Clin Nutr*. 2000;72:694-701.
- [21] American Council on Exercise. *What are the guidelines for percentage of body fat loss? Ask the Expert Blog*. 2009.
- [22] Chumlea WC, Guo SS. Bioelectrical impedance and body composition: present status and future directions. *Nutr Rev*. 1994;52:123-131.
- [23] Gudivaka R, Schoeller DA, Kushner RF, Bolt MJ. Single- and multifrequency models for bioelectrical impedance analysis of body water compartments. *J Appl Physiol*. 1999;87:1087-1096.
- [24] Cornish BH, Thomas BJ, Ward LC. Improved prediction of extracellular and total body water using impedance loci generated by multiple frequency bioelectrical impedance analysis. *Phys Med Biol*. 1993;38:337-346.
- [25] Van Loan MD, Mayclin PL. Use of multi-frequency bioelectrical impedance analysis for the estimation of extracellular fluid. *Eur J Clin Nutr*. 1992;46:117-124.

- [26] Brodie D, Moscrip V, Hutcheon R. Body composition measurement: a review of hydrodensitometry, anthropometry, and impedance methods. *Nutrition*. 1998;14:296-310.
- [27] Elia M, Ward LC. New techniques in nutritional assessment: body composition methods. *Proc Nutr Soc*. 1999;58:33-38.
- [28] Tothill P, Hannan WJ, Wilkinson S. Comparisons between a pencil beam and two fan beam dual energy X-ray absorptiometers used for measuring total body bone and soft tissue. *Br J Radiol*. 2001;74:166-176.
- [29] Sarria A, García-Llop A, Moreno LA, Fleta J, Morellón MP, Bueno M. Skinfold thickness measurements are better predictors of body fat percentage than body mass index in male spanish children and adolescents. *Eur J Clin Nutr*. 1998;52:573-576.
- [30] Bruner R. Fitness and nutrition. *The Jerusalem Post*. 2001.
- [31] Wagner DR, Heyward VH. Measures of body composition in blacks and whites: a comparative review. *Am J Clin Nutr*. 2000;71:1392-1402.
- [32] Slaughter MH, Lohman TG, Boileau RA, et al. Skinfold equations for estimation of body fatness in children and youth. *Hum Biol*. 1988;60:709-723.
- [33] Robert JV, Womersley J. Body fat assessed from total body density and its estimation from skinfold thickness: measurements on 481 men and women aged from 16 to 72 years. *Br J Nutr*. 1974;32:77-97.
- [34] Willett DC, Watts GF, Barrett PH, Burke V. Waist circumference, waist-to-hip ratio and body mass index as predictors of adipose tissue compartments in men. *Q Med*. 2003;96:441-447.
- [35] Willett WC, Stampfer M, Manson J, VanItallie T. New weight guidelines for Americans: justified or injudicious. *Am J Clin Nutr*. 1991;53:1102-1103.

- [36] Manson JE, Colditz GA, Stampfer MJ, et al. A prospective study of obesity and risk of coronary heart disease in women. *N Engl J Med*. 1990;322:882-889.
- [37] Schapira DV, Clark RA, Wolff PA, Jarrett AR, Kumar NB, Aziz NM. Visceral obesity and breast cancer risk. *Cancer*. 1994;74:632-639.
- [38] Wolk A, Lindblad P, Adami HO. Nutrition and renal cell cancer. *Cancer Cause Control*. 1996;7:5-18.
- [39] Van Noord PA, Seidell JC, den Tonkelaar I, Baanders-van Halewijn EA, Ouwehand IJ. The relationship between fat distribution and some chronic disease in 11,825 women participating in the DOM-project. *Int J Epidemiol*. 1990;19:546-570.
- [40] Chan JM, Rimm EB, Colditz GA, Stampfer MJ, Willett WC. Obesity, fat distribution, and weight gain as risk factors for clinical diabetes in men. *Diabetes Care*. 1994;17:961-969.
- [41] Chou P, Liao MJ, Tsai ST. Associated risk factors for diabetes in Kin-HU, Kinmen. *Diabetes Res Clin Pract*. 1994;26:229-235.
- [42] Helmrich SP, Ragland DR, Leung RW, Paffenbarger RS. Physical activities and reduced occurrence of noninsulin dependent diabetes mellitus. *N Engl J Med*. 1991;325:147-152.
- [43] Center for Diseases Control and prevention. Trends in intake of energy and macronutrients – United States, 1971-2000. *Morbidity and Mortality Weekly Report*. 2004;53:80-82.
- [44] World Health Organization. The challenge of obesity in the WHO European Region and the strategies for response. Denmark: WHO Regional Office for Europe. 2007.

- [45] Dastmard B. A statistical analysis of the connection between test results and field claims for ECUs in vehicles. Master of Science Thesis Stockholm, Sweden. 2013.
- [46] Correlation. [Cited on 4 October 2015]. Available from: <http://www.sjsu.edu/faculty/gerstman/StatPrimer/correlation.pdf>.
- [47] Lengerich R. Cohen's Kappa Statistic for Measuring Agreement. Department of Public Health Sciences. Penn State College of Medicine. 2009.
- [48] Landis J, Koch GG. The measurement of observer agreement for categorical data. *Biometrics*. 1977;33:159-174.
- [49] Karimollah HT. Receiver Operating Characteristic (ROC) curve analysis for medical diagnostic test evaluation. *Caspian J Intern Med*. 2013;4:627-635.
- [50] Akobeng AK. Understanding diagnostic test 3: Receiver operating characteristic curves. *Acta Paediatr* 2007;90:644-647.
- [51] Perkins NJ, Schisterman EF. The inconsistency of optimal cut-points obtained using two criteria based on receiver operating characteristic curve. *Am J Epidemiol*. 2006;163:670-675.
- [52] Fluss R, Faraggi D, Reiser B. Estimation of Youden index and its associated cutoff point. *Biom J*. 2005;47:458-472.
- [53] Crichton N. Information point: receiver operating characteristic (ROC) curve. *J Clin Nurs*. 2002;11:136.
- [54] Christensen E. Methodology of diagnostic tests in hepatology. *Ann Hepatol*. 2009;8:177-183.
- [55] Obuchowski NA. Receiver operating characteristic curves and their use in radiology. *Radiology*. 2003;229:3-8.

- [56] Michils A, Louis R, Peché R, Baldassarre S, Van Muylem A. Exhaled nitric oxide as a marker of asthma control in smoking patients. *Eur Respir J*. 2009;33:1295-1301.
- [57] Campbell D, Campbell S. Introduction to Regression and Data Analysis, StatLab Workshop Series. 2008.
- [58] Srivastava R. Polynomial regression. [Cited on 20 July 2015] Available from: <http://www.iasri.res.in/iasriwebsite/DESIGNOFEXPAPPLICATION/Electronic-Book/module5/11PRegres.pdf>.
- [59] Ranasinghe C, Gamage P, Katulanda P, Andraweera N, Thilakarathne S, Tharanga P. Relationship between Body mass index (BMI) and body fat percentage, estimated by bioelectrical impedance, in a group of Sri Lankan adults: a cross sectional study. *BMC Public Health*. 2013;13:797-804.
- [60] Nirav R, Eric R. Measuring adiposity in patients: The utility of body mass index (BMI), Percent body fat, and leptin. *PLoS One*. 2012;7:e33308.
- [61] Bedogni G, Agosti F, De Col A, Marazzi N, Tagliaferri A, Sartorio A. Comparison of dual-energy X-ray absorptiometry, air displacement plethysmography and bioelectrical impedance analysis for the assessment of body composition in morbidly obese women. *Eur J Clin Nutr*. 2013;67:1129-1132.
- [62] Flegal K, Shepherd JA, Looker AC, et al. Comparisons of percentage body fat, body mass index, waist circumference, and waist-stature ratio in adults. *Am J Clin Nutr*. 2009;89:500-508.
- [63] Lazarus R, Baur L, Webb K, Blyth F. Body mass index in screening for adiposity in children and adolescents: systematic evaluation using receiver operating characteristic curves. *Am J Clin Nutr*. 1996;63:500-506.

- [64] Kesavachandran CN, Bihari V, Mathur N. The normal range of body mass index with high body fat percentage among male residents of Lucknow city in north India. *Indian J Med Res.* 2012;135:72-77.
- [65] Meeuwse S, Horgan GW, Elia M. The relationship between BMI and percent body fat, measured by bioelectrical impedance, in a large adult sample is curvilinear and influenced by age and sex. *Clin Nutr.* 2010;29:560-566.
- [66] Macias N, Quezada AD, Flores M, et al. Accuracy of body fat percent and adiposity indicators cut off values to detect metabolic risk factors in a sample of Mexican adults. *BMC public health.* 2014;14:341-349.
- [67] Ho-Pham LT, Lai TQ, Nguyen MT, Nguyen TV. Relationship between Body Mass Index and Percent Body Fat in Vietnamese: Implications for the Diagnosis of Obesity. *PloS one.* 2015;10:e0127198.
- [68] World Health Organization. The WHO Global Database on Body Mass Index. 2009. [Cited on 21 August 2015] Available from: http://apps.who.int/bmi/index.jsp?intro Page=intro_3.html.
- [69] Ruopp MD, Perkins NJ, Whitcomb BW, Schisterman EF. Youden Index and optimal cut-point estimated from observations affected by a lower limit of detection. *Biom J.* 2008;50:419-430.
- [70] De Lorenzo A, Nardi A, Iacopino L, et al. A new predictive equation for evaluating women body fat percentage and obesity-related cardiovascular disease risk. *J Endocrinological Invest.* 2014;37:511-524.
- [71] Fatima SS, Rehman R, Chaudhry B. Body mass index or body fat! Which is a better obesity scale for Pakistani population? *J Pak Med Assoc.* 2014;64:1225-1228.

- [72] Molnar D, Schutz Y. The effect of obesity, age, puberty and gender on resting metabolic rate in children and adolescents. *Eur J Pediatr*. 1997;156:376-381.
- [73] Shah NR, Braverman ER. Measuring Adiposity in Patients: The Utility of Body Mass Index (BMI), Percent Body Fat, and Leptin. *PLoS ONE*. 2012;7:e33308.
- [74] Zeng Q, Dong SY, Sun XN, Xie J, Cui Y. Percent body fat is a better predictor of cardiovascular risk factors than body mass index. *Braz J Med Biol Res*. 2012;45:591-600.
- [75] Karandish M. Validation study of Height-Weight Difference Index (HWDI) for screening underweight and obesity status in Tehranian adults. *Research in Medicine*. 2006;30:91-95.
- [76] Luke A, Durazo-Arvizu R, Rotimi C, et al. Relation between body mass index and body fat in black population samples from Nigeria, Jamaica, and the United States. *Am J Epidemiol*. 1997;145:620-628.
- [77] Ranasinghe C, Gamage P, Katulanda P, Andraweera N, Thilakarathne S, Tharanga P. Relationship between Body mass index (BMI) and body fat percentage, estimated by bioelectrical impedance, in a group of Sri Lankan adults: a cross sectional study. *BMC Public Health*. 2013;13:797-804.
- [78] Romero-Corral A, Somers VK, Sierra-Johnson J, et al. Accuracy of Body Mass Index to Diagnose Obesity In the US Adult Population. *Int J Obes (Lond)*. 2008;32:959-66.
- [79] David E. BMI: The Good, The Bad, and The Ugly. 2015.
- [80] Kerry S. Healthy weight, healthy lives: a toolkit for developing local strategies. 2008.
- [81] Habib SS. Body mass index and body fat percentage in assessment of obesity prevalence in saudi adults. *Biomed Environ Sci*. 2013;26:94-99.

- [82] Wang C, Hou X, Zhang M, et al. Comparison of body mass index with body fat percentage in the evaluation of obesity in Chinese. *Biomed Environ Sci*. 2010;23:173-179.
- [83] Nur A, Huddin B, Arma M. Comparison of body mass index and body fat percentage in the assessment of obesity prevalence among secondary school students in Perak state, Malaysia. *European Science Journal*. 2014;10:115-122.
- [84] Goonasegaran AR, Mat N, Fatin N, Abdul W, Nurul S. Comparison of the effectiveness of body mass index and body fat percentage in defining body composition. *Singapore Med J*. 2012;53:403-408.
- [85] Gallagher D, Visser M, Sepúlveda D, Pierson RN, Harris T, Heymsfield SB. How useful is body mass index for comparison of body fatness across age, sex, and ethnic groups. *Am J Epidemiol*. 1996;143:228-239.
- [86] Sampei MA, Novo NF, Juliano Y, Sigulem DM. Comparison of the body mass index to other methods of body fat evaluation in ethnic Japanese and Caucasian adolescent girls. *Int J Obes Relat Metab Disord*. 2001;25:400-408.
- [87] Senior PA, Bhopal R. Ethnicity as a variable in epidemiological research. *BMJ*. 1994;309:327-330.
- [88] World Health Organization (WHO). Obesity and overweight. 2015. [Cited on 11 April 2016] Available from: <http://www.who.int/mediacentre/factsheets/fs311/en>.
- [89] Shukla A, Kumar K, Singh A. Association between Obesity and Selected Morbidities: A Study of BRICS Countries. *PLoS ONE*. 2014;9:e94433.
- [90] Akmal N, Huddin B, Arma M. Comparison of Body Mass Index and Body Fat Percentage in the Assessment of Obesity Prevalence among Secondary

School Students in Perak State, Malaysia. *European Scientific Journal*. 2014;10:115-122.

- [91] Frank LD, Andresen MA, Schmid TL. Obesity relationships with community design, physical activity, and time spent in cars. *Am J Prev Med*. 2004;27:87-96.
- [92] Hajian-Tilaki KO, Heidari B. Prevalence of obesity, central obesity and the associated factors in urban population aged 20–70 years, in the north of Iran: a population-based study and regression approach. *Obes Rev*. 2007;8:3-10.
- [93] Jia H, Li JZ, Leserman J, Hu Y, Drossman DA. Relationship of abuse history and other risk factors with obesity among female gastrointestinal patients. *Dig Dis Sci*. 2004;49:872-877.
- [94] Haines J, Sztainer DN. Prevention of obesity and eating disorders: a consideration of shared risk factors. *Health Education Research*. 2006;21:770-782.
- [95] Ilman M, Zuhairini Y, Siddiq A. Correlation between Body Mass Index and Body Fat Percentage. *Althea Medical Journal*. 2016;2:575-578.
- [96] Wang J, Thornton JC, Russell M, Burastero S, Heymsfield S, Pierson RN Jr. Asians have lower body mass index (BMI) but higher percent body fat than do whites: comparisons of anthropometric measurements. *Am J clin Nutr*. 1994;60:23-28.
- [97] Kupper J, Bartz M, Schultink JW, Lukito W, Deurenberg P. Measurements of body fat in Indonesian adults: Comparison between a three-compartment model and widely used methods. *Asia Pac J Clin Nutr*. 1998;7:49-54.
- [98] Gurrici S, Hartriyanti Y, Hautvast JG, Deurenberg P. Differences in the relationship between body fat and body mass index between two different

Indonesian ethnic groups: the effect of body build. *Eur J Clin Nutr.* 1999;53:468-472.

- [99] Deurenberg P, Kooy K, Hulshof T, Evers P. Body mass index as a measure of body fatness in the elderly. *Eur J Clin Nutr.* 1989;43:231-236.
- [100] Jackson AS, Stanforth PR, Gagnon J, et al. The effect of sex, age and race on estimating percentage body fat from body mass index: The Heritage Family Study. *Int J Obes Relat Metab Disord.* 2002;26:789-796.
- [101] Rush EC, Freitas I, Plank LD. Body size, body composition and fat distribution: comparative analysis of European, Maori, Pacific Island and Asian Indian adults. *Br J Nutr.* 2009;102:632-641.
- [102] Jackson AS, Pollock ML, Ward A. Generalized equations for predicting body density of women. *Medicine and science in sports and exercise.* 1980;12:175-181.
- [103] Jacoby WG. Regression iii: Advanced methods. Retrieved October. 2005;18:2011.
- [104] Nathans LL, Oswald FL, Nimon K. Interpreting multiple linear regression: A guidebook of variable importance. *Practical Assessment, Research & Evaluation.* 2012;17:1-18.
- [105] Schneider A, Hommel G, Blettner M. Linear Regression Analysis: Part 14 of a Series on Evaluation of Scientific Publications. *Dtsch Arztebl Int.* 2010;107:776-82.