

REFERENCES

- [1] R.K. Singh, U.S. Singh, G.S. Khush, Aromatic Rices, Oxford & IBH, New Delhi, India, 2000.
- [2] R.J. Bryant, A.M. McClung, Volatile profiles of aromatic and non-aromatic rice cultivars using SPME/GC–MS, Food Chem. 124 (2011) 501–513.
- [3] V. Laksanalama, S. Ilangantileke, Comparison of aroma compound (2-acetyl-1-pyrroline) in leaves from pandan (*Pandanus amaryllifolius*) and Thai fragrant rice (Khao Dawk Mali-105), Cereal Chem. 70 (4) (1993) 381–384.
- [4] E.T. Champagne, Rice aroma and flavor: a literature review, Cereal Chem. 85 (4) (2008) 445–454.
- [5] T. Yoshihashi, N.T.T. Huong, H. Inatomi, Precursors of 2-acetyl-1-pyrroline, a potent flavor compound of an aromatic rice variety, J. Agric. Food Chem. 50 (2002), 2001–2004.
- [6] K. Sakthivel, R.M. Sundaram, N. Shobha Rani, S.M. Balachandran, C.N. Neeraja, Genetic and molecular basis of fragrance in rice, Biotechnol. Adv. 27 (2009) 468–473.
- [7] T.L. Fitzgerald, D.L.E. Waters, L.O. Brooks, R.J. Henry, Fragrance in rice (*Oryza sativa*) is associated with reduced yield under salt treatment, Environ. Exp. Bot. 68 (2010), 292–300.
- [8] F. Gay, I. Maraval, S. Roques, Z. Gunata, R. Boulanger, A. Audelbert, C. Mestres, Effect of salinity on yield and 2-acetyl-1-pyrroline content in the grains of three fragrant rice cultivars (*Oryza sativa* L.) in Camargue (France), Field Crop Res. 117 (2010), 154–160.

- [9] J. Poonlaphdecha, I. Maraval, S. Roques, A. Audebert, R. Boulanger, X. Bry, Z. Gunata, Effect of timing and duration of salt treatment during growth of a fragrant rice variety on yield and 2-acetyl-1-pyrroline, proline, and GABA levels, *J. Agric. Food Chem.* 60 (2012) 3824–3830.
- [10] T.L. Fitzgerald, D.L.E. Waters, R.J. Henry, The effect of salt on betaine aldehyde dehydrogenase transcript levels and 2-acetyl-1-pyrroline concentration in fragrant and non-fragrant rice (*Oryza sativa*), *Plant Sci.* 175 (2008) 539–546.
- [11] S. Yang, Y. Zou, Y. Liang, B. Xia, S. Liu, M. Ibrahim, D. Li, Y. Li, L. Chen, Y. Zeng, L. Liu, Y. Chen, P. Li, J. Zhu, Role of soil total nitrogen in aroma synthesis of traditional regional aromatic rice in China, *Field Crop Res.* 125 (2010) 151–160.
- [12] R.G. Buttery, L.C. Ling, B.O. Juliano, 2-Acetyl-1-pyrroline: an important aroma component of cooked rice, *Chem. Ind.* 23 (1982) 958–959.
- [13] S. Chen, Y. Yang, W. Shi, Q. Ji, F. He, Z. Zhang, Badh2, encoding betaine aldehyde dehydrogenase, inhibits the biosynthesis of 2-acetyl-1-pyrroline, a major component in rice fragrance. *Plant Cell* 20 (2008) 1850-1861.
- [14] L.M.T. Bradbury, S.A. Gillies, D.J. Brushett, D.L.E. Waters, R.J. Henry, Inactivation of an aminoaldehyde dehydrogenase is responsible for fragrance in rice. *Plant Mol. Biol.* 68 (2008) 439-449.
- [15] T.C. Huang, C.S. Teng, J.L. Chang, H.S. Chuang, C.T. Ho, M.L. Wu, Biosynthetic mechanism of 2-acetyl-1-pyrroline and its relationship with Δ 1-pyrroline-5-carboxylic acid and methylglyoxal in aromatic rice (*Oryza sativa* L.) callus. *J. Agric. Food Chem.* 56 (2008) 7399-7404.
- [16] K. Sangwijit, P. Thangsunan, R. Cutler, S. Anuntalabchchai, Development of SCAR marker for Thai fragrant rice (*Oryza sativa* L. var.indica cv. Pathum Thani 1) mutants induced by low energy ion beam, *Chiang Mai J. Sci.* 39 (4) (2012) 545–553.

- [17] P.C. Srivastava, U.S. Singh, Effect of graded levels of nitrogen and sulfur and their interaction on yields and quality of aromatic rice, *J. Plant Nutr.* 30 (2007) 811–828.
- [18] R.G. Brereton, *Chemometrics: Data Analysis for the Laboratory and Chemical Plant*, Wiley, Chichester, U.K., 2003
- [19] R.G. Brereton, *Chemometrics for Pattern Recognition*, Wiley, Chichester, U.K., 2009
- [20] S. Kittiwachana, D.L.S. Ferreira, L.A. Fido, D.R. Thompson, R.E.A. Escott, R.G. Brereton, Self-Organizing Map Quality Control Index. *Anal. Chem.* 82 (2010) 5972-5982.
- [21] G.R. Lloyd, S. Ahmad, M. Wasim, R.G. Brereton, Pattern Recognition of Inductively Coupled Plasma Atomic Emission Spectroscopy of Human Scalp Hair for Discriminating between Healthy and Hepatitis C Patients. *Anal. Chim. Acta.* 649 (2009) 33-42.
- [22] P. Geladi, B.R. Kowalski, Partial Least-Squares Regression - a Tutorial. *Anal. Chim. Acta.* 185 (1986) 1-17.
- [23] G.R. Lloyd, K. Wongravee, C.J.L. Silwood, M. Grootveld, R.G. Brereton, Self Organising Maps for Variable Selection: Application to Human Saliva Analysed by Nuclear Magnetic Resonance Spectroscopy to Investigate the Effect of an Oral Healthcare Product. *Chemometr Intell Lab.* 98 (2009) 149-161.
- [24] T.J. Thurston, R.G. Brereton, D.J. Foord, R.E.A. Escott, Principal Components Plots for Exploratory Investigation of Reactions Using Ultraviolet-Visible Spectroscopy: Application to the Formation of Benzophenone Phenylhydrazone. *Talanta* 63 (2004) 757-769.
- [25] B. Levine, J. Workman, *Chemometrics. Anal. Chem.* 78 (2006) 4137-4145.

- [26] M.J. Nueda, A. Conesa, J.A. Westerhuis, H.C.J. Hoefsloot, A.K. Smilde, M. Talon, A. Ferrer, Discovering Gene Expression Patterns in Time Course Microarray Experiments by Anova-Sca. *Bioinformatics* 23 (2007) 1792-1800.
- [27] R.J. Wenning, G.A. Erickson, Interpretation and Analysis of Complex Environmental Data Using Chemometric Methods. *TrAC-Trend. Anal. Chem.* 13 (1994) 446-457.
- [28] J.E. Meng, W. Shiqian, L. Juwei, T. Hock Lye, Face Recognition with Radial Basis Function (Rbf) Neural Networks. *IEEE Trans. Neural Netw. Learn. Syst.* 13 (2002) 697-710.
- [29] T. Lundstedt, E. Seifert, L. Abramo, B. Thelin, Å. Nyström, J. Pettersen, R. Bergman, Experimental design and optimization, *Chemom. Intell. Lab. Syst.* 42 (1998) 3–40.
- [30] D.R. Hoagland, D.I. Arnon, The Water-Culture Method for Growing Plants without Soil, University of California, College of Agriculture, Agricultural Experiment Station, Berkeley, Calif., 1950
- [31] T. Sriseadka, S. Wongpornchai, P. Kitsawatpaiboon, Rapid method for quantitative analysis of the aroma impact compound, 2-acetyl-1-pyrroline, in fragrant rice using automated headspace gas chromatography, *J. Agric. Food Chem.* 54 (2006) 8183–8189.
- [32] J.A. Westerhuis, T. Kourti, J.F. Macgregor, Analysis of multiblock and hierarchical PCA and PLS models, *J. Chemom.* 12 (1998) 301–321.
- [33] A. Berglund, S. Wold, A serial extension of multiblock PLS, *J. Chemom.* 13 (1999) 461–471.
- [34] J.A. Westerhuis, A.K. Smilde, Deflation in multiblock PLS, *J. Chemom.* 15 (2001) 485–493.

- [35] K. Jörgensen, V. Segtnan, K. Thyholt, N. Tormod, A comparison of methods for analysing regression models with both spectral and designed variables, *J. Chemom.* 18 (2004) 451–464.
- [36] P. Yaroshchyk, D.L. Death, S.J. Spencer, Comparison of principal components regression, partial least squares regression, multi-block partial least squares regression, and serial partial least squares regression algorithms for the analysis of Fe in iron ore using LIBS, *J. Anal. At. Spectrom.* 27 (2012) 92–98.
- [37] M. Farrésa, S. Platikanova, S. Tsakovskib, R. Tauler, Comparison of the variable importance in projection (VIP) and of the selectivity ratio (SR) methods for variable selection and interpretation, *J. Chemom.* 29 (2015) 528–536.
- [38] R.G. Brereton, *Applied Chemometrics for Scientists*, Wiley, Chichester, U.K., 2007
- [39] W. Melssen, R. Wehrens, L. Buydens, Supervised Kohonen networks for classification problems, *Chemom. Intell. Lab. Syst.* 83 (2006) 99–113.
- [40] G.R. Lloyd, R.G. Brereton, J.C. Duncan, Self organising maps for distinguishing polymer groups using thermal response curves obtained by dynamic mechanical analysis, *Analyst* 133 (8) (2008) 1046–1059.
- [41] L. Eriksson, E. Johansson, C. Wikstrom, Tutorial mixture design—design generation, PLS analysis and model usage, *Chemom. Intell. Lab. Syst.* 43 (1998) 1–24.
- [42] K. Kumar, M. Kumar, S. Kim, H. Ryu, Y. Cho, Insights into genomics of salt stress response in rice, *Rice* 6 (2013) 27.
- [43] S. Chutipaijit, S. Cha-um, K. Sompornpailin, High contents of proline and anthocyanin increase protective response to salinity in *Oryza sativa* L. ssp. indica, *Aust. J. Crop. Sci.* 5 (10) (2011) 1191–1198.
- [44] A.K. Parida, A.B. Das, Salt tolerance and salinity effects on plants: a review, *Ecotoxicol. Environ. Saf.* 60 (2005) 324–349.

- [45] Z. Zhang, Q. Liu, H. Song, X. Rong, M.I. Abdelbagi, Responses of different rice (*Oryza sativa* L.) genotypes to salts stress and relation to carbohydrate metabolism and chlorophyll content, Afr. J. Agric. Res. 7 (1) (2012) 19–27.




ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่

Copyright© by Chiang Mai University

All rights reserved

LIST OF PUBLICATIONS

- 1) S. Funsueb, C. Krongchai, S. Mahatheeranont, S. Kittiwachana, "Prediction of 2-Acetyl-1-Pyrroline Content in Grains of Thai Jasmine Rice Based on Planting Condition, Plant Growth and Yield Component Data Using Chemometrics", *Chemom. Intell. Lab. Syst.*, 156 (2016), 203-210.
- 2) S. Funsueb, G. Sudtasarn, S. Kittiwachana, "Determination of Amylose Contents in Rice Grains Using Near Infrared Reflectance Spectroscopy (NIRS) and Supervised Self-Organizing Map (SOM)", *Proceedings of the Pure and Applied Chemistry International Conference 2016 (PACCON 2016)*, BITEC, Bangkok, Thailand 9th-11th February 2016, 116-120.



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved