

## REFERENCES

- [1] Songtham, W., Ugai, H., Imsamut, S. and Maranate, S. 2005. Middle miocene molluscan assemblages in Mae Moh Basin, Lampng Province, Northern Thailand. *Science Asia*. 31. 183-191.
- [2] Pengchai, P., Chantara, S., Sopajaree, K., Wangkarn, S., Tengcharoenkul, U. and Rayanakorn, M. 2009. Seasonal variation, risk assessment and source estimation of PM<sub>10</sub> and PM<sub>10</sub>-bound PAHs in the ambient air of Chiang Mai and Lamphun, Thailand. *Environmental Monitoring Assessment*. 154. 197-218.
- [3] Wangkiat, A. 2007. Chemical characterization and source identification of PM<sub>10</sub> by chemical mass balance model in Mae Moh Area. *Rangsit University Journal of Engineering and Technology*. 10(2). 57-67.
- [4] Pollution Control Department (PCD). 2015. Available from : [http://www.aqmthai.com/public\\_report.php](http://www.aqmthai.com/public_report.php)
- [5] Pimonsree, S., Arrin, P. 2010. Status of particulate problems and their effects on human health in Lampang Province. 6<sup>th</sup> Naresuan Conference, Naresuan University, Phitsanulok Thailand, 29-31<sup>th</sup> july, 2010.
- [6] Li, N., Sioutas, C., Cho, A., Schmitz D, Misra C, Sampf, J., Wang, M. Y., Oberley, T., Froines, J. and Nel, A. 2003. Ultrafine particulate pollutants induce oxidative stress and mitochondrial damage. *Environment Health Perspect*. 111(4). 455-460.
- [7] Renggarajan, T., Rajendran, P., Nandakumar, N., Lokeshkumar, B., Rajendran, L. and Nishigaki, I. 2015. Exposure to polycyclic aromatic hydrocarbons with special focus on cancer. *Asian Pacific Journal of Tropical Biomedicine*. 5(3). 182-189.

- [8] Ruwei, W., Jiamei, Z., Jingjing, L. and Liu, G. 2013. Levels of patterns of polycyclic aromatic hydrocarbons in coal-fired power plant bottom ash and fly ash from Huainan, China. Archives of Environmental Contamination and Toxicology. 65. 193-202.
- [9] Fongmoon, D., Pongnikorn, S., Chaisena, A. and Iamsaard, S. 2014. Particulate matters collected from ceramic factories in Lampang Province affecting rat lung. Biomedicine and Biotechnology. 15(1). 75-83.
- [10] Health Information System Development Office (HISO). 2015. Available from <http://www.hiso.or.th/hiso/motionChart/html/lungcancer5.php>
- [11] Sangram, N., Duangsathaporn, K. and Poolsiri, R. 2016. Effect of gases and particulate matter from electricity generation process on the radial growth of teak plantations surrounding Mae Moh power plant, Lampang Province. Agriculture and Natural Resources. 50. 114-119.
- [12] Pongpiachan, S., Tipmanee, D., Khumsup, C., Kittipoon, I. and Hirunyatrakul, P. 2015. Assessing risks to adults and preschool children posed by PM<sub>2.5</sub>-bound polycyclic aromatic hydrocarbons (PAHs) during a biomass burning episode in Northern Thailand. Science of the Total Environment. 508. 435-444.
- [13] Pollution Control Department (PCD). 2014. Air Quality Index. Available from [http://aqmthai.com/aqi\\_info.php](http://aqmthai.com/aqi_info.php)
- [14] Phoothiwut, S. and Junyapoon, S. 2013. Size distribution of atmospheric particulates and particulate-bound polycyclic aromatic hydrocarbons and characteristics of PAHs during haze period in Lampang Province, Northern Thailand. Air Quality Atmospheric Health. 6. 397-405.
- [15] Sukitpaneenit, M., and Oanh, N, T, K. 2014. Satellite monitoring for carbon monoxide and particulate matter during forest fire episodes in Northern Thailand. Environmental Monitoring Assessment. 186. 2495-2504.

- [16] Geo-Informatics Center for Thailand, Chulalongkorn University.  
Available from <http://www.gisthai.org/map-gallery/3d%20north/Lampang/topo/topo.html>.
- [17] Kouprianov, V. I., Tsakunvatana. V. T. and Douglas, P.L. 2003. Effects of fuel quality on emission rates and specific emissions of major pollutants from lignite based power generation in Thailand. 12<sup>th</sup> Conference on Coal Science, Australia, 2-6<sup>th</sup> November 2003.
- [18] Ridd. M.F., Barber, A.J. and Crow, M.J. 2010. Mae Moh Basin. The Geology of Thailand. The Geological Society. 233.
- [19] Chareonpanich, M., Boonfueng, T. and Limtrakul, J. 2002. Production of aromatic hydrocarbons from Mae-Moh lignite. Fuel Processing Technology. 79. 171-179.
- [20] Brigden, K., Santillo, D. and Stringer, R. 2002. Hazardous emissions from Thai coal-fired power plant: Toxic and potentially toxic elements in fly ashes collected from the Mae Moh and Thai Petrochemical Industry coal-fired power plants in Thailand. Greenpeace Research Laboratories, Department of Biological Sciences, University of Exeter, Exeter, UK.
- [21] Verma, S.K., Masto, R.E., Gautam, S., Choudhury, D.P., Maiti, S.K. and Maity, S. 2015. Investigations on PAHs and trace elements in coal and its combustion residues from a power plant. Fuel. 162. 138-147.
- [22] Teeparut, C. and Sthiannopkao, S. 2012. Mae Moh lignite and environmental management. Geosystem Engineering. 14. 85-94.
- [23] College of Public Health Science, 2001. The study of effect of air pollution on health in Mae Moh district, Lampang province. Department of Health, Ministry of Public Health, Bangkok, Thailand.
- [24] Rahman, M. 2004. A study of the health impact of air pollution from the Mae Moh thermal power plant in Thailand. Environmental Management, Faculty of Graduate studies, Mahidol University.
- [25] EPA Office of Research and Delvelopment. Size comparison for particle.

Available from [https://www.epa.gov/sites/production/files/2016-09/pm2.5\\_scale\\_graphic-color\\_2.jpg](https://www.epa.gov/sites/production/files/2016-09/pm2.5_scale_graphic-color_2.jpg)

- [26] Kawanaka, Y., Matsumoto, E., Wang, N., Yun, S.J. and Sakamoto, K. 2008. Contribution of Nitrated Polycyclic Aromatic Hydrocarbons to the mutagenicity of ultrafine particles in the roadside atmosphere. *Atmospheric environment*. 42. 7423-7428.
- [27] Clean Air Initiative for Asian Cities (CAI-Asia) Center, 2010. Particulate Matter (PM) Standards in Asia. *Air Quality in Asia: Status and Trends*.
- [28] Kim, K.H., Jahan, S.R., Kabir, E. and Brown, R.J.C. 2013. A review of airborne polycyclic aromatic hydrocarbons (PAHs) and their human health effects. *Environment International*. 60. 71-80.
- [29] Kong, S., Ji, Y., Li, Z., Lu, B. and Bai, Z. 2013. Emission and profile characteristic of polycyclic aromatic hydrocarbons in PM<sub>2.5</sub> and PM<sub>10</sub> from stationary sources based on dilution sampling. *Atmospheric Environment*. 77. 156-165.
- [30] Herlekar, M., Elizabeth, A., Kumar, R. and Gupta, I. 2012. Chemical speciation and source assignment of particulate phase molecular markers in Mumbai. *Aerosol and Air Quality Research*. 12. 1247-1260.
- [31] Shi, J., Peng, Y., Li, W., Qiu, W., Bai, Z., King, S. and Jin, T. 2010. Characterization and Source identification of PM<sub>10</sub>-bound polycyclic aromatic hydrocarbons in urban air of Tianjin, China. *Aerosol and Air Quality Research*. 10. 507-518.
- [32] Khalili, N.R., Scheff, P.A. and Holsen, T.M. 1995. PAH source fingerprints for coke ovens, diesel and gasoline engines, highway tunnels and wood combustion emissions. *Atmospheric Environment*. 4. 533-542.
- [33] Miguel, A.H. and Pereira, P.A.P. 1989. Benzo[*k*]fluoranthene, Benzo[*ghi*]perylene and Indeno[1,2,3-*cd*]pyrene: New tracers of aromatic emissions in receptor modeling. *Aerosol Science Technology*. 10. 292-295.

- [34] Chuesaard, T., Chetiyankornkul, T., Kameda, T., Hayakawa, K. and Toriba, A. 2014. Influence of biomass burning on the levels of atmospheric polycyclic aromatic hydrocarbons and their nitro derivatives in Chiang Mai, Thailand. *Aerosol and Air Quality Research*. 14. 1247-1257.
- [35] Mesquita, S.R., Drooge, B.L.V., Barata, C., Viera, N., Guimaraes, L. and Pina, B. 2014. Toxicity of atmospheric particle-bound PAHs: an environmental perspective. *Environmental Science Pollution Research*. 21. 11623-11633.
- [36] Topinka, J., Milcova, A., Schmuczerova, J., Krouzek, J. and Hovorka, J. 2013. Ultrafine particles are not major carriers of carcinogenic PAHs and their genotoxicity in size-segregated aerosols. *Mutation Research/Genetic Toxicology and Environmental Mutagenesis*. 754. 1-6.
- [37] Kim, K.H., Jahan, S.R., Kabir, E. and Brown, R.J.C. 2013. A review of airborne polycyclic aromatic hydrocarbons (PAHs) and their human health effects. *Environment International*. 60. 71-80.
- [38] Perera, F.P., Tang, D., Wang, S., Vishnevetsky, J., Zhang, B. and Diaz, D. 2012. Prenatal polycyclic aromatic hydrocarbon exposure and child behavior at age 6-7 years. *Environment Health Perspect*. 120. 921-926.
- [39] Siddens, L.K., Bunde, K.L., Harper Jr, T.A., McQuistan, T.J., Lohr, C.V., Bramar, L.M., Waters, K.M., Tilton, S.C., Krueger, S.K., Williams, D.E. and Baird, W.M. 2015. Cytochrome P450 1b1 in polycyclic aromatic hydrocarbon (PAH)-induced skin carcinogenesis: tumorigenicity of individual PAHs and coal-tar extract, DNA adduction and expression of select genes in the Cyp1b1 knockout mouse. *Toxicology and Applied Pharmacology*. 287. 149-160.
- [40] Raksasap, R., Panyametheekul, S., Chinwetkitvanich, S. and Pluksasith, T. 2007. Pattern of Particle-bound Polycyclic aromatic hydrocarbons (PAHs) in the vicinity of a power plant in Thailand. *The Proceedings of 1<sup>st</sup> Energy, Environment and Materials Conference*. Bangkok, Thailand.

School of Energy, Environment and Materials, King Mongkut's University of Technology Thonburi (KMUTT). 31<sup>th</sup> August, 2007.

- [41] Norramit, P. Cheevaporn, V. Itoh, N. and Tanaka, K. 2005. Characterization and carcinogenic risk assessment of polycyclic aromatic hydrocarbons in the respirable fraction of airborne particles in The Bangkok Metropolitan Area. *Journal of Health Science*. 51. 437-446.
- [42] Takeshi, O.Takashi, A. Masahiro, F. and Hidetsuru, M. 2004. Spatial Distributions and Profiles of Atmospheric Polycyclic Aromatic Hydrocarbons in Two Industrial Cities in Japan. *Environmental Science and Technology*. 38. 49-55.
- [43] Tolis, E. I., Saraga, D.E., Ammari, G.Z., Gkanas, E.I., Gougoulas, T., Papaioannou, C.C., Sarioglou, A.K., Kougioumtzidis, E., Skemper, A., Bartzis, J.G. 2014. Chemical characterization of particulate matter (PM) and source apportionment study during winter and summer period for the city of Kozani, Greece. *Central European Journal of Chemistry*. 12(6). 634-651.
- [44] Punyawadee, V., Pothisuwan, R., Winichakule, N., Satienerakul, K. 2006. Costs and benefits of flue gas desulfurization for pollution control at the Mae Moh power plant, Thailand. *Economy and Environment Program for Southeast Asia*.
- [45] Harrison, R.M., Smith, D.J.T. and Luhana, L.M., 1996. Source apportionment of atmospheric polycyclic aromatic hydrocarbons collected from an urban location in Birmingham, UK. *Environmental Science Technology*. 30. 825-832.
- [46] Ravindra, K., Sokhi, R. and Grieken, R.V. 2008. Atmospheric polycyclic aromatic hydrocarbons: source attribution, emission factors and regulation. *Atmospheric Environment*. 42. 2895-2921.
- [47] Keshtkar, H. and Ashbaugh, L.L. 2007. Size distribution of polycyclic aromatic hydrocarbon particulate emission factors from agricultural burning. *Atmospheric Environment*. 41. 2729–2739.