CONTENTS

Acknowledgement	с
Abstract in Thai	e
Abstract in English	f
List of Tables	i
List of Figures	j
Chapter 1 Introduction	1
1.1 Historical Background	1
1.2 Objectives	2
1.3 Literature Review	3
1.3.1 Coffee in Thailand	3
1.3.2 Coffee pests	4
1.3.3 Coffee berry borer	5
1.3.4 Coffee berry borer control	7
Chapter 2 Materials and Methods	8
2.1 Ant fauna in coffee plantations	8
2.1.1 Ant collection	8
2.1.2 Data analyses	9
2.2 Ant behavior against a CBB	10
2.2.1 Ant collection	10
2.2.2 Behavior observation	11
2.2.3 Data analyses	12

CONTENTS (CONTINUED)

Chapter 3 Results and discussion	13
3.1 Ant fauna in coffee plantations	13
3.1.1 Ants in PM	13
3.1.2 Ants in KCK	20
3.1.3 Comparison between PM and KCK	21
3.1.4 Comparison with other sites	25
3.1.5 Size of each ant against the CBB	26
3.2 Ant behavior against a CBB	27
Chapter 4 Conclusion	36
References	38
Appendix	43
Curriculum Vitae 49 49 49 49 49 49 49 49 49 49 49 49 49	
ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่ Copyright [©] by Chiang Mai University All rights reserved	

LIST OF TABLES

Table 3.1 Dominant species collected by each method at two study sites	15
in Chiang Mai province	
Table 3.2 Number of ant species collected by each collection method,	16
each season and each site	



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่ Copyright[©] by Chiang Mai University All rights reserved

LIST OF FIGURES

Figure 1.1 Coffee berry borer	6
Figure 2.1 Ant collecting methods	9
Figure 2.2 the breeding case of colony of ant	10
Figure 2.3 The behavioral process of ants against the CBB	11
Figure 3.1 Number of ant species collected per coffee plant and per quadrate	14
in Pa Miang during rainy season (PMR) and dry season (PMD),	
and in Khun Chang Kian during rainy season (KCK)	
Figure 3.2 Species accumulation curves for ants on coffee trees and ants in	18
quadrates in Pa Miag during rainy season (PMR) and dry season	
(PMD), and in Khun Chang Kian during rainy season (KCK)	
Figure 3.3 <i>Technomyrmex</i> ants attend green scale insects	19
Figure 3.4 The green scale insect, Coccus viridis	19
Figure 3.5 Comparison of ant fauna between rainy (PMR) and dry seasons (PMD)	22
in Pa Miang and rainy season in Khun Chang Kian (KCK) with	
matched abundance plot	
Figure 3.6 The most dominant ant species of each coffee plantation	23
Figure 3.7 Coffee plantation at Pa Miang village	24
Figure 3.8 Khun Chang Kian Site A	24
Figure 3.9 Head width distribution of 121 ant species collected in two coffee	26
plantations in Chiang Mai	
Figure 3.10 Seven ant species that used in experiment as behavior observation	27
Figure 3.11 The sequence of behaviors of seven ant species in the presence	29
of a CBB in a plastic case under the laboratory	
Figure 3.12 Average time of each behavior performed by each ant species	31
Figure 3.13 The workers of Tapinoma indicum gathered around a CBB	32
Figure 3.14 The body size of each ant species and the CBB	34

LIST OF FIGURES (CONTINUED)

Page

Figure 3.15 A worker of *Dolichoderus* sp. PM1 did antennal contacting to a CBB34Figure 3.16 Frequency of each behavior displayed by the seven ant species35reacted to the CBB35



ลิ<mark>ปสิทธิ์มหาวิทยาลัยเชียงใหม่</mark> Copyright[©] by Chiang Mai University All rights reserved