

CONTENTS

	Page
Acknowledgement	c
Abstract in Thai	d
Abstract in English	f
List of Tables	j
List of Figures	m
Chapter 1 Introduction	1
1.1 Objectives	4
Chapter 2	5
Chapter 3	15
Chapter 4	36
Chapter 5	108
References	111
Appendix	125
Appendix A	126
Appendix B	130
Appendix C	134
Appendix D	136
Appendix E	139
Appendix F	143
Curriculum Vitae	140



LIST OF TABLES

	Page
Table 4.1 Physicochemical properties of different ripening stages of papaya	37
Table 4.2 Color values of papaya peel and flesh affected by fruit ripening stages	40
Table 4.3 Water activity and chemical properties of impregnated papaya affected by ripening stages and papaya sizes	43
Table 4.4 Color values and texture of impregnated papaya affected by ripening stages and papaya sizes	44
Table 4.5 Real porosity (ϵ_r), apparent (ρ_a) and real densities (ρ_r) of impregnated papaya affected by ripening stages and papaya sizes	49
Table 4.6 Effective porosity (ϵ_e), X and γ values of impregnated papaya affected by ripening stages and papaya sizes	50
Table 4.7 Water loss, solid gain and weight reduction of impregnated papaya affected by ripening stages and papaya sizes	51
Table 4.8 Water activity and chemical properties of impregnated papaya affected by impregnation solution ratio and vacuum impregnation condition	56
Table 4.9 Color values and texture of impregnated papaya affected by impregnation solution ratio and vacuum impregnation condition	57
Table 4.10 Weight loss, solid gain and weight reduction of impregnated papaya affected by impregnation solution ratio and vacuum impregnation condition	58
Table 4.11 X value, γ value and effective porosity of impregnated papaya affected by impregnation solution ratio and vacuum impregnation condition	59
Table 4.12 Apparent density, real density and real porosity of impregnated papaya affected by impregnation solution ratio and vacuum impregnation condition	60

Table 4.13 Water activity and chemical properties of partially dried papaya affected by drying methods and temperatures	65
Table 4.14 Yield, texture, real porosity and sample volume deformation of partially dried papaya affected by drying methods and temperatures	66
Table 4.15 Color values of partially dried papaya affected by drying methods and temperatures	70
Table 4.16 Sensory preferences of partially dried papaya affected by drying methods and temperatures	73
Table 4.17 Water activity, chemical properties and yield of partially dried papaya affected by calcium types and concentrations and impregnation temperatures	76
Table 4.18 Real porosity, γ value, shrinkage and texture of partially dried papaya affected by calcium types and concentrations and impregnation temperatures	77
Table 4.19 Color values of partially dried papaya affected by calcium types and concentrations and impregnation temperatures	83
Table 4.20 Sensory preferences of partially dried papaya affected by different calcium lactate concentrations	84
Table 4.21 The physicochemical and microbial properties of partially dried papaya containing lactic acid bacteria	90
Table 4.22 The physicochemical properties of partially dried papaya containing with <i>L.casei</i> TISTR 390 stored at room temperature	96
Table 4.23 The physicochemical properties of partially dried papaya containing with <i>L.casei</i> TISTR 390 stored at 4°C	97
Table 4.23 The physicochemical properties of partially dried papaya containing with <i>L.casei</i> TISTR 390 stored at 4°C (continued)	98
Table 4.24 The physicochemical properties of partially dried papaya containing with <i>L.casei</i> TISTR 390 stored at -18°C	99
Table 4.24 The physicochemical properties of partially dried papaya containing with <i>L.casei</i> TISTR 390 stored at -18°C (continued)	100
Table 4.25 The number of yeast and mold (CFU/g) of partially dried	107

papaya containing with *L.casei* TISTR 390 stored at room temperature, 4 °C and -18°C



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved

LIST OF FIGURES

	Page
2.1 'Pluk Mai Lie' papaya	7
2.2 Physical characteristics of Pluk Mai Lie papaya	7
4.1 <i>L. casei</i> in intracellular space of impregnated papaya, magnification: 5,000 and 10,000	91
4.2 <i>L. casei</i> in intracellular space of dried impregnated papaya, magnification: 1,000 and 10,000	92
4.3 The number of lactic acid bacteria and total microorganism (log cfu/g) in partially dried papaya containing <i>L.casei</i> TISTR 390 at room temperature for 4 weeks	102
4.4 The number of lactic acid bacteria and total microorganism (log cfu/g) in partially dried papaya containing <i>L.casei</i> TISTR 390 at 4 °C for 12 weeks	103
4.5 The number of lactic acid bacteria and total microorganism (log cfu/g) in partially dried papaya containing <i>L.casei</i> TISTR 390 at -18 °C for 12 weeks	103