

## REFERENCES

- Alexopoulos, G. S., & Bruce, M. L. (2009). A model for intervention research in late-life depression. *International Journal of Geriatric Psychiatry*, 24, 1325-1334.
- Almeida, O. P. (2012). Approaches to decrease the prevalence of depression in late life. *Geriatric Psychiatry*, 25(6), 451-456.
- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.) Washington, D.C.: Author.
- American Psychiatric Association. (2016). *Older adults' health and age-related changes*. Retrieved from <http://www.apa.org/pi/aging/resources/guides/older.aspx>
- Amick, H. R., Gartlehner, G., Gaynes, B. N., Forneris, C., Asher, G. N., Morgan, L. C., . . . Lohr, K. N. (2015). Comparative benefits and harms of second generation antidepressants and cognitive behavioral therapies in initial treatment of major depressive disorder: Systematic review and meta-analysis. *The BMJ*, 351, h6019. doi:10.1136/bmj.h6019
- Andreescu, C., & Reynolds, C. F., III (2011). Late-life depression: Evidence-based treatment and promising new directions for research and clinical practice. *The Psychiatric Clinics of North America*, 34, 335-355.
- Arandjelovic, K., Eyre, H. A., Forbes, M., & Lavretsky, H. (2016). Treatment options in late-life treatment-resistant depression. Retrieved from <http://www.psychiatrictimes.com/geriatric-psychiatry/treatment-options-late-life-treatment-resistant-depression>
- Arean, P. A. (2012). Psychotherapy for late-life depression. *Psychiatric Times*, 29(8), 35-38.
- Ari, H., Ouchi, Y., Yokode, M., Ito, H., Uematsu, H., Eto, F., . . . Kita, T. (2012). Toward the realization of a better aged society: Messages from gerontology and geriatric. *Geriatrics & Gerontology International*, 12, 16-22.

Australian Psychological Society. (2010). *Evidence-based psychological interventions in the treatment of mental disorders: A literature review* (3rd ed.). Flinders Lane: The Australian Psychological Society.

Baune, B. T., Stuart, M., Wersching, H., Heindel, W., Arolt, V., & Berger, K. (2012). The relationship between subtypes of depression and cardiovascular disease: A systematic review of biological models. *Translational Psychiatry*, 2, e92. doi:10.1038/tp.2012.18

Beck, A. T. (1967). *Depression: Causes, and treatment*. Philadelphia: University of Pennsylvania Press.

Beck, A. T. (1976). *Cognitive therapy and the emotional disorders*. New York: international Universities Press.

Beck, A. T., & Alford, B. A. (2009). *Depression causes and treatment* (2nd ed.). Philadelphia: University of Pennsylvania Press.

Beck, A. T., Rush, A. J., Shaw, B. F., & Emery, G. (1979). *Cognitive therapy of depression*. New York: The Guilford Press.

Beck, J. S. (1995). *Cognitive therapy: Basic and beyond*. New York: The Guilford Press.

Beck, J. S. (2011). *Cognitive behavior therapy basic and beyond* (2nd ed.). New York: The Guilford Press.

Beekman, A. T. F., Copeland, J. R., & Prince, M. J. (1999). Review of community prevalence of depression in later life. *The British Journal of Psychiatry*, 174(4), 307-311.

Beidas, R. S., Cross, W., & Dorsey, S. (2014). Show me, don't tell me: Behavioral rehearsal as a training and analogue fidelity tool. *Cognitive and Behavioral Practice*, 21, 1-11.

- Beland, D., & Viriot Durandal, J.-P. (2013). Aging in France: Population trends, policy issues, and research institutions. *The Gerontologist*, 53(2), 191-197.
- Beyermann, S., Trippe, R. H., Bähr, A. A., & Püllen, R. (2013). Mini-Mental status examination in geriatrics: An evaluation of diagnostic quality. *Zeitschrift für Gerontologie und Geriatrie*, 46(8), 740-747.
- Bian, C., Li, C., Duan, Q., & Wu, H. (2011). Reliability and validity of patient health questionnaire: Depressive syndrome module for outpatients. *Scientific Research and Essays*, 6(2), 278-282.
- Blazer, D. G. (2003). Depression in late life: Review and commentary. *Journal of Gerontology*, 58(3), 249-265. doi:10.1093/gerona/58.3.M249
- Blazer, D. G. (2009). Depression in late life: Review and commentary. *Focus*, 7(1), 118-136.
- Blazer, D. G., Hybels, C. F., & Pieper, C. F. (2001). The Association of Depression and Mortality in Elderly Persons: A case for multiple, independent pathways. *Journal of Gerontology*, 56A(8), M505-M509.
- Bojorquez-Chapela, I., Villalobos-Daniel, V. E., Manrique-Espinoza, B. S., Tellez-Rojo, M. M., & Salinas-Rodríguez, A. (2009). Depressive symptoms among poor older adults in Mexico: Prevalence and associated factors. *Pan American Journal of Public Health*, 26(1), 70-77.
- Bull World Health Organization, Thailand. (2012). The health-care challenges posed by population ageing. *Bull World Health Organization*, 90, 82-83.  
doi:10.2471/BLT.12.020212
- Bureau of Empowerment for Older Person of Thailand. (2012). *Annual report Thai aging situation 2010*. Foundation of Thai Gerontology Research and Development Institute (TGRI).

- Burnett-Ziegler, I. E., Pfeiffer, P., Zivin, K., Glass, J. E., Ilgen, M. A., Flynn, H. A., Austin, K., & Chermack, S. T. (2012). Psychotherapy utilization for acute depression within the Veterans Affairs health care system. *Psychological Services*, 9(4), 325-335.
- Buszewicz, M., & Murphy, B. (2016). *Supporting people with depression and anxiety: A guide for practice nurses*. Retrieved from [http://www.mind.org.uk/media/944494/MIND\\_ProCEED\\_Training\\_Pack.pdf?ctaid=/about-us/policies-issues/proceed/slices/text-2/](http://www.mind.org.uk/media/944494/MIND_ProCEED_Training_Pack.pdf?ctaid=/about-us/policies-issues/proceed/slices/text-2/)
- Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion. (2015). *Depression is not a normal part of growing older*. US: Division of Population Health.
- Chand, S. P., & Grossberg, G. T. (2013). How to adapt cognitive-behavioral therapy for older adults. *Current Psychiatry*, 12(3), 10-15.
- Chapman, D. P., & Perry, G. S. (2008). Depression as a major component of public health for older adults. *Preventing Chronic Disease*, 5(1), 1-9.
- Charernboon, T., Phanasathit, M., Tangwongchai, S., Hemrungroj, S., & Worakul, P. (2010). Depression and dementia among members of 15 elderly clubs in Bangkok. *Thammasat Medical Journal*, 10(4), 428-436.
- Chen, S., Conwell, Y., Xu, B., Chiu, H., Tu, X., & Ma, Y. (2011). Depression care management for late-life depression in China primary care: Protocol for a randomized controlled trial. *Trials*, 12, 121. doi:10.1186/1745-6215-12-121
- Chisholm, D., Sanderson, K., Ayuso-Mateos, J. L., & Saxena, S. (2004). Reducing the global burden of depression: Population-level analysis of intervention cost-effectiveness in 14 world regions. *The British Journal of Psychiatry*, 184, 393-403.

- Cho, M. J., & Lee, J. Y. (2005). Epidemiology of depressive disorders in Korea. *Psychiatry Investigation*, 2(1), 22-27.
- Chou, K.-L., Jun, L. W., & Chi, I. (2005). Assessing Chinese older adults' suicidal ideation: Chinese version of the Geriatric Suicide Ideation Scale. *Aging & Mental Health*, 9(2), 167-171.
- Clark, D. A., Beck, A. T., & Stewart, B. L. (1990). Cognitive specificity and positive-negative affectivity: Complementary or contradictory views on anxiety and depression? *Journal of Abnormal Psychology*, 99(2), 148-158.
- Colom, F., Vieta, E., Sanchez-Moreno, J., Palomino-Otiniano, R., Reinares, M., Goikolea, J. M., . . . Martinez-Aran, A. (2009). Group psychoeducation for stabilised Bipolar disorders: 5-year outcome of a randomised clinical trial. *British Journal of Psychiatry*, 194(3), 260-265.
- Cooper, C., Katona, C., Lyketsos, K., Blazer, D., Brodaty, H., Rabins, P., . . . Livingston, G. (2011). A systematic review of treatments for refractory depression in older people. *American Journal of Psychiatry*, 168(7), 681-688.  
doi:10.1176/appi.ajp.2011.10081165
- Cox, D., & D'Oyley, H. (2011). Cognitive-behavioral therapy with older. *BC Medical Journal*, 53(7), 348-352.
- Cuijpers, P., van Straten, A., & Smit, F. (2006). Psychological treatment of late-life depression: A meta-analysis of randomized controlled trials. *International Journal of Geriatric Psychiatry*, 21, 1139-1149.
- Cusin, C., Yang, H., Yeung, A., & Fava, M. (2009). Rating Scales for Depression. In L. Baer & M. A. Blais (Eds.), *Handbook of Clinical Rating Scales and assessment in psychiatry and mental health, current clinical psychiatry*. Boston: Human Press.
- D'Zurilla, T. J., & Goldfried, M. R. (1971). Problem solving and behavior modification. *Journal of Abnormal Psychology*, 78, 107-126.

D'Zurilla, T. J., & Nezu, A. M. (1990). Development and preliminary evaluation of the social problem-solving inventory (SPSI) psychological assessment. *Journal of Consulting and Clinical Psychology*, 2, 156-163.

D'Zurilla, T. J., & Nezu, A. M. (2007). *Problem-solving therapy: A positive approach to clinical intervention* (3rd ed.). New York: Springer.

de la Cámare, C., Saz, P., López-Antón, R., Ventura, T., Día, J. L., & Lobo, A. (2008). Depression in the elderly community: I. Prevalence by different diagnostic criteria and clinical profile. *The European Journal of Psychiatry*, 22(3), 131-140.

Department of Economic and Social Affairs. (2011). *Current status of the social situation, well-being, participation in development and rights of older persons worldwide*. New York: United Nations.

Department of Mental Health, Ministry of Public Health of Thailand. (2010). *Clinical practice guideline of major depressive disorder for general practitioner: CPG-MDD-GP*. Nonthaburi: Department of Mental Health.

Department of Mental Health, Ministry of Public Health of Thailand. (2012). *Clinical practice guideline of psychosocial intervention for depressive disorder in secondary care*. Nonthaburi: Department of Mental Health.

Department of Mental Health, Ministry of Public Health, Thailand. (2014). *Guidebook of depressive disorders surveillance and care: Provincial level* (Revised Edition III). Nonthaburi: Department of Mental Health.

DeRubeis, R. J., Webb, C. A., Tang, T. Z., & Beck, A. T. (2010). Cognitive therapy. In K. S. Dobson (Ed.), *Handbook of cognitive-behavioral therapies* (3rd ed.). New York: The Guilford Press.

Dobson, D. J. G., & Dobson, K. S. (2013). In-session structure and collaborative empiricism. *Cognitive and Behavioral Practice*, 20, 410-418.

- Dobson, K. S. (2010). *Handbook of cognitive behavioural therapies*. New York: Guilford Press.
- Donker, T., Griffiths, K. M., Cuijpers, P., & Christensen, H. (2009). Psychoeducation for depression, anxiety and psychological distress: a meta-analysis. *BMC Medicine*, 7, 79. doi:10.1186/174-7015-7-79
- Dykert, D., Der, G., Starr, J. M., & Deary, I. J. (2016). Why is Mini-Mental state examination performance correlated with estimated premorbid cognitive ability? *Psychological Medicine*, 46, 2647- 2654.
- Ekers, D., Richards, D., & Gilbody, S. (2008). A Meta-analysis of randomized trials of behavioural treatment of depression. *Psychological Medicine*, 38(5), 611-623.
- Ekers, D., Webster, L., Van Straten, A., Cuijpers, P., Richards, D., & Gilbody, S. (2014). Behavioural activation for depression: An update of meta-analysis of effectiveness and sub group analysis. *PLoS One*, 9(6), e100100. doi:10.1371/journal.pone.0100100
- Evans, C. (2007). Cognitive-behavioural therapy with older people. *Advances in Psychiatric Treatment*, 13, 111-118.
- Ezeh, A. C. , Bongaarts, J, & Mberu, B. (2012). Global population trends and policy options. *Lancet*, 380, 142-148.
- Farinde, A. (2013). The Beck Depression Inventory. *The Pharma Innovation*, 2(1), 56-62.
- Fernandaz, M. E., Mutran, E. J., Reitzes, D. C., & Sudha, S. (1998). Ethnicity, gender, and depressive symptoms in older workers. *The Gerontologist*, 38(1), 71-79.
- Ferrari, A. J., Charlson, F. J., Norman, R. E., Patten, S. B., Freedman, G., Murray, C. J. L., . . . Whiteford, H. A. (2013). Burden of depressive disorders by country, sex, age, and year: Findings from the Global Burden of Disease Study 2010. *PLOS Medicine*, 10(11), 1-12.

- Field, A. (2009). *Discovering statistics using SPSS* (3rd ed.). London: SAGE.
- Fiske, A., Wetherell, J. L., & Gatz, M. (2009). Depression in older adults. *Annual Review of Clinical Psychology*, 5, 363-389. doi:10.1146/annurev.clinpsy.032408.153621
- Flood, M., & Buckwalter, K. C. (2009). Recommendations for mental health care of older adults: Part 1—An overview of depression and anxiety. *Journal of Gerontological Nursing*, 35(2), 26-34.
- Folstein, M. F., Folstein, S. E., & McHugh, P. R. (1975). “Mini-Mental-State”: A practical method for grading the cognitive state of patients for the clinician. *Journal of Psychiatric Research*, 12, 189-198.
- Foo, K. H., & Kazantzis, N. (2007). Integrating homework assignments based on culture: Working with Chinese patients. *Cognitive and Behavioral Practice*, 14, 333-340.
- Forlani, C., Morri, M., Ferrari, B., Dalmonte, E., Menchetti, M., De Ronchi, D., & Atti, A. R. (2014). Prevalence and gender differences in late-life depression: A population-based study. *American Journal of Geriatric Psychiatry*, 22(4), 370-380.
- Foundation for Older Persons' Development. (2015). *Foundation for older persons' development, Thailand: Situation of the Thai elderly (Population situations)*. Retrieved from <http://fopdev.or.th/situation-of-the-thai-elderly-population-situations/>
- Francis, J. L., & Kumar, A. (2013). Psychological treatment of late-life depression. *Psychiatric Clinical of North America*, 36, 561-575.
- Frazer, C. J., Christensen, H., & Griffiths, K. M. (2005). Systematic review effectiveness of treatments of depression in older people. *The Medical Journal of Australia*, 182(12), 627-632.

- Frisch, N. C., & Frisch, L. (2002). *Psychiatric mental health nursing* (2nd ed.). Australia: Delmar Thomson Learning.
- Froggatt, W. (2006). *A brief introduction to cognitive-behaviour therapy*. Retrieved from <http://www.rational.org.nz/prof-docs/Intro-CBT.pdf>
- Fual, F., Erdfelder, E., Buchner, A., & Lang, A. (2009). Statistical power analyses using G\*Power 3.1: Tests for correlation and regression analyses. *Behavior Research Methods*, 41(4), 1149-1160.
- Gautam, R., & Houde, S. C. (2011). Geriatric Depression Scale for community dwelling older adults in Nepal. *Asian Journal of Gerontology & Geriatrics*, 6(2), 93-99.
- Gonzalez-Pinto, A., Gonzalez, C., Enjuto, S., Fernandez de Corres, B., Lopez, P., Palomo, J., . . . Perez de Heredia, J. L. (2004). Psychoeducation and cognitive-behavioral therapy in bipolar disorder: An update. *Acta Psychiatrica Scandinavica*, 109, 83-90.
- Gould, R. L., Coulson, M. C., & Howard, R. J. (2012). cognitive behavioral therapy for depression in older people: A meta-analysis and meta-regression of randomized controlled trials. *Journal of the American Geriatrics Society*, 60(10), 1817-1830. doi:10.1111/j.1532-5415.2012.04166.x
- Government Public Relations Department, Thailand. (2014). *Thailand prepares for becoming an aging society*. Retrieved from [http://thailand.prd.go.th/ewt\\_news.php?nid=1268&filename=index](http://thailand.prd.go.th/ewt_news.php?nid=1268&filename=index)
- Graham, C. C., Campion, J., Kaiser, P., & Edwards, K. (2011). Management of depression in older people: why this is important in primary care. Retrieved from [http://www.rcgp.org.uk/pdf/NMH\\_10095\\_OPMH%20%20depression%20%20Feb%202011.pdf](http://www.rcgp.org.uk/pdf/NMH_10095_OPMH%20%20depression%20%20Feb%202011.pdf)

- Greenberg, L., & Goldman, R. (2007). Case formulation in Emotion-focused therapy. In T. Ells (Ed.), *Handbook of psychotherapy case formulation* (pp. 379-412). New York: Guilford Press.
- Grove, S. K., Burns, N., & Gray, J. R. (2013). *The practice of nursing research: Appraisal, synthesis, and generation of evidence* (7th ed.). Philadelphia: W.B. Saunders.
- Grove, S. K., Gray, J. R., & Burns, N. (2015). *Understanding nursing research: Building an evidence-based practice* (6th ed.). St. Louis: Elsevier Saunders.
- Hardy, S. (2011). Depression in the elderly: Ways to offer support. *Practice Nursing*, 22(10), 520-525.
- Harris, D. L., & Hiskey, S. (2015). Homework in therapy: a case of it ain't what you do, it's the way that you do it? *The Cognitive Behaviour Therapist*, 8(e20), 1-11.
- Haseen, F., & Prasartkul, P. (2011). Predictors of depression among older people living in rural areas of Thailand. *Bangladesh Medical Research Council Bulletin*, 37(2), 51-56.
- Haynes, S. N., & O'Brien, W. H. (2000). *Principles and practice of behavioral assessment*. New York: Kluwer Academic/Plenum Press.
- Hofmann, S. G., Asmundson, G. J. G., & Beck, A. T. (2013). The science of cognitive therapy. *Behavior Therapy*, 44(2), 199-212.
- Hofmann, S. G., Asnaani, A., Vonk, I. J. J., Sawyer, A. T., & Fang, A. (2012). The efficacy of cognitive behavioral therapy: A review of meta-analyses. *Cognitive Therapy and Research*, 36(5), 427-440. doi:10.1007/s10608-012-9476-1
- Holman, A. J., Serfaty, M. A., Leurent, B. E., & King, M. B. (2011). Cost-effectiveness of cognitive behaviour therapy versus talking and usual care for depressed older people in primary care. *BMC Health Service Research*, 11, 33. doi:10.1186/1472-6963-11-33

Holmes, J. (2002). All you need is cognitive behaviour therapy? *The BMJ*, 324, 288-294.

Holroyd, S., & Clayton, A. H. (2000). Measuring depression in the elderly: Which scale is best? *Medscape General Medicine*, 2(4), 1-4.

Hooyman, N. R., & Kiyak, H. A. (2009). *Social gerontology: A multidisciplinary perspective* (8th ed.). Boston: Pearson Education.

Huntley, A. L., Araya, R., & Sallisbury, C. (2012). Group psychological therapies for depression in the community: Systematic review and meta-analysis. *The British Journal of Psychiatry*, 200(3), 184-190. doi:10.1192/bjp.bp.111.092049

Hwong, W.-C., Wood, J. J., Lin, K.-M., & Cheung, F. (2006). Cognitive-behavioral therapy with Chinese Americans: Research, theory, and clinical practice. *Cognitive and Behavioral Practice*, 13(4), 293-303.

Imel, Z. E., Malterer, M. B., McKay, K. M., & Wampold, B. E. (2008). A meta-analysis of psychotherapy and medication in unipolar depression and dysthymia. *Journal of Affective Disorders*, 110(3), 197-206. doi:10.1016/j.jad.2008.03.018

Institute of Geriatric Medicine, Department of Public Health, Thailand. (2006). *Research Project: Survey and study of health status among older adults in 4 regions of Thailand*. Retrieved from <http://hpc12anamai.moph.go.th/home/LTC53/4regions-R1.pdf>

Institute of Geriatric Medicine, Department of Public Health, Thailand. (2002). *Mini-Mental status examination-Thai 2002*. Nonthaburi: Institute of Geriatric Medicine.

Jang, Y., Gum, A. M., & Chiriboga, D.A. (2011). Knowledge of depression among Korean American older adults. *Journal of Applied Gerontology*, 30(5), 655-665.

Jang, Y., Kim, G., & Chiriboga, D. A. (2011). Gender Differences in Depressive Symptoms Among Older Korean American Immigrants. *Social Work in Public Health*, 26, 96-109.

- Jarrett, R. B., Kraft, D., Schaffer, M., Witt-Browder, A., Risser, R., Atkins, D. H., & Doyle, J. (2000). Reducing relapse in depressed outpatients with atypical features: A pilot study. *Psychotherapy and Psychosomatics*, 69(5), 232-239.
- Joo, J. H., Morales, K. H., de Vries, H. F., & Gallo, J. J. (2010). Disparity in use of psychotherapy offered in primary care between older African-American and White adults: Results from a practice-based depression intervention trial. *Journal of the American Geriatrics Society*, 58(1), 154-159. doi:10.1111/j.1532-5415.2009.02623.x
- Joyst, K. E., Whellan, D. J., & O'Connor, C. M. (2003). Depression and cardiovascular disease: Mechanisms of interaction. *Biological Psychiatry*, 54(3), 248-261.
- Kahlon, S., Neal, A., & Patterson, T. G. (2014). Experiences of cognitive behavioural therapy formulation in clients with depression. *The Cognitive Behaviour Therapist*, 7(e8), 1-15.
- Karel, M. J., & Hinrichsen, G. (2000). Treatment of depression in late life: Interventions. *Clinical Psychology Review*, 20(6), 707-729.
- Kazantzis, N., Pachana, N. A., & Secker, D. L. (2003). Cognitive behavioral therapy for older adults: Practical guidelines for the use of homework assignments. *Cognitive and Behavioral Practice*, 10, 324-332.
- Kazantzis, N., Whittington, C., & Dattilio, F. (2010). Meta-analysis of homework effects in cognitive and behavioral therapy: Replication and extension. *Clinical Psychology: Science and Practice*, 17, 144-156.
- Kessler, R. C., & Bromet, E. J. (2013). The epidemiology of depression across cultures. *Annual Review of Public Health*, 34, 119- 138.

- Khonkaen Psychiatric Hospital, Department of Mental Health, Ministry of Public Health of Thailand. (2011). *Research report: Effectiveness of cognitive behavioral therapy group to reduce negative automatic thought and depressive symptoms with psychiatric patients*. Khonkaen: Khonkaen Publishing.
- Kim, H.-Y. (2012). Statistical notes for clinical researchers: assessing normal distribution (1). *Restorative Dentistry and Endodontic*, 37(4), 245-248.
- Kim, H.-Y. (2013). Statistical notes for clinical researchers: Assessing normal distribution (2) using skewness and kurtosis. *Restorative Dentistry and Endodontic*, 38(1), 52-54.
- Kim, J.-I., Choe, M. A., & Chae, Y. R. (2009). Prevalence and predictors of geriatric depression in community-dwelling elderly. *Asian Nursing Research*, 3(3), 121-129.
- Kitsumban, V., Thapinta, D., Sirindharo, P. B., & Anders, R. L. (2009). Effect of cognitive mindfulness practice program on depression among elderly Thai women. *Thai Journal of Nursing Research*, 13(2), 95-108.
- Kittirattanapaiboon, P., Tantirangsee, N., Chutha, W., Assanangkornchai, S., & Supanya, S. (2016). Thai national mental health survey 2013: Methodology and procedure. *Journal of Mental Health of Thailand*, 24(1), 1-14.
- Klerman, G. L., & Weissman, M. M. (1993). *New application of interpersonal psychotherapy*. London: American Psychiatric Press.
- Kongsuk, T. (2012). *Prevention and management of major depressive disorders for accessibility service problem: Financialy year 2012*. Ubon ratchathani: Prasrimahabhodi Psychiatric Hospital.
- Kroenke, K., & Spitzer, R. L. (2002). The PHQ-9: A new depression and diagnostic severity measure. *Psychiatric Annals*, 32, 509-521.

- Kroenke, K., Spitzer, R. L., & Williams, J. B. (2001). The PHQ-9 validity of a Brief Severity Measure. *Journal of General Internal Medicine*, 16(9), 606-613.
- Kuptniratsaikul, V., & Pekumal, P. (1997). The Center for Epidemiologic Studies-Depression scale (CES-D) in Thai People. *Sriraj Medical Journal*, 49, 442- 448.
- Kurlowicz, L. (2003). Depression in older adults. In M. Mezey, T. Fulmer, & I. Abraham (Eds.), *Geriatric nursing protocols for best practice* (2nd ed., pp. 21-23). New York: Springer.
- Kuyken, W., Padesky, C. A., & Dudley, R. (2008). Process issues: The science and practice of case conceptualization. *Behavioural and Cognitive Psychotherapy*, 36, 757-768.
- Laidlaw, K., Davidson, K., Toner, H., Jackson, G., Clark, S., Law, J., . . . Cross, S. (2008). A randomised controlled trial of cognitive behavior therapy vs treatment as usual in the treatment of mild to moderate late life depression. *International Journal of Geriatric Psychiatry*, 23(8), 843-850. doi:10.1002/gps.1993
- Laidlaw, K., Thompson, L. W., Dick-Siskin, L., & Gallagher-Thompson, D. (2004). *Cognitive behaviour therapy with older people* (Reprinted). England: John Wiley & Sons.
- Lamis, D. A., & Kaslow, N. J. (2014). Mediators of the daily hassles-suicidal ideation link in African American women. *Suicide & Life-Threatening Behavior*, 44(3), 233-245. doi:10.1111/sltb.12099
- Lee, S. Y., Franchetti, M. K., Imanbayev, A., Gallo, J. J., Spira, A. P., & Lee, H. B. (2012). Non-pharmacological prevention of major depression among community-dwelling older adults: A systematic review of the efficacy of psychotherapy interventions. *Archives of Gerontology and Geriatrics*, 55(3), 522-529. doi:10.1016/j.archger.2012.03.003

- Lenze, E. J., Rogers, J. C., Martire, L. M., Mulsant, B. H., Rollman, B. L., Dew, M. A., . . . Reynolds, C. F., III. (2001). The Association of Late-Life Depression and Anxiety With Physical Disability: A review of the literature and prospectus for future research. *The American Journal of Geriatric Psychiatry*, 9(2), 113-135.
- Li, D., Zhang, D. J., Shao, J. J., Qi, X. D., & Tian, L. (2014). A meta-analysis of the prevalence of depressive symptoms in Chinese older adults. *Archives of Gerontology and Geriatrics*, 58(1), 1-9.
- Limpawattana, P., Sawanyawisuth, K., Soonpornrai, S., & Huangthaisong, W. (2011). Prevalence and recognition of geriatric syndromes in an outpatient clinic at atertiary care hospital of Thailand. *Asian Biomedicine*, 5(4), 493-497.
- LoBiondo-Wood, G., & Haber, J. (2014). Nursing research: Methods and critical appraisal for evidence-based practice (8th ed.). St.Louis: Elsevier.
- Lorenzo-Luaces, L., German, R. E., & DeRubeis, R.J. (2015). It's complicated: The relation between cognitive change procedures, cognitive change, and symptom change for depression. *Clinical Psychology Review*, 41, 3-15.
- Lortakul, M., & Saipanish, R. (2009). How do general practioners in Thailand diagnose and treat patients presenting with anxiety and depression? *Psychiatry and Clinical Neurosciences*, 63, 37-42.
- Lortakul, M., & Sukanich, P. (1999). Development of the Thai depression inventory. *Journal of Medical Association of Thailand*, 82(12), 1200-1207.
- Lortakul, M., Sumrithe, S., & Saipanish, R. (2008). Reliability and validity of the Thai version of the PHQ-9. *BMC Psychiatry*, 8, 46. doi:10.1186/1471-244X-8-46
- Luijendijk, H. J., van den Berg, J. F., Dekker, M. J. H. J., van Tuijl, H. R., Otte, W., Smit, F., . . . Tiemeier, H. (2008). Incidence and recurrence of late-life depression. *Archives of General Psychiatry*, 65(12), 1394-1401. doi:10.1001/archpsyc.65.12.1394

- Lukens, E. P., & McFarlane, W. R. (2004). Psychoeducation as evidence-based practice: Consideration for practice, research, and policy. *Brief Treatment and Crisis Intervention*, 4(3), 205-225.
- Lum, T. Y., Parashuram, S., Shippee, T. P., Wysocki, A., Shippee, N. D., Homyak, P., & Kane, P. (2012). Diagnosed prevalence and health care expenditures of mental health disorders among dual eligible older people. *The Gerontologist*, 53(2), 334-344.
- Mackin, R. S., & Arean, P. A. (2005). Evidenced-based psychotherapeutic interventions for geriatric depression. *Psychiatric Clinics of North America*, 28(4), 805-820. doi:10.1016/j.psc.2005.09.009
- Magnil, M., Gunnarsson, R., Bjorkstedt, K., & Bjorkelund, C. (2008). Prevalence of depressive symptoms and associated factors in elderly primary care patients: A descriptive study. *Primary Care Companion to the Journal of Clinical Psychiatry*, 10(6), 462-468.
- Malhotra, R., Chan, A., & Ostbye, T. (2010). Prevalence and correlates of clinically significant depressive symptoms among elderly people in Sri Lanka: Finding from a national survey. *International Psychogeriatrics*, 22(2), 227-236.
- Marc, L.G., Raue, P. J., & Bruce, M. L. (2008). Screening Performance of the Geriatric Depression Scale (GDS-15) in a diverse elderly home care population. *The American Journal of Geriatric Psychiatry*, 16(11), 914-921.
- Marcus, M., Yasamy, M. T., van Ommeren, M., Chisholm, D., & Saxena, S. (2012). *Depression: A global public health concern*. Geneva: WHO, Department of Mental Health and Substance Abuse.
- McAvay, G. J., Bruce, M. L., Raue, P. J., & Brown, E. L. (2004). Depression in elderly homecare patients: Patient versus informant reports. *Psychological Medicine*, 34, 1507-1517.

- McCann, T. V., Songprakul, W., & Stephenson, J. (2015). Effectiveness of guided self-help in decreasing expressed emotion in family caregivers of people diagnosed with depression in Thailand: A randomized controlled trial. *BMC Psychiatry*, 15, 1-8.
- McNaughton, J. L. (2009). Brief interventions for depression in primary care: A systematic review. *Canadian Family Physician*, 55(8), 789-796.
- Mokekhaw, S. (2011). The effect of cognitive behavioral therapy program on depression of patient with major depressive disordered. *Thai Journal of Nursing Council*, 26(2), 126-136.
- Montorio, I., & Izal, M. (1996). The Geriatric Depression Scales: A review of its development and utility. *International Psychogeriatrics*, 8(1), 103-112.
- Mor, N., & Haran, D. (2009). Cognitive-behavioral therapy for depression. *The Israel Journal of Psychiatry and Related Sciences*, 46(4), 269-273.
- Moriarty, A. S., Gilbody, S., McMillan, D., & Manea, L. (2015). Screening and case finding for major depressive disorder using the Patient Health Questionnaire (PHQ-9): A meta-analysis. *General Hospital Psychiatry*, 37(6), 567-576.  
doi:10.1016/j.genhosppsych.2015.06.012
- Mulyalya, K. P., & Varghese, M. (2010). The complex relationship between depression and dementia. *Annals of Indian Academy of Neurology*, 13(Suppl. 2), s69-s73.  
doi:10.4103/0972-2327.74248
- Mumma, G. H., & Fluck J. (2016). How valid is your case formulation? Empirically testing your cognitive behavioural case formulation for tailored treatment. *The Cognitive Behaviour Therapist*, 9(e12), 1-25.
- Mura, G., & Carta, M. G. (2013). Physical activity in depressed elderly: A systematic review. *Clinical Practice & Epidemiology in Mental Health*, 9, 125-135.

Mura, G., Mora, M. F., Patten, S. B., & Carta, M. G. (2014). Exercise an add-on for the treatment of major depressive disorder: A systematic review. *CNS Spectrums*, 19, 496-508.

Mustaffa, M. S., & Alkaff, S. O. A. (2011). Depression among elderly. International Conference on Social Science and Humanity. *IPEDR*, 5, V2-420-423.

Mykletun, A., Bjørkeset, O., Øverland, S., Prince, M., Dewey, M., & Stewart, R. (2009). Levels of anxiety and depression as predictors of mortality: The HUNT study. *British Journal of Psychiatry*, 195(2), 118-125. doi:10.1192/bjp.bp.108.054866

National Institute for Clinical Excellence. (2004). *Depression: Management of depression in primary and secondary care*. London: Author.

National Institute for Clinical Excellence. (2009). *Depression: Management of depression in primary and secondary care*. London: Author.

National Institute of Health. (2016). *NIH senior health: Depression*. USA: NIH Publication.

National Institute of Mental Health. (2011). *Depression*. U.S.: NIH Plublication.

National Institute of Mental Health. (2013). *Depression: What you need to know*. USA: NIH Publication.

National Institute of Mental Health. (2015). *Older adults and depression*. USA: NIH Publication.

National Statistical Office, Ministry of Information and Communication Technology of Thailand. (2012). *The main result survey of working of Thai population 2011*. Retrieved from <http://service.nso.go.th/nso/nsopublish/download/files/lfsSum54.pdf>

Neenan, M., & Dryden, W. (2010). *An overview of cognitive therapy*. Retrieved from [http://www.sagepub.com/upm-data/36593\\_01\\_Neenan\\_&\\_Dryden\\_\(CT\)\\_2e\\_CH\\_01.pdf](http://www.sagepub.com/upm-data/36593_01_Neenan_&_Dryden_(CT)_2e_CH_01.pdf)

Nierenberg, C. (2014). *7 way women and men experience depression differently*. Retrieved from [https://www.huffingtonpost.com/2014/04/28/depression-differences-in-women-and-men\\_n\\_5225719.html](https://www.huffingtonpost.com/2014/04/28/depression-differences-in-women-and-men_n_5225719.html)

Newsma, J. A., Trivedi, R. B., McDuffie, J., Kronish, I., Benjamin, D., & Williams, J. W. (2012). Brief psychotherapy for depression: a systematic review and meta-analysis. *International Journal of Psychiatry in Medicine*, 43(2), 129-151.

Nilmanut, S., Kuptniratsaikul, V., Pekuman, P., & Tosayanonda, O. (1997). The study of The Center for Epidemiologic Studies-Depression Scale (CES-D) in Thai people in Siriraj Hospital. *Journal of Thai Rehabilitation*, 6(3), 25-30.

Nukariya, K. (2001). Psychotherapy and psycho-education of depression. *Japan Medical Association Journal*, 44(5), 230-234.

Office of the National Economic and Social Development Board. (2010). *Population projections for Thailand, 2010-2040*. Retrieved from <http://fopdev.or.th/en/situation-of-the-thai-elderly-population-situations/>

Office of the National Economic and Social Development Broad. (2011). *Summary of the Eleventh National Economic and Social Development Plan (2012-2016)*. Retrieved from [http://www.nesdb.go.th/Portals/0/news/annual\\_meet/54/book/Executive%20Summary%20of%2011th%20Plan.pdf](http://www.nesdb.go.th/Portals/0/news/annual_meet/54/book/Executive%20Summary%20of%2011th%20Plan.pdf)

Palazzolo, J. (2015). Cognitive-behavioral therapy for depression and anxiety in the elderly. *Annals of Depression and Anxiety*, 2(6), 1-4.

Pasterfield, M., Bailey, D., Hems, D., McMillan, D., Richards, D., & Gilbody, S. (2014). Adapting manualized behavioural activation treatment for older adults with depression. *The Cognitive Behavior Therapist*, 7(e5), 1-11.

- Patten, S. B., Wang, J. L., Williams, J. V. A., Currie, S., Beck, C. A., Maxwell, C. J., & El-Guebaly, N. (2006). Descriptive epidemiology of major depression in Canada. *Canadian Journal of Psychiatry*, 51(2), 84- 90.
- Peltzer, K., & Phaswana-Mafuya, N. (2013). Depression and associated factors in older adults in South Africa. *Global Health Action*, 6. doi:10.3402/gha.v6i0.18871
- Pence, B. W., O'Donnell, J. K., & Gaynes, B. N. (2012). The depression treatment cascade in primary care: A public health perspective. *Current Psychiatry Reports*, 14(4), 328-335.
- Persons, J. B., & Tompkins, M. A. (2007). Cognitive-behavioral case formulation. In T. D. Eells (Ed.), *Handbook of psychotherapy case formulation* (2nd ed.). New York: Guilford Press.
- Phelan, E., Williams, B., Meeker, K., Bonn, K., Frederick, J., LoGerfo, J., & Snowden, M. (2010). A study of diagnostic accuracy of the PHQ-9 in primary care elderly. *BMC Family Practice*, 11, 63.
- Piccinelli, M., & Wilkinson, G. (2000). Gender differences in depression: Critical review. *The British Journal of Psychiatry*, 177, 486-492.
- Pilgrim, D., & Bentall, R. (1999). The medicalisation of misery: A critical realist analysis of the concept of depression. *Journal of Mental Health*, 8(3), 261-274.
- Pinquart, M., & Sorensen, S. (2016). *How effective are psychotherapeutic and other psychosocial interventions with older adults?* Center for Reviews and Dissemination, National Institute for Health Research, University of York. Retrieved from <http://www.crd.york.ac.uk/CRDWeb/ShowRecord.asp?ID=12001005478>
- Pirkis, J., Pfaff, J., Williamson, M., Tyson, O., Stocks, N., Goldney, R., . . . Almeida, O. P. (2009). The community prevalence of depression in older Australian. *Journal of Affective Disorder*, 115(1-2), 54-61. doi:10.1016/j.jad.2008.08.014

Polit, D. F., & Beck, C. T. (2004). *Essentials of nursing research* (6th ed.). Philadelphia: Lippincott Williams & Wilkins.

Polit, D. F., & Beck, C. T. (2012). *Nursing research: Generating and assessing evidence for nursing practice* (9th ed.). Philadelphia: Lippincott Williams & Wilkins.

Powell, V. B., Abreu, N., de Oliveira, I. R., & Sudak, D. (2008). Cognitive-behavioral therapy for depression. *Revista Brasileira de Psiquiatria*, 30(Suppl. 2), s73-s80.

Prasrimahaphodi Psychiatric Hospital. (2012). *The annual report 2012*. Retrieved from <http://www.thaidepression.com>

Prasrimahaphodi Psychiatric Hospital. (2015). *Handbook manual of cognitive-behavioral therapy for patient with major depressive disorder* (2nd ed.). Ubonratchathani: Author.

Radloff, L. S. (1977). The CES-D Scale: A self-report depression scale for research in the general population. *Applied Psychological Measurement*, 1(3), 385-401.

Richards, D. A., Lovell, K., Gilbody, S., Gask, L., Torgerson, D., Barkham, M., . . . Richardson, R. (2007). Collaborative care for depression in UK primary care: A randomized controlled trial. *Psychological Medicine*, 38(2), 279-287.  
doi:10.1017/S0033291707001365

Rodda, J., Walker, Z., & Carter, J. (2011). Clinical review: Depression in older adults. *The BMJ*, 343, 683-687.

Rodriguez, M. R., Nuevo, R., Chatterji, S., & Ayuso-Mateos, J. L. (2012). Definitions and factors associated with subthreshold depressive conditions: A systematic review. *BMC Psychiatry*, 12, 181. doi:10.1186/1471-244X-12-181

Rungreangkulkij, S., Chirawatkul, S., Kongsuk, T., Sukavaha, S., Leejongpermpoon, J., & Sutatho, Y. (2012). Sex or gender leading to a high risk of depressive disorder in women. *Journal of Psychiatric Association of Thailand*, 57(1), 61-74.

- Rungreangkulkij, S., Juntachum, W., Chirawatkul, S., & Charatsingha, A. (2010). Gender difference: Sociodemographic risk factors of depression. *Journal of Psychiatric Association of Thailand*, 55(4), 337-346.
- Seedat, S., Scott, K. M., Angermeyer, M. C., Berglund, P., Bromet, E. J., Brugha, T. S., . . . Kessler, R. C. (2009). Cross-national associations between gender and mental disorders in the WHO World Mental Health Surveys. *Archives of General Psychiatry*, 66(7), 785-795.
- Serfaty, M. A., Haworth, D., Blanchard, M., Buszewicz, M., Murad, S., & King, M. (2009). Clinical effectiveness of individual cognitive behavioral therapy for depressed older people in primary care: A randomized controlled trial. *Archives of General Psychiatry*, 66(12), 1332-1340.
- Sheets, D. J., & Gallagher, E. M. (2013). Aging in Canada: State of the art and science. *The Gerontologist*, 53(1), 1-8. doi:10.1093/geront/gns150
- Sheikhjits, J. I., & Yesavage, J. A. (1986). Geriatric Depression Scale (GDS): Recent evidence and development of a short version. In T. L. Brink (Ed.), *Clinical gerontology: A Guide to Assessment and Intervention* (pp. 165-173). New York: The Haworth Press.
- Sin, M.-K., Choe, M.-A., Kim, J., Chae, Y.-R., & Jeon, M.-Y. (2010). Depressive symptoms in community-dwelling elderly korean immigrants and elderly Koreans. *Research in Gerontological Nursing*, 3(4), 262-269.
- Siri, S. (2013). Screening test for dementia in community. *Journal of Public Health Nursing*, 27(1), 115-128.
- Sivrioglu, E. Y., Sivrioglu, K., Ertan, T., Ertan, F. S., Cankurtaran, E., Aki, O., . . . Kirli, S. (2009). Reliability and validity of the Geriatric Depression Scale in detection of poststroke minor depression. *Journal of Clinical and Experimental Neuropsychology*, 31(8), 999-1006.

- Smith, M., & Jaffe, J. (2016). *Depression in women*. Retrieved from <http://www.helpguide.org/articles/depression/depression-in-women.htm>
- Snowden, M., Steinman, L., & Frederick, J. (2008). Treating depression in older adults: Challenges to implementing the recommendations of an expert panel. *Preventing Chronic Disease, 5*(1), 1-7.
- Snowdon, J. (2002). How high is the prevalence of depression in old age? *Revista Brasileira de Psiquiatria, 24*, 42-47.
- Songprakul, W., & McCann, T.V. (2012). Evaluation of a bibliotherapy manual for reducing psychological distress in people with depression: A randomized controlled trial. *Journal of Advanced Nursing, 68*(12), 2674-2684.
- Spangler, D. L., Simon, A. D., Monroc, S. M., & Thase, M. T. (1996). Gender differences in cognitive diathesis-stress domain match: Implications for differential pathways to depression. *Journal of Abnormal Psychology, 105*(4), 653- 657.
- Srisurapanont, M., Pityaratstian, N., Kittiranapaiboon, P., & Charatsigha, A. (2009). Cognitive-behavioral therapy for the treatment of major depressive disorder: Workforce development and service evaluation in Thailand. *Journal of Psychiatric Association of Thailand, 54*(4), 367-374.
- St. John, P. H., Blandford, A. A., & Strain, L. A. (2009). Does a rural residence predict the development of depressive symptoms in older adults? *Canadian Journal of Rural Medicine, 14*(4), 150-156.
- Stage, K. B., Bech, P., Kragh-Sorensen, P., Nair, N. P. V., & Katona, C. (2001). Differences in symptomatology and diagnostic profile in younger and elderly depressed inpatients. *Journal of Affective Disorders, 64*(2-3), 239-248.

- Stanley, M. A., Beck, J. G., Novy, D. M., Averill, P. M., Swann, A. C., Diefenbach, G. J., & Hopko, D. R. (2003) Cognitive-behavioural treatment of late-life generalized anxiety disorder. *Journal of Consulting and Clinical Psychology*, 71(2), 309-319.
- Steel, Z., Marnane, C., Iranpour, C., Chey, T., Jackson, J. W., Patel, V., & Silove, D. (2014). The global prevalence of common mental disorders: A systematic review and meta-analysis 1980-2013. *International Journal of Epidemiology*, 43(2), 476-493. doi:10.1093/ije/dyu038
- Steinman, L. E., Frederick, J. T., Prohaska, T., Satariano, W. A., Dornberg-Lee, S., Fisher, R., . . . Snowden, M. (2007). Recommendations for treating depression in community-based older adults. *American Journal of Preventive Medicine*, 33(3), 175-181.
- Sturmey, P., & McMurran, M. (Eds.).(2011). *Forensic case formulation*. Chichester, UK: Wiley-Blackwell.
- Sugawara, N., Yasui-Furukori, N., Sasaki, G., Umeda, T., Takahashi, I., Danjo, K., . . . Nakaji, S. (2012). Relationships between Suicidal Ideation and the dimensions of depressive symptoms among middle-aged population in Japan. *Journal of Affective Disorders*, 136(3), 819-823.
- Suttajit, S., Punpuing, S., Jirapramukpitak, T., Tangchonlatip, K., Darawuttimaprakorn, N., Stewart, R., Abas, M. A. (2010). Impairment, disability, social support and depression among older parents in rural Thailand. *Psychological Medicine*, 40(10), 1711-1721. doi:10.1017/S003329170999208X
- Thew, G. R., & Krohnert, N. (2015). Formulation as intervention: Case report and client experience for formulating in therapy *The Cognitive Behaviour Therapist*, 8(e25), 1-14.

- Thompson, L.W., Coon, D.W., Gallagher-Thompson, D., Sommer, B. R., & Koin, D. (2016). Comparison of desipramine and cognitive/behavioral therapy in the treatment of older outpatients with mild to moderate depression. *American Journal of Geriatric Psychiatry*, 9(3), 225-240.
- Thongtang, O., Sukhatunga, K., Ngamthipwatthana, T., Chulakadabba, S., Vuthiganond, S., Pooviboonsuk, P., . . . Phattharayuttawat, S. (2002). Prevalence and incidence of depression in the Thais elderly. *Journal of the Medical Association of Thailand*, 85(5), 540-544.
- Tilyard, M. (2011). *Best practice, Special edition: Assessment and management of depression in older adults*. New Zealand: Bpac.
- Tombaugh, T. N., & McIntyre, N. J. (1992). The mini-mental state examination: A comprehensive review. *Journal of the American Geriatrics Society*, 40(9), 922-935.
- Trangle, M., Gursky, J., Haight, R., Hardwig, J., Hinnenkamp, T., Kessler, . . . Myszkowski, M. (2016). Health care guideline depression in primary care. Institute for Clinical Systems Improvement. Retrieved from [https://www.icsi.org/\\_asset/fnhdm3/depr-interactive0512b.pdf](https://www.icsi.org/_asset/fnhdm3/depr-interactive0512b.pdf)
- United Nations Population Fund. (2011). *Impact of demographic change in Thailand*. Bangkok: United Nations Population Fund Country Office in Thailand.
- Unutzer, J. (2007). Late-life depression. *New England Journal of Medicine*, 357(22), 2269-2276. doi:10.1056/NEJMcp073754
- Vaccarino, V., Kasl, S. V., Abramson, J., & Krumholz, H. M. (2001). Depressive symptoms and risk of functional decline and death in patients with heart failure. *Journal of American College of Cardiology*, 38(1), 199-205.

- Valiengo, L. C., Stella, F., & Forlenza, O. V. (2016). Mood disorders in the elderly: Prevalence, functional impact, and management challenges. *Neuropsychiatric Disease and Treatment*, 12, 2105-2114.
- van der Wurff, F. B., Beekman, A. T. F., Dijkshoorn, H., Spijker, J. A., Smits, C. H. M., Stek, M. L., & Werhoeff, A. (2004). Prevalence and risk-factors for depression in elderly Turkish and Moroccan migrants in the Netherlands. *Journal of Affective Disorders*, 83, 33-41.
- Waller, R., Trepka, C., Collerton, D., & Hawkins, J. (2010). Addressing spirituality in CBT. *The Cognitive Behavior Therapist*, 3, 95-106.
- Wang, J.-J. (2004). The Comparative effectiveness among institutionalized and non-institutionalized elderly people in Taiwan of reminiscence therapy as a psychological measure. *Journal of Nursing Research*, 12(3), 237-244.
- Wang, S., & Blazer, D. G. (2015). Depression and cognition in the elderly. *Annual Review of Clinical Psychology*, 11, 331-360.
- Wang, W., Bain, Q., Zhao, Y., Li, X., Wang, W., Du, J., . . . Zhao, M. (2014). Reliability and validity of the Chinese version of the Patient Health Questionnaire (PHQ-9) in the general population. *General Hospital Psychiatry*, 36, 539-544.
- Wang, Y.-P., & Gorenstein, C. (2013). Psychometric properties of the Beck Depression Inventory-II: A comprehensive review. *Revista Brasileira de Psiquiatria*, 35(4), 416-431.
- Weiss, B. D., Mohler, M. J., & Fain, M. J. (2011). Geriatrics in Taiwan: What is the solution? *Journal of Clinical Gerontology & Geriatrics*, 2, 93-95.
- Westbrook, K. (2010). *Theory, development and status of CBT*. Retrieved from [http://www.sagepub.com/upm-data/39347\\_978\\_1\\_84860\\_687\\_6.pdf](http://www.sagepub.com/upm-data/39347_978_1_84860_687_6.pdf)
- Whitbourne, S. K. (2001). *Adult development & aging: Biopsychosocial perspective*. New York: John Wiley& Sons.

Whyte, E. M., & Rovner, B. (2006). Depression in late-life: shifting the paradigm from treatment to prevention. *International Journal of Geriatric Psychiatry*, 21(8), 746-751.

Williams, C., & Garland, A. (2002). A cognitive-behavioural therapy assessment model for use in everyday clinical practice. *Advances in Psychiatric Treatment*, 8, 172-179.

Wongpoom, T., Sukyimng,C., & Udomsubpayakul, U. (2011). Prevalence of depression among the elderly in Chiang Mai Province. *Journal of the Psychiatric Association of Thailand*, 56(2), 103-116.

Worapongsathorn, T., Pandee, W., & Trimchaisri, S.(1990). Validity of the Center for Epidemiologic Studies Depression Scale (CES-D). *Journal of Clinical Psychologist*, 21, 25-45.

World Health Organization. (2008). *The global burden of disease: 2004 update*. Geneva: Author.

World Health Organization. (2010). *Mental health development: Targeting people with mental health conditions an vulnerable group*. Mental health and poverty project.

World Health Organization. (2012). *Depression: A global crisis world mental health day, October 10, 2012*. Geneva: World Federation for Mental Health, World Health Organization.

World Health Organization. (2016a). *Definition of an older or elderly person*. Retrieved from <http://www.who.int/healthinfo/survey/ageingdefolder/en/>

World Health Organization. (2016b). *Depression*. Retrieved from <http://www.who.int/mediacentre/factsheets/fs369/en/>

Wright, J. H. (2006). Cognitive behavior therapy: Basics principles and recent advances. *The Focus*, 4(2), 173-178.

- Wuthrich, V. M., & Rapee, R. M. (2013). Randomised controlled trial of group cognitive behavioural therapy for comorbid anxiety and depression in older adults. *Behaviour Research and Therapy*, 51(12), 779-786.
- Wuthrich, V. M., Rapee, R. M., Kangas, M., & Perini, S. (2016). Randomized controlled trial of group cognitive behavioral therapy compared to a discussion group for co-morbid anxiety and depression in older adults. *Psychological Medicine*, 46(4), 785-795.
- Xu, J., Wang, J., Wimo, A., & Qui, C. (2016). The economic burden of mental health disorders in China, 2005-2013: Implications for health policy. *BMC Psychiatry*, 16, 1-9.
- Yesavage, J. M., Brink, T. L., Rose, T. L., & Lum, O. (1983). Development and validation of a Geriatric Depression Screening Scale: A Preliminary report. *Journal of Psychiatric Research*, 17(1), 37-49.
- Yu, J., Li, J., Cuijpers, P., Wu, S., & Wu, Z. (2012). Prevalence and correlates of depressive symptoms in Chinese older adults: A population-based study. *International journal of Geriatric Psychiatry*, 27, 305-312.
- Zimmerman, M. (2011). *Tools for depression: Standardized rating scales*. Retrieved from <http://www.medscape.org/viewarticle/749921>
- Zuithoff, N. P. A., Vergouwe, Y., King, M., Nazareth, I., an Wezep, M. J., Moons, K. G. M., . . . Geerlings, M. I. (2010). The Patient Questionnaire-9 for detection of major depressive disorder in primary care: Consequences of current threshold in a crossectional study. *BMC Family Practice*, 11, 98. doi:10.1186/1471-2296-11-98