

## REFERENCES

- Abalain-Colloc MLC, Chastel JG, Tully JM, Bove RF, Whitcomb B, Gilo et al. Spiroplasma sabaudiense sp. nov. from mosquitos collected in France. *Int J Syst Bacteriol* 1987;37:260-5.
- Abraham EG, Jacobs-Lorena M. Mosquito midgut barriers to malaria parasite development. *Insect Biochem Mol Biol* 2004;34:667-71.
- Akaki M, Dvorak JA. A chemotactic response facilitates mosquito salivary gland infection by malaria sporozoites. *J Exp Biol* 2005;208:3211-18.
- Akhouayri IG, Habtewold T, Christophides GK. Melanotic Pathology and Vertical Transmission of the Gut Commensal *Elizabethkingia meningoseptica* in the Major Malaria Vector *Anopheles gambiae*. *PLoS ONE* 2013;8(10):e77619.
- Alvarenga PH, Francischetti IM, Calvo E, Sa-Nunes A, Ribeiro JM, Andersen JF. The function and three-dimensional structure of a thromboxane A<sub>2</sub>/cysteinyl leukotriene-binding protein from the saliva of a mosquito vector of the malaria parasite. *PLoS Biol* 2010;8: e1000547.
- Alves CR, Albuquerque-Cunha JM, Mello CB, Garcia ES, Nogueira NF, Bourguignon SC, et al. *Trypanosoma cruzi*: attachment to perimicrovillar membrane glycoproteins of *Rhodnius prolixus*. *Exp Parasitol* 2007;116:44-52.
- Amino R, Thibierge S, Martin B, Celli S, Shorte S, Frischknecht F, et al. Quantitative imaging of *Plasmodium* transmission from mosquito to mammal. *Nature Med*. 2006;12:220-4.
- Andrade BB, Teixeira CR, Barral A, Barral-Netto M. Haematophagous arthropod saliva and host defense system: a tale of tear and blood. *An Acad Bras Cienc* 2005;77:665-93.

Apte-Deshpande A, Paingankar M, Gokhale MD, Deobagkar DN. *Serratia odorifera* a Midgut Inhabitant of *Aedes aegypti* Mosquito Enhances Its Susceptibility to Dengue-2 Virus. PLoS ONE 2012;7:e40401.

Arca` B, Lombardo F, Lanfrancotti A, Spanos L, Veneri M, Louis C, et al. A cluster of four D7-related genes is expressed in the salivary glands of the African malaria vector *Anopheles gambiae*. Insect Mol Biol 2002;11:47-55.

Assumpcao TC, Ma D, Schwarz A, Reiter K, Santana JM, Andersen JF, et al. Salivary antigen-5/CAP family members are Cu<sup>2+</sup>-dependent antioxidant enzymes that scavenge O<sub>2</sub>(-). and inhibit collagen-induced platelet aggregation and neutrophil oxidative burst. J Biol Chem 2013;288:14341-61.

Atmosoedjono S, Partono F, Dennis DT, Purnomo : *Anopheles barbirostris* (Diptera: Culicidae) as a vector of the Timor filaria on Flores Island: preliminary observations. J Med Entomol 1977;13:611-3.

Baker GC, Smith JJ, Cowan DA. Review and re-analysis of domain-specific 16S primers. J microbial methods. 2003;55:541-55.

Barrow PM, McIver SB, Wright KA. Salivary glands of female *Culex pipiens*: morphological changes associated with maturation and blood-feeding. Can Ent 1975;107:1153-60.

Baton LA, Ranford-Cartwright LC. How do malaria ookinetes cross the mosquito midgut wall? Trends Parasitol 2005a;21:22-8.

Beard CB, Cordon-Rosales C, Durvasula RV. Bacterial symbionts of the triatominae and their potential use in control of Chagas disease transmission. Annu Rev Entomol 2002;47:123-41.

Beckett EB. Development and ageing of the salivary glands of adult female *Aedes aegypti* (L.) and Aedes mosquito *Aedes togoi* (Theobald) mosquitoes (Diptera: Culicidae). Int J Insect Morphol Embryol 1990;19:277-90.

Beier JC. Malaria parasite development in mosquitoes. Annu Rev Entomol 1998;43:519-43.

Billingsley PF. Blood digestion in the mosquito, *Anopheles stephensi* Liston (Diptera: Culicidae): partial characterization and post-feeding activity of midgut aminopeptidases. Arch Insect Biochem Physiol 1990;15:149-63.

Billingsley PF, Rudin WTJ. The role of the mosquito peri-trophic membrane in blood meal digestion and infectivity of *Plasmodium* species. J Parasitol 1992;78:430-40.

Boissière A, Tchioffo MT, Bachar D, Abate L, Marie A, Nsango SE, et al. Midgut microbiota of the malaria mosquito vector *Anopheles gambiae* and interactions with *Plasmodium falciparum* infection. PLoS Pathog 2012;8:e1002742.

Boisson B, Jacques JC, Choumet V, Martin E, Xu J, Vernick K, et al. Gene silencing in mosquito salivary glands by RNAi. FEBS letters. 2006;580:1988-1992.

Briones AM, Shililu J, Githure J, Novak R, Raskin L. *Thorsellia anophelis* is the dominant bacterium in a Kenyan population of adult *Anopheles gambiae* mosquitoes. The ISME journal 2008;2:74-82.

Calvo E, Andersen J, Francischetti IM, deL Capurro M, deBianchi AG, James AA, et al. The transcriptome of adult female *Anopheles darlingi* salivary glands. Insect Biochem Mol Biol 2004;13:73-88.

Calvo E, Dao A, Pham VM, Ribeiro JMC. An insight into sialome of *Anopheles funestus* reveals an emerging pattern in anopheline salivary protein families. Insect Biochem Mol Biol 2006;37:164-75.

Calvo E, Pham VM, Marinotti O, Andersen JF, Ribeiro JM. The salivary gland transcriptome of the neotropical malaria vector *Anopheles darlingi* reveals accelerated evolution of genes relevant to hematophagy. BMC Genomics 2009;10:57-66.

Carlson J. Genetic manipulation of mosquitoes an approach to controlling disease. Trends Biotechnol 1996;1:447-8.

Caroci AS, Calvo E, Ribolla PE, de Bianchi AG, Marinotti O. Two digestive trypsins occur in three species of neotropical anophelines. J Med Entomol 2003;40:991-5.

Cázares-Raga FE, Chávez-Munguía B, González-Calixto C, Ochoa-Franco AP, Gawinowicz MA, Rodríguez MH, et al. Morphological and proteomic characterization of midgut of the malaria vector *Anopheles albimanus* at early time after a blood feeding. J Proteomics 2014;111:100-12.

Champagne DE, Ribeiro JMC. Sialokinin I and II: vasodilatory tachykinins from the yellow fever mosquito *Aedes aegypti*. Proc Natl Acad Sci 1994;91:138-42.

Chandel K, Mendki MJ, Parikh RY, Kulkarni G, Tikar SN, Sukumaran D, et al. Midgut microbial community of *Culex quinquefasciatus* mosquito populations from India. PLoS ONE 2013;8:e80453.

Choochote W and Saeung A. Systematic techniques for the recognition of Anopheles species complexes, in: S. Manguin (Ed.). Anopheles mosquitoes-New insights into malaria vectors. InTech 2013;57-79.

Choumet V, Attout T, Chartier L, Khun H, Sautereau J, Robbe-Vincent A, et al. Visualizing non infectious and infectious *Anopheles gambiae* blood feedings in naive and saliva-immunized mice. PLoS ONE. 2012;7:e50464.

Clements AN : “Adult digestion.” In The biology of mosquitoes. Chapman & Hall, London, New York. 1992, pp.272-91.

Conte JE. A novel approach to preventing insect-borne diseases. N Engl J Med 1997;337:785-6.

Cotama S, Dekumyoy P, Samung Y, Lek-Uthai U. Salivary glands proteins expression of *Anopheles dirus* A fed on *Plasmodium vivax* and *Plasmodium falciparum* infected human blood. J Parasitol Res 2013;2013:535267.

Crotti E, Damiani C, Pajoro M, Gonella E, Rizzi A, Ricci I, et al. *Asaia*, a versatile acetic acid bacterial symbiont, capable of cross-colonizing insects of phylogenetically distant genera and orders. *Environ Microbiol* 2009;11: 3252-3264. doi:10.1111/j.1462-2920.2009.02048.x.

da Cunha Sais T, de Moraes RM, Ribolla PE, de Bianchi AG, Marinotti O, Bijovsky AT. Morphological aspects of *Culex quinquefasciatus* salivary glands. *Arthropod Struct Dev* 2003;32:219-26.

Damiani C, Ricci I, Crotti E, Rossi P, Rizzi A, Scuppa P, et al. Mosquito-Bacteria symbiosis: The case of *Anopheles gambiae* and *Asaia*. *Microb Ecol* 2010;60:644-54.

Das S, Radtke A, Choi YJ, Mendes AM, Valenzuela JG, Dimopoulos G. Transcriptomic and functional analysis of the *Anopheles gambiae* salivary gland in relation to blood feeding. *BMC Genomics* 2010;11:566.

Delong EF 1992. Archaea in coastal marine environments. *Proc Natl Acad Sci USA* 89:5685-9.

Demaio J, Pumpuni CB, Kent M, Beier JC. The midgut bacterial flora of wild *Aedes triseriatus*, *Culex pipiens* and *Psorophora columbiae* mosquitoes. *Am J Trop Med Hyg* 1996;54:219-23.

Devenport M, Fujioka H, Jacobs-Lorena M. Storage and secretion of the peritrophic matrix protein Ag-Aper1 and trypsin in the midgut of *Anopheles gambiae*. *Insect Mol Biol* 2004;13:349-58.

Dhar R, Kumar N. Role of mosquito salivary glands. *Curr Sci* 2003;85:1308-13.

Dong Y, Manfredini F, Dimopoulos G. Implication of the mosquito midgut microbiota in the defense against malaria parasites. *PLoS Pathog* 2009;5:e1000423.

Favia G, Ricci I, Damiani C, Raddadi N, Crotti E, Marzorati M, et al. Bacteria of the genus *Asaia* stably associate with *Anopheles stephensi*, an Asian malarial mosquito vector. *Proc Natl Acad Sci USA* 2007;104:9047-51.

- Ferguson M J, Micks DW. Microorganisms associated with mosquitoes: I. Bacteria isolated from the midgut of adult *Culex fatigans* Wiedemann. *J Insect Pathol* 1961;3:112-9.
- Ferreira M, Lima J, Rocha R, Pimenta P. Morphological Aspect of the Midgut of *Anopheles aquasalis* (Curry, 1932) Insecta: Diptera. *Int J Morphol* 2008;26:577-82.
- Fialho MCQ, Zanuncio JC, Neves CA, Ramalho FS, Serrão JE. Ultrastructure of the Digestive Cells in the Midgut of the Predator *Brontocoris tabidus* (Heteroptera: Pentatomidae) after Different Feeding Periods on Prey and Plants. *Ann Entomol Soc Am* 2009;102:119-27.
- Francischetti IM, Valenzuela JG, Ribeiro JMC. Anophelin: kinetics and mechanism of thrombin inhibition. *Biochemistry* 1999;38:16678-85.
- Frischknecht F, Baldacci P, Martin B, Zimmer C, Thiberge S, Olivo-Marin JC, et al. Imaging movement of malaria parasites during transmission by *Anopheles* mosquitoes. *Cell Microbiol* 2004;6:687-94.
- Frischknecht F, Martin B, Thiery I, Bourgouin C, Menard R. Using green fluorescent malaria parasites to screen for permissive vector mosquitoes. *Malar J* 2006; 5:23.
- Fouda MA, Hassan MI, Al-Daly AG, Hammad KM. Effect of midgut bacteria of *Culex pipiens* L. on digestion and reproduction. *J Egypt Soc Parasitol* 2001;31:767-80.
- Ghosh A, Edwards MJ, Jacobs-Lorena M. The journey of the malaria parasite in the mosquito: hopes for the new century. *Parasitol Today* 2000;16:196-201.
- Girard YA, Popov V, Wen J, Han V, Higgs S. Ultrastructural study of West Nile virus pathogenesis in *Culex pipiens quinquefasciatus* (Diptera: Culicidae). *J Med Entomol* 2005;42:429-44.

Godoy RS, Fernandes KM, Martins GF. Midgut of the non-hematophagous mosquito *Toxorhynchites theobaldi* (Diptera, Culicidae). Sci rep 2015;5.

Gonzalez-Ceron L, Santillan F, Rodriguez MH, Mendez D, Hernandez-Avila JE. Bacteria in midguts of field-collected *Anopheles albimanus* block *Plasmodium vivax* sporogonic development. J Med Entomol 2003;40:371-4.

Gonzalez P, Marin C, Rodriguez-Gonzalez I, Illana A, Mateo H, Longoni SS, et al. Diterpenoid alkaloid derivatives as potential chemotherapeutic agents in American trypanosomiasis. Pharmacology 2006;76:123-8.

Gross TL, Myles KM, Adelman ZN. Identification and characterization of heat shock 70 genes in *Aedes aegypti* (Diptera: Culicidae). J Med Entomol 2009;46:496-504.

Gouveia C, Asensi MD, Zahner V, Rangel EF, Oliveria S. Study on the Bacterial midgut microbiota associated to different brazilian populations of *Lutzomyia longipalpis* (Lutz & Neiva) (Diptera: Psychodidae). Neotrop Entomol 2008;37:597-601.

Gupta L, Kumar S, Han YS, Pimenta PFP, Barillas-Mury C. Midgut epithelial responses of different mosquito-*Plasmodium* combinations: The actin cone zipper repair mechanism in *Aedes aegypti*. Proc Natl Acad Sci USA 2005;102:4010-5.

Gusmaõ DS, Santos AV, Marini DC, Russo ES, Peixoto AMD. First isolation of microorganisms from the gut diverticulum of *Aedes aegypti* (Diptera: Culicidae): new perspectives for an insect-bacteria association. Mem Inst Oswaldo Cruz 2007;102:919-24.

Gusmaõ DS, Santos AV, Marini DC, Bacci M, Berbert-Molina MA, Lemos FJ. Culture-dependent and culture-independent characterization of microorganisms associated with *Aedes aegypti* (Diptera: Culicidae) (L.) and dynamics of bacterial colonization in the midgut. Acta Tropica 2010;115:275-81.

Hagedorn HH. The endocrinology of the adult female mosquito. Adv. Dis Vect Res 1994;10:109-48.

- Han YS, Thompson J, Kafatos FC, Barillas-Mury C. Molecular interactions between *Anopheles stephensi* midgut cells and *Plasmodium berghei*: the time bomb theory of ookinete invasion of mosquitoes. EMBO J 2000;19:6030-40.
- Hecker H, Brun R, Reinhardt C, Burri PH. Morphometric analysis of the midgut of female *Aedes aegypti* (L.) (Insecta Diptera) under various physiological conditions. Cell Tissue Res 1974;152:31-49.
- Hecker H. Structure and function of midgut epithelial cells in culicidae mosquitoes (insecta, diptera) Cell Tissue Res 1977;184: 321.doi:10.1007/BF00219894.
- Huber M, Cabib E, Miller LH. Malaria parasite chitinase and penetration of the mosquito peritrophic membrane. Proc Natl Acad Sci USA 1991; 88:2807-10.
- Ikeda H, Omura S. Control of avermectin biosynthesis in *Streptomyces avermitilis* for the selective production of a useful component. J Antibiot 1995;48:549-62.
- Intakhan N, Jariyapan N, Sor-Suhan S, Phattanawiboon B, Taai K, Chanmol W, et al. Exsheathment and midgut invasion of nocturnally subperiodic *Brugia malayi* microfilariae in a refractory vector, *Aedes aegypti* (Thailand strain). Parasitol Res 2014;113:4141-9.
- Isawa H, Orito Y, Iwanaga S, Jingushi N, Morita A, Chinzei Y, et al. Identification and characterization of a new kallikrein-kinin system inhibitor from the salivary glands of the malaria vector mosquito *Anopheles stephensi*. Insect Biochem Mol Biol 2007;37:466-77.
- James AA. Blocking malaria parasite invasion of mosquito salivary glands. J Exp Biol 2003;206:3817-21.
- Janzen HG, Wright KA. The salivary glands of *Aedes aegypti* (L.): an electron microscope study. Can J Zool 1971;49:1343-46.
- Jaresithikunchai J, Phaonakrop N, Kittisenachai S, Roytrakul S. Rapid in-gel digestion protocol for protein identification by peptide mass fingerprint. In: Proceeding of the 2nd biochemistry and molecular biology conference: biochemistry and molecular biology for regional sustainable development. 2009;May7-8 Khon Kaen, Thailand, p 29.
- Jariyapan N, Choochote W, Jitpakdi A, Harnnoi T, Siriyasatein P, Wilkinson MC, et al. Salivary gland proteins of the human malaria vector, *Anopheles dirus* B

- (Diptera: Culicidae). Revista do Instituto de Medicina Tropical de São Paulo 2007;49:5-10.
- Jariyapan N, Baimai V, Poovorawan Y, Roytrakul S, Saeung A, Thongsahuan S, et al. Analysis of female salivary gland proteins of the *Anopheles barbirostris* complex (Diptera: Culicidae) in Thailand. Parasitol Res 2010;107:509-16.
- Jariyapan N, Roytrakul S, Paemanee A, Junkum A, Saeung A, Thongsahuan S, et al. Proteomic analysis of salivary glands of female *Anopheles barbirostris* species A2 (Diptera: Culicidae) by two-dimensional gel electrophoresis and mass spectrometry. Parasitol Res 2012;111:1239-49.
- Jariyapan N, Saeung A, Intakhan N, Chanmol W, Sor-Suwan S, Phattanawiboon B, et al. Peritrophic matrix formation and *Brugia malayi* microfilaria invasion of the midgut of a susceptible vector, *Ochlerotatus togoi* (Diptera: Culicidae). Parasitol Res 2013;112:2431-440.
- Juhn J, Naeem-Ullah U, Guedes B, Majid A, Coleman J, Pimenta P, et al. Spatial mapping of gene expression in the salivary glands of the dengue vector mosquito, *Aedes aegypti*. Parasit Vectors 2011;4.
- Laemmli UK. Cleavage of Structural Proteins during the Assembly of the Head of Bacteriophage T4. Nature 1970;226:607-10.
- Lehane, M.J. Biology of Blood-Sucking Insects, Second Edition, 2005, Cambridge University Press. 321 pp.
- Lane DJ, Pace B, Olsen GJ, Stahl DA, Sogin ML, Pace NR. Rapid determination of 16S ribosomal sequences for phylogenetic analyses. Proc Natl Acad Sci USA 1985;82:6955-9.
- Lindh JM, Terenius O, Faye I. 16S rRNA gene-based identification of midgut bacteria from field-caught *Anopheles gambiae* Sensu Lato and *A. funestus* Mosquitoes reveals new species related to known insect symbionts. Appl Environ Microbiol 2005;71:7217-23.
- Lombardo F, Ronca R, Rizzo C, Mestres-Simòn M, Lanfrancotti A, Currà C, et al. The *Anopheles gambiae* salivary gland protein gSG6: An anopheline-specific protein with a blood-feeding role. Insect Biochem Mol Biol 2009;39:457-66.

- Matsuoka H, Yoshida S, Hirai M, Ishii A. A rodent malaria, *Plasmodium berghei*, is experimentally transmitted to mice by merely probing of infective mosquito, *Anopheles stephensi*. Parasitol Int. 2002;51:17-23.
- Matsuoka H, Tomita H, Hattori R, Arai M, Hirai M. Visualization of malaria parasites in the skin using the luciferase transgenic parasite, *Plasmodium berghei*. Trop Med Health 2015;43:53-61.
- Marinotti O, Calvo E, Nguyen QK, Dissanayake S, Ribeiro JMC, James AA. Genome-wide analysis of gene expression in adult *Anopheles gambiae*. Insect Mol Biol 2006;15:1-12.
- Moll RM, Romoser WS, Modrzakowski MC, Moncayo AC, Lerdthusnee K. Meconial peritrophic membranes and the fate of midgut bacteria during mosquito (Diptera: Culicidae) metamorphosis. J Med Entomol 2001;38:29-32.
- Moncayo AC, Lerdthusnee K, Leon R, Moll RM, Romoser W. Meconial peritrophic membrane structure, formation and meconial degeneration in mosquito pupae/pharate adults: histological and ultrastructural aspects. J Med Entomol 2005;42:939-44.
- Moore JE, McCalmont M, Xu J, Millar BC, Heaney N. *Asaia* sp., an unusual spoilage organism of fruit-flavored bottled water. Appl Environ Microbiol 2002;68:4130-1.
- Moreira-Ferro CK, Daffre S, James AA, Morinotti O. A lysozyme in the salivary glands of malaria vector *Anopheles darlingi*. Insect Mol Biol 1998;7:257-64.
- Moreira-Ferro CK, Morinotti O, Bijovsky AT. Morphological and biochemical analyses of the salivary glands of the malaria vector, *Anopheles darlingi*. Tissue Cell 1999;31:264-73.
- Morinotti O, James AA, Ribeiro JMC. Diet and salivation in female *Aedes aegypti* mosquitoes. J Insect Physiol 1990;36:545-8.
- Myung JM, Marshall P, Sinnis P. The *Plasmodium* circumsporozoite protein is involved in mosquito salivary gland invasion by sporozoites. Mol Biochem Parasitol 2004;133:53-9.
- Na-Bangchang K, Congpuong K. Current malaria status and distribution of drug resistance in East and Southeast Asia with special focus to Thailand. Tohoku J Exp Med 2007;211:99-113.

- Nair PMG, Choi J. Characterization and transcriptional regulation of thioredoxin reductase 1 on exposure to oxidative stress inducing environmental pollutants in *Chironomus riparius*. Comp Biochem Physiol B Biochem Mol Biol 2012;161:134-9.
- Nascimento EP, dos Santos Malafronte R, Marinotti O. Salivary gland proteins of the mosquito *Culex quinquefasciatus*. Arch Insect Biochem Physiol 2000;43:9-15.
- Ngo CT, Aujoulat F, Veas F, Jumas-Bilak E, Manguin S. Bacterial Diversity Associated with Wild Caught Anopheles Mosquitoes from Dak Nong Province, Vietnam Using Culture and DNA Fingerprint. PLoS ONE. 2015;10(3):e0118634.
- Ngwa CJ, Glöckner V, Abdelmohsen UR, Scheuermayer M, Fischer R, Hentschel U, et al. 16S rRNA gene-based identification of *Elizabethkingia meningoseptica* (Flavobacteriales: Flavobacteriaceae) as a dominant midgut bacterium of the Asian malaria vector *Anopheles stephensi* (Diptera: Culicidae) with antimicrobial activities. J Med Entomol 2013;50:404-14.
- Noriega FG, Shah DK, Wells MA. Juvenile hormone controls early trypsin gene transcription in the midgut of *Aedes aegypti*. Insect Mol Biol. 1997;6:63-6.
- Novak MG, Ribeiro JM, Hildebrand JG. 5-hydroxytryptamine in the salivary glands of adult female *Aedes aegypti* and its role in regulation of salivation. The J Exp Biol 1995;198:167-74.
- Okuda K, Caroci AS, Ribolla PEM, Bianchi AG, Bijovsky AT. Functional morphology of adult female *Culex quinquefasciatus* midgut during blood digestion. Tissue Cell Res 2002;255:1-9.
- Okuda K, Caroci AS, Ribolla P, Marinotti O, de Bianchi AG, Bijovsky AT. Morphological and enzymatic analysis of the midgut of *Anopheles darlingi* during blood digestion. J Insect Physiol 2005;51:769-76.
- Orr CWM, Hudson A, West AS. The salivary glands of *Aedes aegypti*, histological-histochemical studies. Can J Zool 1961;39:265-72.
- Pampiglione S, Majori G, Petrangeli G, Romi R. Avermectins, MK-933 and MK-936, for mosquito control. Trans R Soc Trop Med Hyg 1985;79:797-9.

- Perkins DN, Pappin DJC, Creasy DM, Cottrell JS. Probability-based protein identification by searching sequence databases using mass spectrometry data. *Electrophoresis* 1999;20:3551-67.
- Perrone JB, Spielman A. Time and site of assembly of the peritrophic membrane of the mosquito *Aedes aegypti*. *Cell Tissue Res* 1988;252:473-8.
- Phattanawiboon B, Jariyapan N, Roytrakul S, Paemanee A, Sor-suwan S, Intakhan N, et al. Morphological and protein analyses of adult female salivary glands of *Anopheles barbirostris* species A1 (Diptera: Culicidae). *Trop Biomed* 2014;31:813-27.
- Phumee A, Preativatanyou K, Kraivichain K, Thavara U, Tawatsin A, Phusup Y, et al. Morphology and protein profiles of salivary glands of filarial vector mosquito *Mansonia uniformis*; possible relation to blood feeding process. *Asian Biomed* 2011;5:353-60.
- Pidiyar VJ, Jangid K, Patole M, Shouche YS. Studies on cultured and uncultured microbiota of wild *Culex quinquefasciatus* mosquito midgut based on 16S ribosomal RNA gene analysis. *Am J Trop Med Hyg* 2004;70:597-603.
- Popova-Butler A, Dean DH. Proteomic analysis of the mosquito *Aedes aegypti* midgut brush border membrane vesicles. *J Insect Physiol* 2009;55:264-72.
- Pollard TD, Cooper JA. Actin, a central player in cell shape and movement. *Science* 2009;326:1208-12.
- Prevot GI, Laurent-Winter C, Feldmann AM, Rodhain F, Bourgouin C. Two-dimensional gel analysis of midgut proteins of *Anopheles stephensi* lines with different susceptibility to *Plasmodium falciparum* infection. *Insect Mol Biol* 1998;7: 375-83.
- Prevot GI, Laurent-Winter C, Rodhain F, Bourgouin C. Sex-specific and blood meal-induced proteins of *Anopheles gambiae* midguts: analysis by two-dimensional gel electrophoresis. *Malaria J* 2003;2: 1.

Pumpuni CB, Beier MS, Nataro JP, Guers LD, Davis JR. *Plasmodium falciparum*: inhibition of sporogonic development in *Anopheles stephensi* by gram-negative bacteria. *Exp Parasitol* 1993;77:195-9.

Pumpuni CB, DeMaio J, Kent M, Davis JR, Beier JC. Bacterial population dynamics in three anopheline species: the impact on *Plasmodium* sporogonic development. *Am J Trop Med Hyg* 1996;54:214-8.

Raikhel AS, Brown MR, Belles X. Hormonal control of reproductive processes. In: Gilbert LI, Iatrou K, Gill SS, editors. *Comprehensive molecular insect science*. Vol. 3. Elsevier; Boston: 2005. pp.433-91.

Ramirez JL, Short SM, Bahia AC, Saraiva RG, Dong Y, Kang S, et al. *Chromobacterium Csp\_P* Reduces Malaria and Dengue Infection in Vector Mosquitoes and Has Entomopathogenic and *In Vitro* Anti-pathogen Activities. *PLoS Pathog* 2014;10:e1004398.

Ramirez JL, Souza-Neto J, Torres Cosme R, Rovira J, Ortiz A, Pascale JM, et al. Reciprocal Tripartite Interactions between the *Aedes aegypti* Midgut Microbiota, Innate Immune System and Dengue Virus Influences Vector Competence. *PLoS Negl Trop Dis* 2012;6:e1561.

Rani A, Sharma A, Rajagopal R, Adak T, Bhatnagar RK. Bacterial diversity analysis of larvae and adult midgut microflora using culture-dependent and culture-independent methods in lab-reared and field-collected *Anopheles stephensi*-an Asian malarial vector. *BMC Microbiol* 2009;9: 96.

Ranson H, Hemingway J. Mosquito glutathione transferases. *Methods in enzymology*. 2005;401:226-41.

Rao KV, Chattopadhyay SK, Reddy GC. Flavonoids with mosquito larval toxicity. *J Agr Food Chem* 1990; 38:1427-30.

Reid JA: Anopheline mosquitoes of Malaya and Borneo Malaysia; 1968.

Reid JA, Harrison BA, Atmosoedjono S: Variation and vector status in *Anopheles barbirostris*. Mosq Syst 1979;11:235-51.

Ribeiro JMC, Francischetti IM. Role of arthropod saliva in blood feeding: sialome and post-sialome perspectives. Annu Rev Entomol 2003;48:73-88.

Rodrigues J, Agrawal N, Sharma A, Malhotra P, Adak T, Chauhan VS, Bhatnagar RK. Transcriptional analysis of an immune-responsive serine protease from Indian malarial vector, *Anopheles culicifacies*. BMC Molecular Biol 2007;8:33.

Rodrigues J, Sharma A, Kajla M, Agrawal N, Adak T, Bhatnagar RK. *Plasmodium* infection upregulates prophenoloxidase (AcPPO6A) in *Anopheles culicifacies*. Innate Immun 2009;1:1-8.

Rodrigues J, Brayner FA, Alves LC, Dixit R, Barillas-Mury C. Hemocyte differentiation mediates innate immune memory in *Anopheles gambiae* mosquitoes. Science 2010;329:1353-5.

Romoser WS : "The vector alimentary system." In: Beaty BJ, Marquardt WC (Eds.), Biology of Disease Vectors. University Press of Colorado, Niwot, pp.298-317, 1996.

Saboiia-Vahia L, Borges-Veloso A, Cuervo P, Junqueira M, Mesquita-Rodrigues C, Britto C, et al. Protein expression in the midgut of sugar-fed *Aedes albopictus* females. Parasit Vectors 2012;5:1-25.

Saeung A, Otsuka Y, Baimai V, Somboon P, Pitasawat B, Tuetun B, et al. Cytogenetic and molecular evidence for two species in the *Anopheles barbirostris* complex (Diptera: Culicidae) in Thailand. Parasitol Res 2007;101:1337-44.

Saeung A, Baimai V, Otsuka Y, Rattanarithikul R, Somboon P, Junkum A, et al. Molecular and cytogenetic evidence of three sibling species of the *Anopheles barbirostris* Form A (Diptera: Culicidae) in Thailand. Parasitol Res 2008;102: 499-507.

Sato Y, Montagna GN, Matuschewski K. *Plasmodium berghei* sporozoites acquire virulence and immunogenicity during mosquito hemocoel transit. Infect Immun 2014;82:1164-72.

Sawabe K, Moribayashi A. Lipid Utilization for Ovarian Development in an Autogenous Mosquito, *Culex pipiens molestus* (Diptera: Culicidae). J Med Entomol 2000;37:726-31.

Scaraffia PY, Zhang Q, Thorson K, Wysocki VH, Miesfeld RL. Differential ammonia metabolism in *Aedes aegypti* fat body and midgut tissues. J insect physiol 2010;56:1040-9.

Serrano-Pinto V, Acosta-Perez M, Luviano-Bazan D, Huritado-Sil G, Batista CVF, Martínez-Barnetche J, et al. Differential expression of proteins in the midgut of *Anopheles albimanus* with *Plasmodium berghei*. Insect Biochem Mol Biol 2010;40:752-8.

Shahabuddin M, Pimenta PF. *Plasmodium gallinaceum* preferentially invades vesicular ATPase-expressing cells in *Aedes aegypti* midgut. Proc Natl Acad Sci USA 1998;95:3385-9.

Siden-Kiamos I, Louis C. Interactions between malaria parasites and their mosquito hosts in the midgut. Insect Biochem Mol Biol 2004;34:679-85.

Sidjanski S, Vanderberg JP. Delayed migration of *Plasmodium* sporozoites from the mosquito bite site to the blood. Am J Trop Med Hyg 1997;57:426-9.

Silva JR, Mury FB, Oliveira MF, Oliveira PL, Silva CP, Dansa-Petretski M. Perimicrovillar membranes promote hemozoin formation into *Rhodnius prolixus* midgut. Insect Biochem Mol Biol 2007;37:523-31.

Sim C, Denlinger DL. Catalase and superoxide dismutase-2 enhance survival and protect ovaries during overwintering diapause in the mosquito *Culex pipiens*. J Insect physiol 2011;57:628-34.

- Siriwasatien P, Tangthongchaiwiriy K, Jariyapan N, Kaewsaitiam S, Poovorawan Y, Thavara U. Analysis of salivary gland proteins of the mosquito *Armigeres subalbatus*. Southeast Asian J Trop Med Public Health 2005;36:64-7.
- Sor-Suhan S, Jariyapan N, Roytrakul S, Paemanee A, Saeung A, Thongsahuan S, et al. Salivary gland proteome of the human malaria vector, *Anopheles campestris*-like (Diptera: Culicidae). Parasitol Res 2013;112:1065-75.
- Sor-suwan S, Jariyapan N, Roytrakul S, Paemanee A2, Phumee A3, Phattanawiboon B, et al. Identification of salivary gland proteins depleted after blood feeding in the malaria vector *Anopheles campestris*-like mosquitoes (Diptera: Culicidae). PLoS One 2014;9:1-9.
- Stark KR, James AA. Salivary gland anticoagulants in culicine and anopheline mosquitoes (Diptera:Culicidae). J Med Entomol 1996;33:645-50.
- Straif SC, Mbogo CN, Toure AM, Walker ED, Kaufman M, Toure YT, et al. Midgut bacteria in *Anopheles gambiae* and *An. funestus* (Diptera: Culicidae) from Kenya and Mali. J Med Entomol 1998;35:222-6.
- Sun D, McNicol A, James AA, Peng Z. Expression of functional recombinant mosquito salivary apyrase: a potential therapeutic platelet aggregation inhibitor. Platelets 2006;17:178-84.
- Suhan N, Wilkinson MC, Crampton JM, Bates PA. Expression of D7 and D7-related proteins in the salivary glands of the human malaria mosquito *Anopheles stephensi*. Insect Mol Biol 2002;11:223-32.
- Suwannamit S, Baimai V, Otsuka Y, Saeung A, Thongsahuan S, Tuetun B, et al. Cytogenetic and molecular evidence for an additional new species within the taxon *Anopheles barbirostris* (Diptera: Culicidae) in Thailand. Parasitol Res 2009;104:905-18.
- Taai K, Harbach RE. Systematics of the *Anopheles barbirostris* species complex (Diptera: Culicidae: Anophelinae) in Thailand. Zool J Kinnean Soc 2015;174:244-64.

Tchankouo-Nguetcheu S, Khun H, Pincet L, Roux P, Bahut M, Huerre M, et al. Differential protein modulation in midguts of *Aedes aegypti* infected with Chikungunya and Dengue 2 viruses. PloS ONE 2010;5:e13149.

Terenius O, de Oliveira CD, Pinheiro WD, Tadei WP, James AA, Marinotti O. 16S rRNA gene sequences from bacteria associated with adult *Anopheles darlingi* (Diptera: Culicidae) mosquitoes. J Med Entomol 2008;45: 172-5.

Terenius O, Lindh JM, Eriksson-Gonzales K, Bussiere L, Laugen AT, Bergquist H, et al. Midgut bacterial dynamics in *Aedes aegypti*. FEMS Microbiol Ecol 2012;80:556-65.

Thongsahuan S, Baimai V, Junkum A, Saeung A, Min GS, Joshi D, et al. Susceptibility of *Anopheles campestris*-like and *Anopheles barbirostris* species complexes to *Plasmodium falciparum* and *Plasmodium vivax* in Thailand. Mem Inst Oswaldo Cruz 2011;106:105-12.

Tucker CM, Fukami T. Environmental variability counteracts priority effects to facilitate species coexistence: evidence from nectar microbes. Proc R Soc Lond B Biol Sci 2014;281:1778.

Valenzuela JG, Francischetti IM, Pham VM, Garfield MK, Ribeiro JMC. Exploring the salivary gland transcriptome and proteome of the *Anopheles stephensi* mosquito. Insect Biochem Mol Biol 2003;33:717-32.

Vanderberg JP, Frevert U. Intravital microscopy demonstrating antibody-mediated immobilisation of *Plasmodium berghei* sporozoites injected into skin by mosquitoes. Int J Parasitol 2004;34:991-6.

Vega-Rodriguez J, Ghosh AK, Kanzok SM, Dinglasan RR, Wang S, Bongio NJ, et al. Multiple pathways for *Plasmodium* ookinete invasion of the mosquito midgut. Proc Natl Acad Sci USA 2014;111:492-500.

Villalon JM, Ghosh A, Jacobs-Lorena M. The peritrophic matrix limits the rate of digestion in adult *Anopheles stephensi* and *Aedes aegypti* mosquitoes. J Insect Physiol 2003;49:891-5.

Vlachou D, Zimmermann T, Cantera R, Janse CJ, Waters AP, Kafatos FC. Real-time, in vivo analysis of malaria ookinete locomotion and mosquito midgut invasion. *Cell Microbiol* 2004;6:671-85.

Wang Y, Gilbreath TM, Kukutla P, Yan G, Xu J. Dynamic gut microbiome across life history of the malaria mosquito *Anopheles gambiae* in Kenya. *PLoS One* 2011;6:e24767.

Wasinpiyamongkol L, Patramool S, Luplertlop N, Surasombatpattana P, Doucoure S, Mouchet F, et al. Blood-feeding and immunogenic *Aedes aegypti* saliva proteins. *Proteomics* 2010;10:1906-16.

Whitten MM, Shiao SH, Levashina EA. Mosquito midguts and malaria: cell biology, compartmentalization and immunology. *Parasite Immunol* 2006;28:121-30.

World Health Organization. World Malaria report 2014. WHO Global Malaria Programme 2014; p 238.

Wright KA. The anatomy of salivary glands of *Anopheles stephensi* Liston. *Can J Zool* 1969;47:579-87.

Yamauchi LM, Coppi A, Snounou G, Sinnis P. Plasmodium sporozoites trickle out of the injection site. *Cell Microbiol* 2007;9:1215-22.

Yoshida S, Sudo T, Niimi M, Tao L, Sun B, Kambayashi J, et al. Inhibition of collagen-induced platelet aggregation by anopheline antiplatelet protein, a saliva protein from a malaria vector mosquito. *Blood* 2008;111:2007-14.

Zhou G, Isoe J, Day WA, Miesfeld RL. Alpha-COPI coatomer protein is required for rough endoplasmic reticulum whorl formation in mosquito midgut epithelial Cells. *PLoS One* 2011;6:e18150.

Zieler H, Garon CF, Fischer ER, Shahabuddin M. A tubular network associated with the brush-border surface of the *Aedes aegypti* midgut: implications for pathogen transmission by mosquitoes. *J Exp Biol* 2000;203:1599-611.

Zieler H and Dvorak JA. Invasion in vitro of mosquito midgut cells by the malaria parasite proceeds by a conserved mechanism and results in death of the invaded midgut cells. Proc Natl Acad Sci USA 2000; 97:11516-21.

Zizka Z, Weiser J, Blumauerova M, Jizba J. Ultrastructural effects of macrotetrolides of *Streptomyces griseus* LKS-1 in tissues of *Culex pipiens* larvae. Cytobios 1988;233:85-91.

Zocevic A, Carmi-Leroy A, Sautereau J, d'Alayer J, Lenormand P, Rousselle JC, et al. New markers in *Anopheles gambiae* salivary glands after *Plasmodium berghei* infection. Vector Borne Zoonotic Dis 2013;13:119-27.

Zouache K, Voronin D, Tran-Van V, Laurence M, Failloux AB, Mavingui P. Persistent *Wolbachia* and cultivable bacteria infection in the reproductive and somatic tissues of the mosquito vector *Aedes albopictus*. PLoS ONE 2009;4:e6388.

Zouache K, Raharimalala FN, Raqui V, Tran-Van V, Harivelo L, Ravelonandro P, et al. Bacterial diversity of field-caught mosquitoes, *Aedes albopictus* and *Aedes aegypti*, from different geographic region of Madagascar. FEMS Microbiol Ecol 2010;75:377-89.

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่  
Copyright© by Chiang Mai University  
All rights reserved

## LIST OF PUBLICATIONS

- 1) **Phattanawiboon B**, Jariyapan N, Roytrakul S, Paemanee A, Sor-suwan S, Intakhan N, Chanmol W, Siriyasatien P, Saeung A, Choochote W. Morphological and protein analyses of adult female salivary glands of *Anopheles barbirostris* species A1 (Diptera: Culicidae). *Trop Biomed* 2014;31:813-27.
- 2) **Phattanawiboon B**, Jariyapan N, Mano C, Roytrakul S, Paemanee A, Sor-Suwan S, Sriwichai P, Saeung A, Bates PA. Salivary gland proteome during adult development and after blood feeding of female *Anopheles dissidens* mosquitoes (Diptera: Culicidae). *PLoS ONE* 2016;11:e0163810.

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่  
Copyright© by Chiang Mai University  
All rights reserved