CONTENTS

Acknowledgement	c
Abstract in Thai	d
Abstract in English	f
List of Tables	j
List of Figures	1
Statement of Originality	р
Chapter 1 Introduction	1
Chapter 2 Literature Review	4
2.1 Caddisflies; Trichoptera	4
2.2 Trichoptera and man-made freshwater ecosystem	7
2.3 Affects to upstream and downstream	10
Chapter 3 Method	12
3.1 Sampling area; Mae Ngat Somboonchol dam	12
3.2 Site descriptions	12
3.3 Physicochemical water sampling	18
3.4 Trichoptera sampling and identification	20
3.5 Data analysis 19 hts reserved	23
Chapter 4 Results	24
4.1 Physicochemical water analyzed	24
4.2 Trichoptera assemblage	30
Chapter 5 Discussion	67
Chapter 6 Conclusion	71

References	7	4
Appendix	8	2
Appendix A	8	2
Appendix B	9	5
Appendix C	1	04
Curriculum vitae	1	54
	. พมยนต์	



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม Copyright[©] by Chiang Mai University All rights reserved

LIST OF TABLES

Table 3.1

Table 4.1

Physicochemical of water parameters method of sampling	19
and procedure for analysis.	
	20
Average and standard deviation of Physicochemical	29
parameters of sampling sites in Mae Ngat Somboonchol	
dam during June 2013 - May 2014. (one way ANOVA	

Page

	p<0.05)	
Table 4.2	Total number of larvae Trichoptera individuals, species	36
	richness, diversity index and evenness index in Mae Ngat	
	Somboonchol dam during June 2013 - May 2014	

Table 4.3	Total number of adult male Trichoptera individuals,	46
	species richness, diversity index and evenness index in	
	Mae Ngat Somboonchol dam during June 2013 - May	
	2014.	

Table 4.4	Comparison of abundance and occurrence of larvae and	49
	adult stage of Trichoptera in SU1 on Mage Ngat	
0	Somboonchol dam during June 2013 - May 2014.	

- Table 4.5
 Comparison of abundance and occurrence of larvae and
 50 adult stage of Trichoptera in SU2 on Mage Ngat Somboonchol dam during June 2013 - May 2014. d
- Table 4.6 Comparison of abundance and occurrence of larvae and 51 adult stage of Trichoptera in SD3 on Mage Ngat Somboonchol dam during June 2013 - May 2014.
- Table 4.7 Comparison of abundance and occurrence of larvae and 52 adult stage of Trichoptera in SD4 on Mage Ngat Somboonchol dam during June 2013 - May 2014.

- Table 4.8Comparison of abundance and occurrence of larvae and53adult stage of Trichoptera in SD5 on Mage NgatSomboonchol dam during June 2013 May 2014.
- Table 4.9Comparison of abundance and occurrence of larvae and54adult stage of Trichoptera in SD6 on Mage NgatSomboonchol dam during June 2013 May 2014.
- Table 4.10Comparison of abundance and occurrence of larvae and
adult stage of Trichoptera in SD7 on Mage Ngat
Somboonchol dam during June 2013 May 2014.55
- Table 4.11Species occurrence and abundance of adult male56Trichoptera in Mae Ngat Somboonchol, compared with
previous data from Thapanya *et al.* (2013)56



ลิฮสิทธิ์มหาวิทยาลัยเชียงใหม่ Copyright[©] by Chiang Mai University All rights reserved

k

LIST OF FIGURES

Figures 2.1	A framework for assessing the impact of dams on river	10
	ecosystems	
Figures 2.2	Hydrological changes on dammed river.	11
Figures 3.1	All 7 sampling sites sorting by upstream to downstream.	15
Figures 3.2	Drawing map showed all sampling sites in Mae Ngat	15
	Somboonchol dam.	
Figures 3.3	Upstream sampling site 1 (SU1) on August 2013.	16
Figures 3.4	Upstream sampling site 2 (SU2) on August 2013	16
Figures 3.5	Downstream sampling site (SD3) on September 2013.	16
Figures 3.6	Downstream sampling site 4 (SD4) on October 2013.	17
Figures 3.7	Downstream sampling site 5 (SD5) on September 2013.	17
Figures 3.8	Downstream sampling site 6 (SD6) on May 2014.	17
Figures 3.9	Downstream sampling site 7 (SD7) on May 2014.	17
Figures 3.10	Genitalia of a hypothetical Cheumatopsyche species on	22
	genital structures	
Figures 3.11	Light pan trap for adult Trichoptera in sampling site.	22
Figures 4.1	Monthly of air temperatures (°C) of all sampling site in Mae	25
А	Ngat Somboonchol dam during June 2013 - May 2014.	
Figures 4.2	Monthly of water temperatures (°C) of all sampling site in	25
	Mae Ngat Somboonchol dam during June 2013 - May 2014.	
Figures 4.3	Monthly total dissolved solid (TDS: mg/l – bar graph) and	26
	conductivity: μ s/cm – line graph) of all sampling sites in Mae	
	Ngat Somboonchol dam during June 2013 – May 2014.	

Figures 4.4	Monthly of pH of all sampling site in Mae Ngat Somboonchol	26
	dam during June 2013 - May 2014.	
Figures 4.5	Monthly of DO (mg/l) of all sampling site in Mae Ngat	27
	Somboonchol dam during June 2013 - May 2014.	
Figures 4.6	Monthly of BOD ₅ (mg/l) of all sampling site in Mae Ngat	27
	Somboonchol dam during June 2013 - May 2014.	
Figures 4.7	Monthly water nutrients (mg/l) of sampling sites in Mae Ngat	28
	Somboonchol dam during June 2013 – May 2014	
Figures 4.8	The dendrogram of cluster analysis based on the correlation of	30
	water parameter among all sampling sites in Mae Ngat	
	Somboonchol dam from June 2013 – May 2014.	
Figures 4.9	Numbers of individuals of larvae Trichoptera in SU1, Mae	31
	Ngat Somboonchol dam during June 2013 – May 2014	
Figures 4.10	Numbers of individuals of larvae Trichoptera in SU2, Mae	32
	Ngat Somboonchol dam during June 2013 – May 2014.	
Figures 4.11	Numbers of individuals of larvae Trichoptera in SD3, Mae	33
	Ngat Somboonchol dam during June 2013 – May 2014.	
Figures 4.12	Numbers of individuals of larvae Trichoptera in SD4, Mae	33
	Ngat Somboonchol dam during June 2013 – May 2014.	
Figures 4.13	Numbers of individuals of larvae Trichoptera in SD5, Mae	34
	Ngat Somboonchol dam during June 2013 – May 2014.	
Figures 4.14	Numbers of individuals of larvae Trichoptera in SD6, Mae	35
C	Ngat Somboonchol dam during June 2013 – May 2014	
Figures 4.15	Numbers of individuals of larvae Trichoptera in SD7, Mae	35
<u></u>	Ngat Somboonchol dam during June 2013 – May 2014.	
Figures 4.16	The dendrogram of cluster analysis based on larvae	36
	Trichoptera diversity and abundance among all sampling sites	
	in Mae Ngat Somboonchol dam from June 2013 – May 2014.	
Figures 4.17	Number of larvae individuals (dotted graph) and water depth	37
	(bar graph) monthly found during June 2013 – May 2014 in	
	Mae Ngat Somboonchol dam	

- Figures 4.18 The comparison of number of individuals and species richness 39 in SU1 during June 2013 May 2014.
- Figures 4.19 The comparison of number of individuals and species richness 40 in SU2 during June 2013 May 2014.
- Figures 4.20 The comparison of number of individuals and species richness 41 in SD3 during June 2013 May 2014.
- Figures 4.21 The comparison of number of individuals and species richness 42 in SD4 during June 2013 May 2014.
- Figures 4.22 The comparison of number of individuals and species richness 43 in SD5 during June 2013 May 2014.
- Figures 4.23 The comparison of number of individuals and species richness 43 in SD6 during June 2013 May 2014.
- Figures 4.24 The comparison of number of individuals and species richness 44 in SD7 during June 2013 May 2014.
- Figures 4.25Monthly in species richness of Trichoptera between all45sampling sites in Mae Ngat Somboonchol dam during June2013 May 2014.
- Figures 4.26 Monthly in abundance of adult male Trichoptera between all 46 sampling sites in Mae Ngat Somboonchol dam during June 2013 – May 2014.
- Figures 4.27 Comparison of average water width (m) between upstream 64 and downstream sampling sites in Mae Ngat Somboonchol dam during June 2013 – May 2014.
- Figures 4.28 Comparison of average water depth (cm) between upstream 65 and downstream sampling sites in Mae Ngat Somboonchol dam during June 2013 – May 2014.
- Figures 4.29Comparison of average water velocity (m/s) between65upstream and downstream sampling sites in Mae NgatSomboonchol dam during June 2013 May 2014.

- Figures 4.30 Comparison of average larvae family richness and number of 66 individuals between upstream and downstream sampling sites in Mae Ngat Somboonchol dam during June 2013 May 2014.
- Figures 4.31 Comparison of average adult male of Trichoptera family richness and number of individuals between upstream and downstream sampling sites in Mae Ngat Somboonchol dam during June 2013 – May 2014.



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่ Copyright[©] by Chiang Mai University All rights reserved

66

ข้อความแห่งการริเริ่ม

้ ได้นำเสนอความรู้เกี่ยวกับการเปลี่ยนแปลงทางความหลากหลายและความอุดม วิทยาบิพบซ์นี้ แมลงหนอนปลอกน้ำในแม่น้ำที่ถูกควบคุมการไหล โดยเงื่อนแม่งัดสมบูรณ์ชล ซึ่งแมลงหนอน ปลอกน้ำเป็นแมลงที่มีระยะตัวอ่อนอาศัยอยู่ในน้ำ และตัวเต็มวัยอาศัยอยู่บนบก

ามยนต

 เพื่อเพิ่มฐานข้อมูลชนิดของแมลงหนอนปลอกน้ำในประเทศไทยให้ครอบคลุมมากขึ้น การเก็บ ข้อมูลของแมลงหนอนปลอกน้ำจากแหล่งน้ำที่มีมนุษย์ควบคุมยังมีการศึกษาน้อย ซึ่งข้อมลใน แสดงให้เห็นความเปลี่ยนแปลงของกลุ่มประชากรของแมลงหนอนปลอกน้ำที่ วิทยานิพนธ์นี้ แตกต่างกันระหว่างพื้นที่เหนือเบื่อนและท้ายเบื่อน จะนำไปสู่ความเข้าใจของผลกระทบจากการ ้ควบคุมการใหลของน้ำ อันจะนำไปสู่การอนุรักษ์ทรัพยากรแหล่งน้ำจืดโดยใช้แมลงหนอนปลอก น้ำ หรือสิ่งมีชีวิตขนาดเล็กเป็นดัชนีชี้วัดต่อไป



Copyright[©] by Chiang Mai University All rights reserved

STATEMENTS OF ORIGINALITY

- The knowledge of changes on diversity and abundance of Trichoptera from water regulation by Mae Ngat Somboonchol dam, which larvae stage lived in water and adults are terrestrial insects.
- 2) To fill up the Thai Trichoptera database. This study will show the changes of Trichoptera that vary between upstream and downstream of the dam and will lead to a better understanding of input of the regulated river. This will contribute to the conservation of freshwater ecosystem used by Trichoptera or macroinvertebrate as bioindicators.



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่ Copyright[©] by Chiang Mai University All rights reserved