

CONTENTS

| | Page |
|---|-------------|
| Acknowledgements | iii |
| Abstract in English | iv |
| Abstract in Thai | vi |
| Table of Contents | viii |
| List of Figures | xv |
| List of Tables | xix |
| Abbreviations and symbols | xx |
| Chapter 1 Introduction | 1 |
| Chapter 2 Literatures review | 5 |
| 2.1 Avian influenza virus | 5 |
| 2.1.1 Influenza virus proteins | 6 |
| 2.1.1.1 Haemagglutinin (HA) | 6 |
| 2.1.1.2 Neuraminidase (NA) | 8 |
| 2.1.1.3 The others protein | 8 |
| 2.1.2 Influenza virus receptors and replication | 10 |
| 2.1.3 Antigenic shift and drift | 11 |
| 2.1.4 Human Infections by Avian Influenza H5N1 | 12 |
| 2.1.5 Epidemiology of Human Cases of Avian Influenza H5N1 | 13 |

| | Page |
|--|-------------|
| 2.1.6 Laboratory Diagnosis | 13 |
| 2.1.6.1 Virus isolation | 13 |
| 2.1.6.2 Antigen detection | 14 |
| 2.1.6.3 RT-PCR | 14 |
| 2.1.6.4 Antibody detection | 15 |
| 2.2 <i>Pichia pastoris</i> | 15 |
| 2.2.1 Methanol metabolism in <i>Pichia pastoris</i> | 16 |
| 2.2.2 The <i>Pichia</i> expression systems | 17 |
| 2.2.2.1 Expression vector | 17 |
| 2.2.2.2 The promoter in <i>Pichia pastoris</i> | 20 |
| 2.2.2.3 Transformation and integration into the <i>P. pastoris</i> genome | 21 |
| 2.2.2.4 Intracellular and secretory protein expression | 23 |
| 2.2.2.5 Posttranslational modification | 23 |
| 2.2.2.6 Process of <i>Pichia pastoris</i> fermentation | 25 |
| 2.2.3 Application of <i>Pichia pastoris</i> | 26 |
| 2.2.3.1 Therapeutics application | 26 |
| 2.2.3.2 Enzymes production | 27 |
| 2.2.3.3 Recombinant methylotrophs as a biocatalyst | 27 |
| 2.2.3.4 Methanol removing | 27 |

| | Page |
|--|-------------|
| Chapter 3 Materials and methods | 28 |
| 3.1 Bacteria, yeast, plasmid, primers and gene | 28 |
| 3.1.1 Bacteria | 28 |
| 3.1.2 Yeast | 28 |
| 3.1.3 Plasmid | 28 |
| 3.1.4 Gene | 28 |
| 3.1.5 Primers | 29 |
| 3.1.6 Positive serum and negative serum of avian influenza A virus CMU strain | 30 |
| 3.2 General equipments and chemical | 30 |
| 3.2.1 Equipments | 30 |
| 3.2.2 Chemicals | 31 |
| 3.3 Media | 33 |
| 3.4 Buffers and reagents | 33 |
| 3.5 Construction of recombinant vectors | 33 |
| 3.5.1 Vector preparation | 33 |
| 3.5.2 Amplification of H5N1 HA2 gene | 34 |
| 3.5.3 Purification of DNA product | 35 |
| 3.5.3.1 Sample Preparation | 35 |
| 3.5.3.2 DNA Purification | 36 |
| 3.5.4 <i>Xho</i> I and <i>Not</i> I digestion | 36 |
| 3.5.5 Ligation | 37 |
| 3.5.6 Preparation of competent <i>Escherichia coli</i> | 38 |

| | Page |
|--|-------------|
| 3.5.7 Transformation of recombinant plasmid into <i>Escherichia coli</i> | 38 |
| 3.5.8 Confirmation of recombinant clones using colony PCR technique | 38 |
| 3.5.9 Isolation of plasmid DNA and restriction analysis | 39 |
| 3.6 Transformation of recombinant vector in <i>Pichia pastoris</i> | 40 |
| 3.6.1 Vector preparation | 40 |
| 3.6.2 Linearization of recombinant vector by <i>SacI</i> | 40 |
| 3.6.3 Precipitation of DNA with isopropanol | 41 |
| 3.6.4 Preparation of competent yeast cell | 41 |
| 3.6.5 Transformation of yeast cells by electroporation | 42 |
| 3.7 PCR analysis of <i>Pichia</i> integrants | 42 |
| 3.8 Agarose gel electrophoresis | 43 |
| 3.9 Cultivation and induction of <i>Pichia pastoris</i> containing H5N1 HA1_OPT or H5N1 HA2 gene | 44 |
| 3.10 Preparation of yeast cell lysate using ultrasonicator | 44 |
| 3.11 Purification of recombinant H5N1 HA protein from <i>Pichia pastoris</i> using HisPur™ Ni-NTA Spin Column under native condition | 45 |
| 3.11.1 Sample preparation | 45 |
| 3.11.2 Purification of H5N1 HA protein | 45 |
| 3.12 Analysis of H5N1 HA1_OPT or H5N1 HA2 protein using SDS-PAGE and Western blot | 46 |
| 3.12.1 SDS-PAGE gel electrophoresis | 46 |
| 3.12.2 Coomassie Blue Staining of SDS-PAGE gels | 46 |

| | Page |
|---|-------------|
| 3.12.3 Western blot | 46 |
| 3.12.3.1 Detection with HisDetector™ Western Blot Kits | 46 |
| 3.12.1.2 Detection with Anti-myc antibody | 47 |
| 3.12.1.3 Detection with human serum | 47 |
| 3.13 Estimation of protein concentration | 47 |
| Chapter 4 Results | 48 |
| 4.1 Construction of recombinant vectors | 48 |
| 4.1.1 Preparation of expression vectors | 48 |
| 4.1.2 Construction of recombinant plasmid containing H5N1 HA genes | 49 |
| 4.1.2.1 Construction of pPICZA-H5N1 HA1_OPT and transformation into <i>Escherichia coli</i> XL1-blue | 49 |
| 4.1.2.2 Construction of pPICZαB-H5N1 HA1_OPT and transformation into <i>Escherichia coli</i> XL1-blue | 52 |
| 4.1.2.3 Construction of pPICZA-H5N1 HA2 and transformation into <i>Escherichia coli</i> XL1-blue | 55 |
| 4.1.2.4 Construction of pPICZαB -H5N1 HA2 and transformation into <i>Escherichia coli</i> XL1-blue | 60 |
| 4.2 Linearization of recombinant vector with <i>SacI</i> | 63 |
| 4.3 Transformation of recombinant vectors into <i>Pichia pastoris</i> | 65 |
| 4.4 PCR analysis of <i>Pichia</i> Integrants | 67 |
| 4.4.1 Analysis of recombinant <i>Pichia</i> containing H5N1 HA1_OPT gene | 67 |

| | Page |
|---|-------------|
| 4.4.2 Analysis of recombinant <i>Pichia</i> containing H5N1 HA2 gene | 68 |
| 4.5 Expression of H5N1 HA gene by recombinant <i>Pichia pastoris</i> | 70 |
| 4.5.1 Expression of H5N1 HA1_OPT gene by recombinant <i>Pichia pastoris</i> | 70 |
| 4.5.1.1 Expression of H5N1 HA1_OPT gene by recombinant <i>Pichia pastoris</i> containing pPICZA-H5N1 HA1_OPT vector | 70 |
| 4.5.1.2 Expression of H5N1 HA1_OPT gene by recombinant <i>Pichia pastoris</i> containing pPICZ α B-H5N1 HA1_OPT vector | 70 |
| 4.5.2 Expression of H5N1 HA2gene by recombinant <i>Pichia pastoris</i> | 72 |
| 4.5.2.1 Expression of H5N1 HA2gene by recombinant <i>Pichia pastoris</i> containing pPICZA-H5N1 HA2 vector | 72 |
| 4.5.2.2 Expression of H5N1 HA2gene by recombinant <i>Pichia pastoris</i> containing pPICZ α B-H5N1 HA2 vector | 76 |
| 4.6 Estimation of protein concentration | 77 |
| Chapter 5 Discussions | 78 |
| 5.1 Cloning of haemagglutinin domain (H5N1 HA gene) in <i>Pichia pastoris</i> | 79 |
| 5.2 Expression of Haemagglutinin domain (H5N1 HA gene) by <i>Pichia pastoris</i> | 80 |
| 5.2.1 Secretory expression of haemagglutinin domain (H5N1 HA gene) by <i>Pichia pastoris</i> | 80 |
| 5.2.2 Intracellular expression of haemagglutinin domain (H5N1 HA gene) by <i>Pichia pastoris</i> | 81 |

| | Page |
|-----------------------------|-------------|
| Chapter 6 Conclusion | 84 |
| Reference | 86 |
| Appendices | 101 |
| Appendix A | 102 |
| Appendix B | 104 |
| Appendix C | 105 |
| Appendix D | 108 |
| Appendix E | 112 |
| Appendix F | 113 |
| Appendix G | 114 |
| Curriculum Vitae | 115 |



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
 Copyright© by Chiang Mai University
 All rights reserved

LIST OF FIGURES

| | Page | |
|------------|---|----|
| Figure 2.1 | Schematic diagram of influenza A virus | 6 |
| Figure 2.2 | Schematic diagram of influenza virus replication | 11 |
| Figure 2.3 | <i>Pichia pastoris</i> | 15 |
| Figure 2.4 | The methanol pathway in <i>Pichia pastoris</i> | 17 |
| Figure 2.5 | General diagram of <i>P. pastoris</i> expression vector | 18 |
| Figure 2.6 | Vector diagram of pPICZ and pPICZ α | 19 |
| Figure 2.7 | Gene insertions in <i>Pichia pastoris</i> | 22 |
| Figure 2.8 | Gene replacements in <i>Pichia pastoris</i> | 22 |
| Figure 2.9 | Protein glycosylation patterns | 25 |
| Figure 4.1 | Agarose gel electrophoresis of expression vectors | 49 |
| Figure 4.2 | H5N1 HA1_OPT gene and fusion tags | 50 |
| Figure 4.3 | The colonies of <i>Escherichia coli</i> containing recombinant plasmid (pPICZA-H5N1 HA1_OPT) on LB low salt agar containing of 50 μ g/ml zeocin | 50 |
| Figure 4.4 | Analysis of the recombinant plasmid (pPICZA H5N1 HA1_OPT) by colony PCR technique using specific primer | 51 |
| Figure 4.5 | Agarose gel electrophoresis of recombinant plasmid (pPICZA-H5N1 HA1_OPT) with <i>Xho</i> I and <i>Not</i> I restriction analysis | 51 |
| Figure 4.6 | Agarose gel electrophoresis of H5N1 HA1_OPT gene purification | 52 |

| | Page |
|-------------|---|
| Figure 4.7 | The colonies of <i>Escherichia coli</i> XL1-blue containing recombinant plasmid (pPICZ α B-H5N1 HA1_OPT) on LB low salt agar containing 50 μ g/ml zeocin 53 |
| Figure 4.8 | Analysis of the recombinant plasmid (pPCZ α B-H5N1 HA1_OPT) by colony PCR technique 54 |
| Figure 4.9 | Agarose gel electrophoresis of recombinant plasmid (pPICZ α B-H5N1 HA2) with <i>Xho</i> I and <i>Not</i> I restriction analysis 54 |
| Figure 4.10 | H5N1 HA2 gene and restriction enzymes sites 55 |
| Figure 4.11 | Agarose gel electrophoresis of PCR products (H5N1 HA2 gene) amplified by using High Fidelity PCR 56 |
| Figure 4.12 | The colonies of <i>Escherichia coli</i> XL1-blue containing pPICZA-H5N1 HA2 on LB low salt agar containing 50 μ g/ml zeocin 57 |
| Figure 4.13 | Analysis of the recombinant plasmid (pPICZA-H5N1 HA2) by colony PCR technique 58 |
| Figure 4.14 | Agarose gel electrophoresis of recombinant plasmid (pPICZA-H5N1 HA2) with <i>Xho</i> I and <i>Not</i> I restriction analysis 58 |
| Figure 4.15 | Nucleotide sequence of the H5N1 HA2 gene from recombinant plasmid (pPICZA-H5N1 HA2) compared with the haemagglutinin domain gene of avian influenza A (H5N1) virus strain CMU H5 (A/Chicken/Chiang Mai/1/2004) 59 |
| Figure 4.16 | Agarose gel electrophoresis of H5N1 HA2 gene purification 60 |
| Figure 4.17 | The colonies of <i>Escherichia coli</i> XL1-blue containing recombinant plasmid (pPICZ α B-H5N1 HA2) on low salt LB agar containing 50 μ g/mL zeocin 61 |

| | Page | |
|-------------|---|----|
| Figure 4.18 | Analysis of the recombinant plasmid (pPCZ α B-H5N1 HA2) by colony PCR technique | 62 |
| Figure 4.19 | Restriction analysis of recombinant plasmid (pPICZ α B-H5N1 HA2) with <i>Xho</i> I and <i>Not</i> I | 62 |
| Figure 4.20 | Agarose gel electrophoresis of the recombinant vectors digested with <i>Sac</i> I | 64 |
| Figure 4.21 | The colonies of <i>Pichia pastoris</i> containing recombinant plasmid on selective medium using 100 μ l of transformant suspension on YPD agar containing 100 μ g/ml zeocin | 66 |
| Figure 4.22 | PCR products of twelve selected recombinant <i>Pichia</i> colonies containing pPICZA-H5N1 HA1_OPT vector | 67 |
| Figure 4.23 | PCR products of twelve selected recombinant <i>Pichia</i> colonies pPICZ α B-H5N1 HA1_OPT vector | 68 |
| Figure 4.24 | PCR products of twelve selected recombinant <i>Pichia</i> colonies containing pPICZA-H5N1 HA2 vector | 69 |
| Figure 4.25 | PCR products of twelve selected recombinant <i>Pichia</i> colonies pPICZ α B-H5N1 HA2 vector | 69 |
| Figure 4.26 | SDS-PAGE analysis of H5N1 HA2 protein with coomassie blue staining | 72 |
| Figure 4.27 | Western blot analysis of H5N1 HA2 protein detected by HisDetector TM Western Blot Kits (KPL, Inc., USA) | 73 |
| Figure 4.28 | Western blot analysis of H5N1 HA2 protein detected by Anti-myc Antibody | 74 |
| Figure 4.29 | Western blot analysis of H5N1 HA2 protein detected by PO3 (positive serum) | 75 |

Figure 4.30 Western blot analysis of H5N1 HA2 protein detected by CS serum negative)

75



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved

LIST OF TABLES

| | | Page |
|------------|---|-------------|
| Table 2.1 | Another protein of avian influenza segments and their encoded polypeptides | 9 |
| Table 3.1 | Sequence of specific primers for H5N1 HA1_OPT gene | 29 |
| Table 3.2 | Sequence of specific primers for H5N1 HA2 gene | 29 |
| Table 3.3 | Equipment | 30 |
| Table 3.4 | Chemicals | 31 |
| Table 3.5 | The reaction mixture of high-fidelity PCR | 35 |
| Table 3.6 | The digestion mixture of expression vector and H5N1 gene | 36 |
| Table 3.7 | The ligation mixture of expression vectors and H5N1 HA genes | 37 |
| Table 3.8 | The reaction mixture of colony PCR | 39 |
| Table 3.9 | The digestion mixture of recombinant expression vector (<i>XhoI</i> and <i>NotI</i>) | 40 |
| Table 3.10 | The digestion mixture of expression vector | 41 |
| Table 3.11 | The reaction mixture of colony PCR | 43 |
| Table 4.1 | Estimated protein concentration of H5N1 HA2 protein purified by HisPur™ Ni-NTA Spin Column under native condition | 77 |

ABBREVIATION AND SYMBOLS

| | |
|--------------------|------------------------|
| rpm | revolutions per minute |
| bp | base pair |
| kDa | kilodalton |
| kb | kilobase |
| μg | microgram |
| mg | milligram |
| g | gram |
| μM | micromole |
| μm | micrometer |
| dH ₂ O | distilled water |
| l | liter |
| ml | milliliter |
| U | unit |
| % | percent |
| $^{\circ}\text{C}$ | degree Celsius |
| h | hour |
| min | minute |
| sec | second |
| MW | molecular weight |

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright © by Chiang Mai University
All rights reserved