

REFERENCES

- Alam, F., Date, A., Rasjidin, R., Mobin, S., Moria, H. and Baqui, A. 2012. Biofuel from algae- is it a viable alternative?. *Procedia Eng.* 49: 221–227.
- Anastassiadis, S. and Rehm, H.J. 2006. Oxygen and temperature effect on continuous citric acid secretion in *Candida oleophila*. *J Biotechnol.* 9(4): 1–12.
- Angerbauer, C., Siebenhofer, M., Mittelbach, M. and Guebitz, G.M. 2008. Conversion of sewage sludge into lipids by *Lipomyces starkeyi* for biodiesel production. *Bioresour Technol.* 99: 3051–3056.
- Behera, S., Singh, R., Arora, R., Sharma, N.K., Shukla, M. and Kumar, S. 2014. Scope of algae as third generation biofuels. *Front Bioeng Biotechnol.* 2: 1–13.
- Beopoulos, A. and Nicaud, J.M. 2012. Yeast: A new oil producer?. *OCL.* 19: 22–28.
- Bessey, O.A., Lowry, O.H. and Brock, M.J. 1946. A method for the rapid determination of alkaline phosphates with five cubic millimeters of serum. *J Biol Chem.* 164: 321–329.
- Bligh, E.G. and Dyer, W.J. 1959. A rapid method of total lipid extraction and purification. *Can J Biochem Phys.* 37: 911–917.
- Braunwald, T., Schwemmlin, L., Graeff-Hönninger, S., French, W.T., Hernandez, R., Holmes, W.E. and Claupein, W. 2013. Effect of different C/N ratios on carotenoid and lipid production by *Rhodotorula glutinis*. *Appl Microbiol Biotechnol.* 97: 6581–6588.
- Cardoso, L.A.C., Jäckel, S., Karp, S.G. Framboisier, X., Chevalot, I. and Marc, I. 2016. Improvement of *Sporobolomyces ruberrimus* carotenoids production by the use of raw glycerol. *Bioresour Technol.* 200: 374–379.

- Chaiyaso, T., Kuntiya, A., Techapun, C., Leksawasdi, N., Seesuriyachan, P. and Hanmoungjai, P. 2011. Optimization of cellulase-free xylanase production by thermophilic *Streptomyces thermovulgaris* TISTR1948 through Plackett-Burman and response surface methodological approaches. *Biosci Biotechnol Biochem.* 75: 531–537.
- Chaiyaso, T., Seesuriyachan, P., Zimmermann, W. and H-Kittikun, A. 2012. Purification and characterization of lipase from newly isolated *Burkholderia multivorans* PSU-AH130 and its application for biodiesel production. *Ann Microbiol.* 62: 1615–1624.
- Chandi, GK. and Gill, B.S. 2011. Production and characterization of microbial carotenoids as an alternative to synthetic colors: A review. *Int J Food Prop.* 14: 503–513.
- Cheirsilp, B., Suwannarat, W. and Niyomdecha, R. 2011. Mixed culture of oleaginous yeast *Rhodotorula glutinis* and microalga *Chlorella vulgaris* for lipid production from industrial wastes and its use as biodiesel feedstock. *New Biotechnol.* 28: 362–368.
- Chi, Z., Zheng, Y., Jiang, A. and Chen, S. 2011. Lipid production by culturing oleaginous yeast and algae with food waste and municipal wastewater in an integrated process. *Appl Biochem Biotechnol.* 165: 442–453.
- Chodok, P., Kanjana-Opas, A. and Kaewsuan, S. 2010. The Plackett-Burman design for evaluating the production of polyunsaturated fatty acids by *Physcomitrella patens*. *J Am Oil Chem Soc.* 87: 521–529.
- Christophe, G., Kumar, V., Nouaille, R., Gaudet, G., Fontanille, P., Pandey, A. Soccol, C.R. and Larroche, C. 2012. Recent developments in microbial oils production: a possible alternative to vegetable oils for biodiesel without competition with human food?. *Braz Arch Biol Technol.* 55: 29–46.
- Derrien, M., Badr, A., Gosselin, A., Desjardins, Y. and Angers, P. 2017. Optimization of a green process for the extraction of lutein and chlorophyll from spinach by-products using response surface methodology (RSM). *Food Sci Technol.* 79: 170–177.

- Dong, T., Knoshaug, E.P., Pienkos, P.T. and Laurens, L.M.L. 2016. Lipid recovery from wet oleaginous microbial biomass for biofuel production: A critical review. *Appl Energ.* 177: 879–895.
- Donot, F., Fontana, A., Baccou, J.C., Strub, C. and Schorr-Galindo, S. 2014. Single cell oils (SCOs) from oleaginous yeasts and moulds: Production and genetics. *Biomass Bioenerg.* 68: 135–150.
- Dubois, M., Gilles, K.A., Hamilton, J.K., Rebers, P.A. and Smith, F. 1956. Colorimetric method for determination of sugars and related substances. *Anal Chim.* 28: 350–356.
- Dwivedi, G. and Sharma, M.P. 2015. Investigation and improvement in cold flow properties of pongamia biodiesel. *Waste Biomass Valori.* 6: 73–79.
- Easterling, E.R, French, W.T., Hernandez, R. and Licha, M. 2009. The effect of glycerol as a sole and secondary substrate on the growth and fatty acid composition of *Rhodotorula glutinis*. *Bioresour Technol.* 100: 356–361.
- Elbeshbishy, E., Hafez, H., Dhar, B.R. and Nakhla, G. 2011. Single and combined effect of various pretreatment methods for biohydrogen production from food waste. *Int J Hydrogen Energ.* 36: 11379–11387.
- El-Banna, A.A., El-Razek, A.M.A. and El-Mahdy, AR. 2012. Some factors affecting the production of carotenoids by *Rhodotorula glutinis* var. *glutinis*. *J Food Nutr Sci.* 3: 64–71.
- Gambelli, D., Alberti, F., Solfanelli, F., Vairo, D. and Zanolli, R. 2017. Third generation algae biofuels in Italy by 2030: A scenario analysis using Bayesian networks. *Energ Policy.* 103: 165–178.
- Gen, Q., Wang, Q. and Chi, Z.M. 2014. Direct conversion of cassava starch into single cell oil by co-cultures of the oleaginous yeast *Rhodospiridium toruloides* and immobilized amylases-producing yeast *Saccharomycopsis fibuligera*. *Renew Energ.* 62: 522–526.

- Hadi, R.S. and Ivan, M. 2006. Effect of temperature and pH on the growth kinetics and carotenoid production by *Sporobolomyces ruberrimus* H110 using technical glycerol as carbon source. Iran J Chem Chem Eng. 25: 59–64.
- Hall, T.A. 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. Nucl Acids Symp Ser. 41: 95–98.
- Han, M., Xu, Z., Du, C., Qian, H. and Zhang, W.G. 2016. Effects of nitrogen on the lipid and carotenoid accumulation of oleaginous yeast *Sporidiobolus pararoseus*. Bioproc Biosyst Eng. 39: 1425–1433.
- Helrich, K. 1990. official methods of analysis of AOAC international, 15th ed. Association of official analytical chemists. Arlington, USA.
- Hong, Y.S. and Yoon, H.H. 2011. Ethanol production from food residues. Biomass Bioenerg. 35: 3271–3275.
- Hu, C., Zhao, X., Zhao, J., Wu, S. and Zhao, Z.K. 2009. Effects of biomass hydrolysis by-products on oleaginous yeast *Rhodospiridium toruloides*. Bioresour Technol. 100: 4843–4847.
- Jiménez Callejón, M.J., Robles Medina, A., Macías Sánchez, M.D., Hita Peña, E., Esteban Cerdán, L., González Moreno, P.A. and Molina Grima, E. 2014. Extraction of saponifiable lipids from wet microalgal biomass for biodiesel production. Bioresour Technol. 169: 198–205.
- Karatay, S.E. and Dönmez, G. 2010. Improving the lipid accumulation properties of the yeast cells for biodiesel production using molasses. Bioresour Technol. 101: 7988–7990.
- Karimi, K., Kheradmandinia, S. and Taherzadeh, M.J. 2006. Conversion of rice straw to sugars by dilute-acid hydrolysis. Biomass Bioenerg. 30: 247–253.
- Kastner, V., Somitsch, W. and Schnitzhofer, W. 2012. The anaerobic fermentation of food waste: A comparison of two bioreactor systems. J Clean Prod. 34: 82–90.
- Kavšček, M., Bhutada, G., Madl, T. and Natter, K. 2015. Optimization of lipid production with a genome-scale model of *Yarrowia lipolytica*. BMC Syst Biol. 9: 1–13.

- Kim, D.H., Kim, S.H., Kim, K.Y. and Shin, H.S. 2010. Experience of a pilot-scale hydrogen-producing anaerobic sequencing batch reactor (ASBR) treating food waste. *Int J Hydrogen Energ.* 35: 1590–1594.
- Kim, J.H., Lee, J.C. and Pak, D. 2011. Feasibility of producing ethanol from food waste. *Waste Manage.* 31: 2121–2125.
- Kirti, K., Amita, S., Priti, S., Mukesh Kumar, A. and Jyoti, S. 2014. Colorful world of microbes: carotenoids and their applications. *J Adv Biol.* 2014: 1–13.
- Kitcha, S. and Cheirsilp, B. 2011. Screening of oleaginous yeasts and optimization for lipid production using crude glycerol as a carbon source. *Energy Procedia.* 9: 274–282.
- Kolouchová, I., Sigle, K., Schreiberová, O., Masák, J. and Řezanka, T. 2015. New yeast-based approaches in production of palmitoleic acid. *Bioresour Technol.* 192: 726–734.
- Kraisintu, P., Yongmanitchai, W. and Limtong, S. 2010. Selection and optimization for lipid production of a newly isolated oleaginous yeast, *Rhodosporidium toruloides* DMKU3-TK16. *Kasetsart J. (Nat. Sci.).* 44: 436–445.
- Kumara, C.G. and Takagi, H. 1999. Microbial alkaline proteases: From a bioindustrial viewpoint. *Biotechnol Adv.* 17: 561–594.
- Lamers, D., Biezen, N.V., Martens, D., Peters, L., Zilver, E.V.D., Dreumel, N.J.V., Wiffels, R.H. and Likman, C. 2016. Selection of oleaginous yeasts for fatty acid production. *BMC Biotechnol.* 16: 1–10.
- Lau, K.Y., Pleissner, D. and Lin, C.S.K. 2014. Recycling of food waste as nutrients in *Chlorella vulgaris* cultivation. *Bioresour Technol.* 170: 144–151.
- Li, H., Yang, L., Kim, Y.J. and Kim, S.J. 2011. Continuous ethanol production by the synchronous saccharification and fermentation using food wastes. *Korean J Chem Eng.* 28: 1085–1089.
- Li, M., Liu, G.L., Chi, Z. and Chi, Z.M. 2010. Single cell oil production from hydrolysate of cassava starch by marine-derived yeast *Rhodotorula mucilaginosa* TJY15a. *Biomass Bioenerg.* 34: 101–107.

- Liang, M.H. and Jiang, J.G. 2013. Advancing oleaginous microorganisms to produce lipid via metabolic engineering technology. *Prog Lipid Res.* 52: 395–408.
- Lin, H., Wang, Q., Shen, Q., Zhan, J. and Zhao, Y. 2013. Genetic engineering of microorganisms for biodiesel production. *Bioeng.* 4: 292–304.
- Lin, J., Shen, H., Tan, H., Zhao, X., Wu, S., Hu, C. and Zhao, Z.K. 2011. Lipid production by *Lipomyces starkeyi* cells in glucose solution without auxiliary nutrients. *J Biotechnol.* 152: 184–188.
- Liu, J.X., Yue, Q.Y., Gao, B.Y., Ma, Z.H. and Zhang, P.D. 2012. Microbial treatment of the monosodium glutamate wastewater by *Lipomyces starkeyi* to produce microbial lipid. *Bioresour Technol.* 106: 69–73.
- Ma, H., Wang, Q., Qian, D., Gong, L. and Zhan, W. 2009. The utilization of acid-tolerant bacteria on ethanol production from kitchen garbage. *Renew Energ.* 34: 1466–1470.
- Malisorn, C. and Suntornsuk, W. 2008. Optimization of β -carotene production by *Rhodotorula glutinis* DM28 in fermented radish brine. *Bioresour Technol.* 99: 2281–2287.
- Manowattana, A., Seesuriyachan, P., Techapun, C. and Chaiyaso, T. 2012. Optimization of carotenoids production by red yeast *Sporobolomyces pararoseus* TISTR5213 using waste glycerol as the sole carbon source. *KKU Res J.* 17(4): 607–621.
- Manowattana, A., Techapun, C., Seesuriyachan, P., Hanmoungjai, P. and Chaiyaso, T. 2015. β -Carotene production by *Sporobolomyces pararoseus* TISTR5213 using crude glycerol as the sole carbon source. *Chiang Mai J Sci.* 42(1): 17–33.
- Manysoat, S. 2013. Production of sato and distilled spirit from job's tears. Master degree thesis, Chiang Mai University, Chiang Mai, Thailand.
- Martinez, E.J., Raghavan, V., Gonzalez-Andres, F. and Gomez, X. 2015. New biofuel alternatives: Integrating waste management and single cell oil production. *Int J Mol Sci.* 16: 9385–9405.

- Mata-Gómez, L.C., Montañez, J.C., Méndez-Zavala, A. and Aguilar, C.N. 2014. Biotechnological production of carotenoids by yeasts: An overview. *Microb Cell Fact.* 13: 1–11.
- Matsakas, L., Kekos, D., Loizidou, M. and Christakopoulos, P. 2014. Utilization of household food waste for the production of ethanol at high dry material content. *Biotechnol Biofuels.* 7: 1–9.
- McCurry, J.D. 2011. GC-MS analysis of trace fatty acid methyl esters (FAME) in Jet fule using energy institute method P585. Agilent Technologies, Inc., USA. pp. 1-8.
- Meng, X., Yang, J., Xu, X., Zhang, L., Nie, Q. and Xian, M. 2009. Biodiesel production from oleaginous microorganisms. *Renew Energ.* 34: 1–5.
- Miller, G.L. 1959. Use of dinitrosalicylic acid reagent for determination of reducing sugar. *Anal Chem.* 31: 426–428.
- Moon, H.C., Song, S., Kim, J.K., Shirai, Y., Lee, D.H., Kim, J.K., Chung, S.O., Kim, D.H., Oh, K.K. and Cho, Y.S. 2009. Enzymatic hydrolysis of food waste and ethanol fermentation. *Int J Energ Res.* 33: 164–172.
- Muniraj, I.K., Xiao, L., Hu, Z., Zhan, X. and Shi, J. 2013. Microbial lipid production from potato processing wastewater using oleaginous filamentous fungi *Aspergillus oryzae*. *Water Res.* 47: 3477–3483.
- Naik, S.N., Goud, V.V., Rout, P.K. and Dalai, A.K. 2010. Production of first and second generation biofuels: A comprehensive review. *Renew Sust Energ Rev.* 14: 578–597.
- Papanikolaou, S., Dimou, A., Fakas, S., Diamantopoulou, P., Philippoussis, A., Galiotou-Panayotou, M. and Aggelis, G. 2011. Biotechnological conversion of waste cooking olive oil into lipid-rich biomass using *Aspergillus* and *Penicillium* strains. *J Appl Microbiol.* 110: 1138–1150.
- Patel, A., Arora, N., Sartaj, K., Pruthi, V. and Pruthi, PA. 2016. Sustainable biodiesel production from oleaginous yeasts utilizing hydrolysates of various non-edible lignocellulosic biomasses. *Renew Sust Energ Rev.* 62: 836–855.

- Patnayak, S. and Sree, A. 2005. Screening of bacterial associates of marine sponges for single cell oil and PUFA. *Lett Appl Microbiol.* 40(5): 358–363.
- Peramuna, A. and Summers, M.L. 2014. Composition and occurrence of lipid droplets in the cyanobacterium *Nostoc punctiforme*. *Arch Microbiol.* 196: 881–890.
- Pisutpaisal, N., Nathao, C. and Sirisukpoka, U. 2014. Biological hydrogen and methane production in from food waste in two-stage CSTR. *Energy Procedia.* 50: 719–722.
- Pleissner, D., Lam, W.C., Sun, Z. and Lin, C.S.K. 2013. Food waste as nutrient source in heterotrophic microalgae cultivation. *Bioresour Technol.* 137: 139–146.
- Poontawee, R., Yongmanitchai, W. and Limtong, S. 2017. Efficient oleaginous yeasts for lipid production from lignocellulosic sugars and effects of lignocellulose degradation compounds on growth and lipid production. *Process Biochem.* 53: 44–60.
- Priatni, S. 2014. Review: Potential production of carotenoids from *Neurospora*. *Nus Biosci.* 6: 63–68.
- Probst, K.V., Schulte, L.R., Durrett, T.P., Rezac, M.E. and Vadlani, P.V. 2016. Oleaginous yeast: a value-added platform for renewable oils. *Crit Rev Biotechnol.* 36: 942–955.
- Saenge, C., Cheirsilp, B., Suksaroge, T. and Bourtoom, T. 2011a. Efficient concomitant production of lipids and carotenoids by oleaginous red yeast *Rhodotorula glutinis* cultured in palm oil mill effluent and application of lipids for biodiesel production. *Biotechnol Bioeng.* 16: 23–33.
- Saenge, C., Cheirsilp, B., Suksaroge, T.T. and Bourtoom, T. 2011b. Potential use of oleaginous red yeast *Rhodotorula glutinis* for the bioconversion of crude glycerol from biodiesel plant to lipids and carotenoids. *Process Biochem.* 46: 210–218.

- Schneider, T., Graeff-Honninger, S., French, W.T., Hernandez, R., Claupein, W., Holmes, W.E. and Merkt, N. 2012. Screening of industrial wastewaters as feedstock for the microbial production of oils for biodiesel production and high-quality pigments. *J Combust.* 2012: 1–9.
- Schneider, T., Rempp, T., Graeff-Hönninger, S., French, W.T., Hernandez, R. and Claupein, W. 2013. Utilization of soluble starch by oleaginous red yeast *Rhodotorula glutinis*. *Sust Bioenerg Syst.* 3: 57–63.
- Sha, Q. 2013. A comparative study on four oleaginous yeasts on their lipid accumulating capacity. Master degree Thesis, Swedish University of Agricultural Sciences.
- Sharp, A. and Sang-Aru, J. 2012. A Guide for sustainable urban organic waste management in Thailand: Combining food, energy, and climate co-benefits. Institute for Global Environmental Strategies (IGES), Japan.
- Sitepu, I.R., Garay, L.A., Sestric, R., Levin, D., Block, D.E., German, J.B. and Boundy-Mills, K.L. 2014. Oleaginous yeasts for biodiesel: Current and future trends in biology and production. *Biotechnol Adv.* 32: 1336–1360.
- Sitepu, I.R., Sestric, R., Ignatia, L., Levin, D., German, J.B., Gillies, L.A., Luis A.G., Almada, L.A.G. and Boundy-Mills, K.L. 2013. Manipulation of culture conditions alters lipid content and fatty acid profiles of a wide variety of known and new oleaginous yeast. *Bioresour Technol.* 144: 360–369.
- Soccol, C.R., Dalmas Neto, C.J., Soccol, V.T., Sydney, E.B., Costa, E.S.F., Medeiros A.B.P. and Vandenberghe, L.P.S. 2017. Pilot scale biodiesel production from microbial oil of *Rhodosporidium toruloides* DEBB 5533 using sugarcane juice: Performance in diesel engine and preliminary economic study. *Bioresour Technol.* 223: 259–268.
- Somashekar, D. and Joseph, R. 2000. Inverse relationship between carotenoid and lipid formation in *Rhodotorula gracilis* according to the C/N ratio of the growth medium. *World J Microbiol Biotechnol.* 16: 491–493.

- Subramaniam, R., Dufreche, S., Zappi, M. and Bajpai, R. 2010. Microbial lipids from renewable resources: Production and characterization. *J Ind Microbiol Biot.* 37: 1271–1287.
- Sundarram, A. and Murthy, T.P.K. 2014. α -Amylase production and applications: A review. *Appl Environ Microb.* 2: 166–175.
- Tanimura, A., Takashima, A., Sugita, T., Endoh, R., Kikukawa, M., Yamaguchi, S., Sakuradani, E., Ogawa, J., Ohkuma, M. and Shima, J. 2014. *Cryptococcus terricola* is a promising oleaginous yeast for biodiesel production from starch through consolidated bioprocessing. *Sci Rep.* 4: 1–6.
- Taskin, M., Ortucu, S., Aydogan, M.N. and Arslan, N.P. 2016. Lipid production from sugar beet molasses under non-aseptic culture conditions using the oleaginous yeast *Rhodotorula glutinis* TR29. *Renew Energ.* 99: 198–204.
- Teste, M.A., François, J.M. and Parrou, J.L. 2010. Characterization of a new multigene family encoding isomaltases in the yeast *Saccharomyces cerevisiae*, the IMA family. *J Biol Chem.* 285: 26815–26824.
- Thevenieau, F. and Nicaud, J.M. 2013. Microorganisms as sources of oils. *OCL.* 20: 1–8.
- Thi, N.B.D., Kumar, G. and Lin, C.Y. 2015. An overview of food waste management in developing countries: Current status and future perspective. *J Environ Manage.* 157: 220–229.
- Tsigie, Y.A., Wang, C.Y., Truong, C.T. and Ju, Y.H. 2011. Lipid production from *Yarrowia lipolytica* Po1g grown in sugarcane bagasse hydrolysate. *Bioresour Technol.* 102: 9216–9222.
- Uçkun Kiran, E., Trzcinski, A.P., Jern, N.W. and Liu, Y. 2014. Bioconversion of food waste to energy: A review. *Fuel.* 134: 389–399.
- Verma, P., Sharma, M.P. and Dwivedi, G. 2016. Evaluation and enhancement of cold flow properties of palm oil and its biodiesel. *Energ Rep.* 2: 8–13.

- Viegas, C.V., Hachemi, I., Mäki-Arvela, P., Smeds, A., Aho, A., Freitas, S.P., da Silva Gorgônio, C.M., Carbonetti, G., Peurla, M. Paranko, J., Kumar, N., Aranda, D.A.G. and Murzin, D.Y. 2015. Algal products beyond lipids: Comprehensive characterization of different products in direct saponification of green alga *Chlorella* sp. *Algal Research*. 11: 156–164.
- Wang, Q., Guo, F.J., Rong, Y.J. and Chi, Z.M. 2012. Lipid production from hydrolysate of cassava starch by *Rhodospiridium toruloides* 21167 for biodiesel making. *Renew Energ*. 46: 164–168.
- Watanabe, M., Ichinose, K., Sasano, K., Ozaki, Y., Tsuiki, T., Hidaka, H. and Kanemoto, S. 2011. Effect of enzymatic treatment on sedimentation and flocculation abilities of solid particles in rice washing drainage and its relationship with protein profiles. *J Biosci Bioeng*. 112: 67–70.
- Wiebe, M.G., Koivuranta, K., Penttilä, M. and Ruohonen, L. 2012. Lipid production in batch and fed-batch cultures of *Rhodospiridium toruloides* from 5 and 6 carbon carbohydrates. *BMC Biotechnol*. 12: 1–10.
- Wild, R., Patil, S., Popovi, M., Zappi, M., Dufreche, S. and Bajpai, R. 2010. Lipids from *Lipomyces starkeyi*. *Food Technol Biotech*. 48: 329–335.
- Woiciechowski, A.L., Nitsche, S., Pandey, A. and Soccol, C.R. 2002. Acid and enzymatic hydrolysis to recover reducing sugars from cassava bagasse: An economic study. *Braz Arch Biol Technol*. 45(3): 393–400.
- Wu, S., Hu, C., Jin, G., Zhao, X. and Zhao, Z.K. 2010. Phosphate-limitation mediated lipid production by *Rhodospiridium toruloides*. *Bioresour Technol*. 101: 6124–6129.
- Wu, S., Zhao, X., Shen, H., Wang, Q. and Zhao, Z.K. 2011. Microbial lipid production by *Rhodospiridium toruloides* under sulfate-limited conditions. *Bioresour Technol*. 102: 1803–1807.
- Xie, F., Quan, S., Liu, D., Ma, H., Li, F., Zhou, F. and Chen, G. 2014. Purification and characterization of a novel α -amylase from a newly isolated *Bacillus methylotrophicus* strain P11-2. *Process Biochem*. 49: 47–53.

- Xu, J., Du, W., Zhao, X., Zhan, G. and Liu, D. 2013. Microbial oil production from various carbon sources and its use for biodiesel preparation. *Biofuels Bioprod Bior.* 7: 65–77.
- Xu, J., Zhao, X., Wang, W., Du, W. and Liu, D. 2012. Microbial conversion of biodiesel byproduct glycerol to triacylglycerols by oleaginous yeast *Rhodospiridium toruloides* and the individual effect of some impurities on lipid production. *Biochem Eng J.* 65: 30–36.
- Xue, F., Gao, B., Zhu, Y., Zhang, X., Feng, W. and Tan, T. 2010. Pilot-scale production of microbial lipid using starch wastewater as raw material. *Bioresour Technol.* 101: 6092–6095.
- Yan, S., Li, J., Chen, X., Wu, J., Wang, P., Ye, J. and Yao, J. 2011. Enzymatical hydrolysis of food waste and ethanol production from the hydrolysate. *Rev Energ.* 36: 1259–1265.
- Yang, X., Jin, G., Wang, Y., Shen, H. and Zhao, Z.K. 2015. Lipid production on free fatty acids by oleaginous yeasts under non-growth conditions. *Bioresour Technol.* 193: 557–562.
- Yasin, N.H.M., Rahman, N.A.A., Man, H.C., Yusoff, M.Z.M. and Hassan, M.A. 2011. Microbial characterization of hydrogen-producing bacteria in fermented food waste at different pH values. *Int J Hydrogen Energ.* 36: 9571–9580.
- Yen, H.W. and Liu, Y.X. 2014. Application of airlift bioreactor for the cultivation of aerobic oleaginous yeast *Rhodotorula glutinis* with different aeration rates. *J Biosci Bioeng.* 118: 195–198.
- Yen, H.W. and Zhang, Z. 2011. Enhancement of cell growth rate by light irradiation in the cultivation of *Rhodotorula glutinis*. *Bioresour Technol.* 102: 9279–9281.
- Yu, X., Zheng, Y., Dorgan, K.M. and Chen, S. 2011. Oil production by oleaginous yeasts using the hydrolysate from pretreatment of wheat straw with dilute sulphuric acid. *Bioresour Technol.* 102: 6134–6140.

- Zeng, Y., Bian, D., Xie, Y., Jiang, X., Li, X., Li, P., Zhang, Y. and Xie, T. 2017. Utilization of food waste hydrolysate for microbial lipid and protein production by *Rhodospiridium toruloides* Y2. J Chem Technol Biotechnol. 92: 666–673.
- Zhang, D., Ong, Y.L., Li, Z. and Wu, J.C. 2013. Biological detoxification of furfural and 5-hydroxyl methyl furfural in hydrolysate of oil palm empty fruit bunch by *Enterobacter* sp. FDS8. Biochem Eng J. 72: 77–82.
- Zhang, Y., Wang, Y., Wang, Z.G., Wang, X., Guo, H.S., Meng, D.F. and Wong, P.K. 2012. Optimization of fermentation medium for the production of atrazine degrading strain *Acinetobacter* sp. DNS₃₂ by statistical analysis system. J Biomed Biotechnol. 7: 1–7.
- Zhang, Z., Schwartz, S., Wagner, L. and Miller, W. 2000. A greedy algorithm to aligning DNA sequences. J Comput Biol. 7: 203–214.
- Zhang, Z., Zhang, X. and Tan, T. 2014. Lipid and carotenoid production by *Rhodotorula glutinis* under irradiation/high-temperature and dark/low-temperature cultivation. Bioresour Technol. 157: 149–153.
- Zhang, G., French, W.T., Hernandez, R., Alley, E. and Paraschivescu, M. 2011. Effects of furfural and acetic acid on growth and lipid production from glucose and xylose by *Rhodotorula glutinis*. Biomass Bioeng. 35: 734–740.
- Zhu, L.Y., Zong, M.H. and Wu, H. 2008. Efficient lipid production with *Trichosporon fermentans* and its use for biodiesel preparation. Bioresour Technol. 99: 7881–7885.
- Zoz, L., Carvalho, J.C., Soccol, V.T., Casagrande, T.C. and Cardoso, L. 2015. Torularhodin and torulene: Bioproduction, properties and prospective applications in food and cosmetics - a review. Braz Arch Biol Technol. 58: 278–288.