

LIPIDS AND CAROTENOIDS PRODUCTIONS FROM OLEAGINOUS YEASTS USING RICE RESIDUE FROM FOOD WASTE AS A CARBON SOURCE



GRADUATE SCHOOL
CHIANG MAI UNIVERSITY
MAY 2017


LIPIDS AND CAROTENOIDS PRODUCTIONS FROM OLEAGINOUS YEASTS USING RICE RESIDUE FROM FOOD WASTE AS A CARBON SOURCE



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่ Copyright[©] by Chiang Mai University All rights reserved

GRADUATE SCHOOL
CHIANG MAI UNIVERSITY
MAY 2017

LIPIDS AND CAROTENOIDS PRODUCTIONS FROM OLEAGINOUS YEASTS USING RICE RESIDUE FROM FOOD WASTE AS A CARBON SOURCE



A THESIS SUBMITTED TO CHIANG MAI UNIVERSITY IN PARTIAL FULFILLMENT OF THE REQUIREMENTS FOR THE DEGREE OF

MASTER OF SCIENCE
IN BIOTECHNOLOGY

l rights reserved

GRADUATE SCHOOL, CHIANG MAI UNIVERSITY

MAY 2017

LIPIDS AND CAROTENOIDS PRODUCTIONS FROM OLEAGINOUS YEASTS USING RICE RESIDUE. FROM FOOD WASTE AS A CARBON SOURCE

WIMADA SRISUWAN

THIS THESIS HAS BEEN APPROVED TO BE A PARTIAL FULFILLMENT OF THE REQUIREMENTS FOR THE DEGREE OF MASTER OF SCIENCE IN BIOTECHNOLOGY

Examination Committee:	Advisory Committee:
A Dit Chairman	T. Chaiyaso Advisor
(Dr. Punyawatt Pintathong)	(Asst. Prof. Dr. Thanongsak Chaiyaso)
Cham Techapu Member	Chami Techapen Co-advisor
(Asst. Prof. Dr. Charin Techapun)	(Asst. Prof. Dr. Charin Techapun)
T. Chaiy aso Member (Asst. Prof. Dr. Thanongsak Chaiyaso)	
Chamint Mallemoner Member	
(Dr. Churairat Moukamnerd)	

25 May 2017 Copyright © by Chiang Mai University

ACKNOWLEDGEMENT

I would like to express my special thanks of gratitude to my advisor, Asst. Prof. Dr. Thanongsak Chaiyaso, Division of Biotechnology, School of Agro-Industry, Faculty of Agro-Industry, Chiang Mai University, for his kindness, invaluable advice, generous support and consistent encouragement throughout my research, and for also educating me to develop my scientific thinking and problem solving skills. This research was accomplished with his kindness understanding and assistance.

Similarly, I also would like to thank my co-advisor, Asst. Prof. Dr. Charin Techapun, Division of Biotechnology, School of Agro-Industry, Faculty of Agro-Industry, Chiang Mai University for valuable comments and suggestions, Dr. Punyawatt Pintathong, School of Cosmetic Science, Mae Fah Luang University, for his helpful suggestions, and also the external chairperson, Dr. Churairat Moukamnerd, Division of Biotechnology, Faculty of Agro-Industry, Chiang Mai University, for her helpful suggestions to an external member. Moreover, I would like to express my gratitude to all staff members in the Division of Biotechnology, Faculty of Agro-Industry, Chiang Mai University, for their teaching and guidance.

This work has been financially supported by the Thesis Grant for Master Degree Student FY2014, Tetra Pak (Thailand) Ltd., Thailand, Faculty of Agro-Industry and the Graduate School, Chiang Mai University, FY2016 Thesis Grant for Master Degree Student, The National Research Council of Thailand (NRCT), Thailand, Japan Student Services Organization (JASSO) Student Exchange Support Program (Scholarship for Short-Term Study in Japan) and Japan Science and Technology Agency, Japan-Asia Youth Exchange program in Science, Japan.

Finally, I would like to express my deepest gratitude to my family, especially my parents for their love and encouragement to overcome every difficulty and understanding during the period of this study.

Wimada Srisuwan