

REFERENCES

- [1] Peng L., Song X., Shi X., Li J. and Ye C., “An improved HPLC method for simultaneous determination of phenolic compounds, purine alkaloids and theanine in *Camellia species*,” *Journal of Food Composition and Analysis*, 21, 2008, 559-563.
- [2] Saito S.T., Fröhlich P.E., Gosmann G. and Bergold A.M., “Full validation of a simple method for determination of catechins and caffeine in brazilian green tea (*Camellia sinensis* var. *assamica*) using HPLC,” *Chromatographia*, 65, 2007, 607-610.
- [3] Yang X.R., Ye C.X., Xu J.K., Jiang M., “Simultaneous analysis of purine alkaloids and catechins in *Camellia sinensis*, *Camellia ptilophylla* and *Camellia assamica* var. *kucha* by HPLC.” *Food Chemistry*, 100, 2007, 1132-1136.
- [4] Chan E.W.C., Lim Y.Y. and Chew Y.L., “Antioxidant activity of *Camellia sinensis* leaves and tea from a lowland plantation in Malaysia,” *Food Chemistry*, 102, 2007, 1214-1222.
- [5] Gramzał A., Korczak J. and Amarowicz R., “Tea polyphenols – their antioxidant properties and biological activity,” *Polish Journal of Food and Nutrition Sciences*, 3, 2005, 219-235.
- [6] [Online] “Non-fermented tea”, from [https://m.teavivre.com/info/fermentation of-teas](https://m.teavivre.com/info/fermentation-of-teas), 16 June 2017.
- [7] [Online] “Oolong tea”, from <http://www.tenren.com/fermentation>, 16 June 2017.
- [8] [Online] “Black tea”, from [https://www.teatulia.com/tea-varieties/what-is blacktea.htm](https://www.teatulia.com/tea-varieties/what-is-blacktea.htm), 16 June 2017.
- [9] Graham H. N., “Green tea composition, consumption, and polyphenol chemistry.” *Preventive Medicine*, 21, 1992, 334-350.
- [10] Nonaka G.I., Kawahara O. and Nishioka I., “Tannins and related compounds XV11 A new class of dimeric flavan-3-ol gallates, theasinensins A and B, and

- proanthocyanidingallates from green tea leaf,” *Chemical and Pharmaceutical Bulletin*, 11, 1983, 3906-3914.
- [11] Nishitani E. and Sagesaka M., “Simultaneous determination of catechins, caffeine and other phenolic compounds in tea using new HPLC method,” *Journal of Food Composition and Analysis*, 17, 2004, 675-685.
- [12] Samuel T.S., Grace G., Jenifer S., Márcio P., Marc F.R. and Ana M.B., “Characterization of the constituents and antioxidant activity of Brazilian green tea (*Camellia sinensis* var. *assamica* IAC-259 Cultivar) extracts,” *Journal of Agricultural and Food Chemistry*, 55, 2007, 9409-9414.
- [13] Hattori S., Takagaki H. and Fujimori T., “A comparison of the volatile compounds in several green teas,” *Food Science and Technology Research*, 11, 2005, 82-86.
- [14] Kumazawa K. and Masuda H., “Identification of potent odorants in Japanese green tea (Sen-cha),” *Journal of Agricultural and Food Chemistry*, 47, 1999, 5169-5172.
- [15] Pripdeevech P. and Machan T., “Fingerprint of volatile flavour constituents and antioxidant activities of teas from Thailand,” *Food Chemistry*, 125, 2011, 797-802.
- [16] Hashimoto F, Nonaka G I., Nishoka I., “Tannins and related compounds. LXIX.1 isolation and structure elucidation of B,B'-linked bisflavanoids, theasinensins D-G and oolongtheanin from oolong tea. (2),” *Chemical and Pharmaceutical Bulletin*, 36, 1988, 1676-1684.
- [17] Hashimoto F, Nonaka G I. and Nishoka I., “Tannins and related compounds. XC. 8-C-Ascorbyl (-)-epigallocatechin 3-O-gallate and novelb dimeric flavan-3-ols, oolonghomobisflavans A and B, from oolong tea. (3),” *Chemical and Pharmaceutical Bulletin*, 37, 1989, 3255-3263.
- [18] Kato M. and Shibamoto T., “Variation of major volatile constituents in various green teas from southeast Asia,” *Journal of Agricultural and Food Chemistry*, 49, 2001, 1394-1396.
- [19] Wang, K.B., Liu, F. and Liu, Z.H., “Analysis of chemical components in oolong tea in relation to perceived quality,” *International Journal of Food Science & Technology*, 45, 2010, 913-920.

- [20] Takeo, T., "Variations in the aroma compound content of semi-fermented and black tea," *Journal of Agricultural and Food Chemistry*, 57, 1983, 457-459.
- [21] Takami C., Shimotsukasa A. and Kobayashi A., "Aroma components of Oolong tea," *Journal of Agricultural and Food Chemistry*, 64, 1990, 1349-1354.
- [22] Hashimoto F, Nonaka G. I., Nishoka I., "Tannins and related compounds. CXIV.1 structures of novel fermentation products, theogallinin, theaflavonin and desgalloyl theaflavonin from black tea, and changes of tea leaf polyphenols during fermentation," *Chemical and Pharmaceutical Bulletin*, 40, 1992, 1383-1389.
- [23] Hashimoto F., Nonaka G.I., and Nishioka I., "Tannins and related compounds. LXXVII. novel chalcone-flavan dimers, assamicains A, B and C, and a new flavan-3-ol and proanthocyanidins from the fresh leaves of *Camellia sinensis* L. var. *assamica* KITAMURA," *Chemical and Pharmaceutical Bulletin*, 37, 1989, 77-85.
- [24] Wang B. S., Yu M. H., Chang L. W., Yen W. J. and Duh, P. D., "Protective effects of pu-erh tea on LDL oxidation and nitric oxide generation in macrophage cells. *Journal of Food Science and Technology*, 41, 2008a, 1122-1132.
- [25] Wang L.F., Lee J.Y., Chung J.O., Baik J.H., So S. and Park, S.K., "Discrimination of teas with different degrees of fermentation by SPME-GC analysis of the characteristic volatile flavor compounds," *Food Chemistry*, 109, 2008b, 196-206.
- [26] Ravichandran R., "Carotenoid composition distribution and degradation to flavour volatiles during black tea manufacture and the effect of carotenoid supplementation on tea quality and aroma," *Food Chemistry*, 78, 2002, 23-28.
- [27] Morita M., Wakabayashi M., Kubota K., Kobayashi A. and Herath N.L., "Aglycone constituents in fresh tea leaves cultivated for green and black tea," *Bioscience, Biotechnology, and Biochemistry*, 58, 1994, 687-690.
- [28] Mick W. and Schreie P., "Additional volatiles of black tea aroma," *Journal of Agricultural and Food Chemistry*, 32, 1984, 924-929.
- [29] Sanderson G.W. and Graham H.N., "On the formation of black tea aroma," *Journal of Agricultural and Food Chemistry*, 21, 1973, 576-585.

- [30] Kawakami M., Chairote G. and Kobayashi A., "Flavor constituents of pickle tea miang in Thailand," *Agricultural and Biological Chemistry*. 51, 1987, 1683-1687.
- [31] Korsamphan C., Boonma W. and Srigarm K., "Diversity of local tea on highland of northern Thailand," *Journal of Agricultural and Food Chemistry*, 26, 2010, 93-99.
- [32] Phromrukachat S., Tiengburanatum N. and Meechui J., "Assessment of active ingredients in pickled tea," *Asian Journal of Food and Agro-Industry*, 3, 2010, 312-318.
- [33] Klayraung S. and Siriporn O., "Antibacterial and antioxidant activities of acid and bile resistant strains of *Lactobacillus fermentum* isolated from miang," *Brazilian Journal of Microbiology*, 40, 2009, 326-367.
- [34] Ketwal S., Chueamchaitrakun P., Theppakorn T. and Wongsakul S., "Contents of total polyphenol, microorganisms and antioxidant capacities of pickled tea (Miang) commercially available in Chiang Rai, Thailand," *The 16th Food Innovation Asia Conference 2014*, at Bangkok, Thailand.
- [35] Halliwell B. and Gutteridge J.M.C., "*Free Radicals in Biology and Medicine*", 3rd ed., Oxford: Clarendon Press, 1999, 416-494.
- [36] Karadag A., Ozcelik B. and Saner S., "Review of Methods to Determine Antioxidant Capacities," *Food Analytical Methods*, 2, 2009, 41-60.
- [37] Schillacia C., Nepravishtaa R. and Bellomariaa A., "Antioxidants in food and pharmaceutical research," *Albanian Journal of Pharmaceutical Sciences*, 1, 2013, 2313-1772.
- [38] Sies H., "Oxidative stress: oxidants and antioxidants," *Experimental Physiology*, 82, 1997, 29-295.
- [39] Chu Y.H., Chang C.L. and Hsu H.F., "Flavonoid content of several vegetables and their antioxidant activity," *Journal of the Science of Food and Agriculture*, 80, 2000, 561-566.
- [40] Nollet L. M. L., "*Food Analysis by HPLC*," 2nd ed., Marcel, Inc., New York, 2000, pp. 775-824.
- [41] Tsao, R. and Yang, R., "Optimization of a new mobile phase to know the complex and Real polyphenolic composition: towards a total phenolic index

- using highperformance liquid chromatography,” *Journal of Chromatography A*, 29, 2003, 29-40.
- [42] Ezoubeiri A., Gadhi C.A., Fdil N., Benharref A., Jana M. and Vanhaelen M., “Isolation and antimicrobial activity of two phenolic compounds from *Pulicaria odora* L.,” *Journal of Ethnopharmacology*, 99, 2005, 287-292.
- [43] Pedreschi R. and Cisneros-Zevallos, L., “Phenolic profiles of andean purple corn (*Zea mays* L.),” *Food Chemistry*, 100, 2007, 956-963.
- [44] Kähkönen M. P. and Heinonen M., “Antioxidant activity of anthocyanins and their aglycons,” *Journal of Agricultural and Food Chemistry*, 51, 2003, 628-633.
- [45] Cook N.C. and Samman S., “Flavonoids-chemistry, metabolism, cardioprotective effects, and dietary sources,” *The Journal of Nutritional Biochemistry*, 7, 1996, 66-76.
- [46] Harborne, J. B., “*Plant phenolics*. In: BELL EA, CHARLWOOD BV (eds) *Encyclopedia of Plant Physiology*,” volume 8 Secondary Plant Products, SpringerVerlag, Berlin Heidelberg, New York.1980, pp: 329-395.
- [47] Singleton V.L and Rossi J.A., “Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents,” *American Journal of Enology and Viticulture*, 16, 1965, 144-158.
- [48] Magalhaes L.M., Segundo M.A., Reis S. and Lima. J.L.F.C., “Methodological aspects about in vitro evaluation of antioxidant properties. *Analytica Chimica Acta*, 1, 2008, 613.
- [49] MacDonald-Wicks L.K., Wood L.G. and Garg M.L., “Methodology for the determination of biological antioxidant capacity in vitro: a review,” *Journal of the Science of Food and Agriculture*, 86, 2006, 2046.
- [50] Yamaguchi T., Takamura H., Matoba T. and Terao, J., “HPLC method for evaluation of the free radical-scavenging activity of foods by using 1,1-diphenyl-2-picrylhydrazyl,” *Bioscience, Biotechnology, and Biochemistry*, 62, 1998, 1201-1204.
- [51] Nur Alam M.D., Jahan Bristi N. and Rafiquzzaman M.D., “Review on in vivo and in vitro methods evaluation of antioxidant activity,” *Saudi Pharmaceutical Journal*, 21, 2013, 143-152.

- [52] Siddhuraju P., Mohan P. S. and Becker K., "Studies on the antioxidant activity of Indian Laburnum (*Cassia fistula* L.): a preliminary assessment of crude extracts from stem bark, leaves, flowers and fruit pulp," *Food Chemistry*, 79, 2002, 61-67.
- [53] Chung Y.C., Chang C.T., Chao W.W., Lin C.F. and Chou S.T., "Antioxidative activity and safety of the 50% ethanolic extract from red bean fermented by *Bacillus subtilis* IMR-NK1," *Journal of Agricultural and Food Chemistry*, 50, 2002, 2454.
- [54] Gu I., Huyut Z., Elmastas M. and Aboul-Enein H.Y., "Radical scavenging and antioxidant activity of tannic acid," *Arabian Journal of Chemistry*, 3, 2010, 43-53.
- [55] Guo Q., Zhao B., Shen S., Hou J., Hu, J. and Xin W., "ESR study on the structure antioxidant activity relationship of tea catechins and their epimers," *Biochimica et Biophysica Acta*, 1427, 1999, 13-23.
- [56] Michalak, "Phenolic compounds and their antioxidant activity in plants growing under heavy metal stress," *Polish Journal of Environmental Studies*, 15, 2006, 523-530.
- [57] Nakagawa, T., Yokozawa, T., "Direct scavenging of nitric oxide and superoxide by green tea," *Food and Chemical Toxicology*, 40, 2002, 1745-1750.
- [58] Nanjo F., Honda M., Okushio K., Matsumoto N., Ishigaki F., Ishigami T., and Hara, Y., "Effects of dietary tea catechins on alpha-tocopherol levels, lipid peroxidation, and erythrocyte deformability in rats fed on high palm oil and perilla oil diet," *Biological and Pharmaceutical Bulletin*, 16, 1993, 1156-1159.
- [59] Velayutham P., Babu A. and Liu, D., "Green tea catechins and cardiovascular health: an update," *Current Medicinal Chemistry*, 15, 2008, 1840-1850.
- [60] Gramza A., Pawlak-Lemańska K., Korczak J. and Wsowicz E., "Tea extracts as free radicals scavengers," *Polish Journal of Environmental Studies*, 32, 2005, 78-89.
- [61] Chen Z.Y., Chan P.T., Ma H.M., Fung K.P. and Wang J., "Antioxidative effect of ethanol tea extracts on oxidation of canola oil," *Journal of the American Oil Chemists Society*, 73, 1996b, 375-380.

- [62] Su Y., Xu J., Ng C.H., Leung L.K., Huang Y. and Chen Z., "Antioxidant activity of tea theaflavins and methylated catechins in canola oil. *Journal of the American Oil Chemists Society*, 81, 2004, 269-272.
- [63] Cao G., Sofic E. and Prior R.L., "Antioxidant capacity of tea and common vegetables," *Journal of Agricultural and Food Chemistry*, 44, 1996, 3426-343.
- [64] Gadow A., Joubert E. and Hansmann C.F., "Comparison of the antioxidant activity of rooibos tea (*Aspalathus linearis*) with green, oolong and black tea," *Food Chemistry*, 60, 1997, 73-77.
- [65] Matsuzaki T.L. and Hara Y., "Antioxidative activity of tea leaf catechins," *International Journal of Food Science & Technology*, 59, 1985, 129-134.
- [66] Tanizawa H., Sazuka Y., Komatsu A., Toda S. and Taniko Y., "A new efficacy test of antioxidants based on air-oxidation of linoleic," *Chemical and Pharmaceutical Bulletin*, 31, 1983, 4139-4143.
- [67] Amarowicz R. and Shahidi F., "Antioxidant activity of green tea catechins in a b-carotene-linoleate model system," *Journal of Food Lipids*, 2, 1995, 47-56.
- [68] Chen C.-W. and Ho C.-T., "Antioxidant properties of polyphenols extracted green and black teas," *Journal of Food Lipids*, 2, 1995, 35-46.
- [69] Wanasundara P. K. J. P. D. and Shahidi, F., "*Antioxidants: science, technology and applications*," In F. Shahidi (Ed.), *Bailey's industrial oil and fat products*, 2005, pp. 431-489.
- [70] Frankel E. N. and Huang S.-W. and Aescbach R., "Antioxidant activity of green teas in different lipid systems," *Journal of the American Oil Chemists Society*, 74, 1997, 1309-1315.
- [71] Nanjo F., Goto K., Seto R., Suzuki M., Sakai M. and Hara Y., "Scavenging effects of tea catechins and their derivatives on 1,1-diphenyl-2-picrylhydrazyl radical," *Free Radical Biology & Medicine*, 21, 1996, 895-902.
- [72] Salah N., Miller N.J., Paganga G. and Tijburg L., "Polyphenolic flavanols as scavengers of aqueous phase radicals and as chain-breaking antioxidants," *Archives of Biochemistry and Biophysics*, 322, 1995, 339-346.
- [73] Terao J., Piskula M. and Yao Q., "Protective effect of epicatechin, epicatechin gallate, and quercetin on lipid peroxidation in phospholipid bilayers". *Archives of Biochemistry and Biophysics*, 231, 1994, 278-284.

- [74] Kondo K., Masaaki K., Miyata N., Suzuki T. and Toyoda M., “Mechanistic studies of catechins as antioxidants against radical oxidation,” *Archives of Biochemistry and Biophysics*, 362, 1999, 79-86.
- [75] Adams R. P. *“Identification of Essential Oil Components by Gas Chromatography/ Quadrupole Mass Spectrometry”*, Carol Stream, Illinois: Allured, 2007.
- [76] Kähkönen M.P., Hopia A.I., Vuorela H.J., Rauha J.P., Pihlaja K., Kujala T.S. and Heinonen M., “Antioxidant activity of plant extracts containing phenolic compounds,” *Journal of Agricultural and Food Chemistry*, 47(10), 1999, 3954-3962.
- [77] Chu Y. H., Chang C. L., and Hsu H. F., “Flavonoid content of several vegetables and their antioxidant activity,” *Journal of the Science of Food and Agriculture*, 80, 2000, 561-566.
- [78] Palleros D. R., *“Extraction and Isolation of Caffeine from Tea Leaves”* Experimental Organic Chemistry, Wiley: New York, 2000.
- [79] Pawin B., *“Identification and quantification of some phenolic compounds in fresh and fermented tea using high performance liquid chromatography-mass spectrometry”*. M.S. Thesis, Chaing Mai University, 2006.
- [80] Panda S.H., Panda S., Sivakumar P.S. and Ray R.C., “Anthocyanin-Rich Sweet Potato Lacto-Pickle: Production, Nutritional and Proximate Composition,” *International Journal of Food Science & Technology*, 44, 2009, 445-455.
- [81] Xiao P., Huang Y. Y., Yang W. P., Zhang B. W. and Quan, X. X., “Screening lactic acid bacteria with high yielding-acid capacity from pickled tea for their potential uses of inoculating to ferment tea products,” *Journal of Food Science and Technology*, 52(10), 2015, 6727-6734.
- [82] Liu H., Qiu N., Ding H., and Yao R., “Polyphenols contents and antioxidant capacity of 68 Chinese herbals suitable for medical or food uses,” *Food Research International*, 41, 2008, 363-370.
- [83] Luo Z.M., Ling T.J., Li L.X., Zhang Z.Z., Zhu H.T., Zhang Y.J. and Wan X.C., “A new norisoprenoid and other compounds from Fuzhuan brick tea,” *Molecules*, 17, 2012, 3539-3546.

- [84] Yang Q.M., Pan X. and Kong W., “Antioxidant activities of malt extract from barley (*Hordeum vulgare* L.) toward various oxidative stress *in vitro* and *in vivo*,” *Food Chemistry*, 118, 2010, 84-89.
- [85] Rawat R., Gulati A., Babu G. D. K., Acharya R., Kaul, V. K. and Singh B., “Characterization of volatile components of Kangra orthodox black tea by gas chromatography-mass spectrometry,” *Food Chemistry*, 105, 2007, 229-235.
- [86] Lewinsohn E., Sitrit Y. and Bar E., “Standardized Methods for the Determination of Antioxidant Capacity and Phenolics in Foods and Dietary Supplements,” *Journal of Agricultural and Food Chemistry*, 53, 2005, 3142-3148.
- [87] Drużyńska B., Stępniewska A. and Wołosiak R. “ The influence of time and type of solvent on efficiency of the extract of polyphenols from green and antioxidant properties obtained extracts”, *Acta Scientiarum Polonorum Technologia Alimentaria*, 6(1), 2007, 27-36.
- [88] Hseu Y.C., Chang W.H. and Chen C.S., “Antioxidant activities of *C. sinensis* leaves extracts using different antioxidant models,” *Food and Chemical Toxicology*, 46, 2008, 105-114.
- [89] Liu S.C., Lin J.T. and Wang C.K., “Antioxidant properties of various solvent extracts from lychee (*Litchi chinensis* sonn.) flowers,” *Food Chemistry*, 114, 2009, 577-581.
- [90] Luo Z.M., Ling T.J., Li L.X., Zhang Z.Z., Zhu H.T., Zhang Y.J. and Wan X.C., “A New Norisoprenoid and Other Compounds from Fuzhuan Brick Tea,” *Molecules*, 17, 2012, 3539-3546.
- [91] Krishnamurty H. G., Krishnamoorth, V. and Seshadri T. R., “Microbiological transformation of flavanone,” *Phytochemistry*, 60, 2003, 247-253.
- [92] Miller, J. C. and Miller, J. N., “*Statistics for Analytical Chemistry*,” 3rd ed., 2009, Ellis Horwood, New York, pp. 96-100.
- [93] Phromrukachat S., Tiengburanatum N. and Meechui J., “Assessment of active ingredients in pickled tea,” *Asian Journal of Food and Agro-Industry*, 3(03), 2010, 312-318.
- [94] Wulandari R.A., Amano M., Yanagita T., Tanaka T., Kouno I., Kawamura D. and Ishimaru K., “New phenolic compounds from *Camellia sinensis* L. leaves

- fermented with *Aspergillus* sp.,” *Journal of Natural Medicines*, 65, 2011 594-597.
- [95] Zhang L., Li N., Ma Z.Z. and Tu P., “Comparison of the chemical constituents of aged puerh tea, ripened pu-erh tea, and other teas using HPLC-DAD-ESI-MS,” *Journal of Agricultural and Food Chemistry*, 59, 2011, 8754-8760.
- [96] Kahkonen M. P., Hopia A. I., Vuorela H. J., Rauha J. P., Pihlaja K. and Kujala, T. S., “Antioxidant activity of plant extracts containing phenolic compounds,” *Journal of Agricultural and Food Chemistry*, 47, 1999, 3954-3962.
- [97] Ratty A. K. and Das, N. P., “Effects of flavonoids on nonenzymatic lipid peroxidation: structure-activity relationship,” *Biochemical Medicine and Metabolic Biology*, 39, 1988, 69-79.
- [98] Rice-Evans C. A., Miller N. J. and Paganga, G., “Structure-antioxidant activity relationships of flavonoids and phenolic acids,” *Free Radical Biology and Medicine*, 20, 1996, 933-956.
- [99] Miller N. J., Castelluccio C., Tijburg, L. and Rice-Evans C., “The antioxidant properties of theaflavins and their gallate esters—radical scavengers or metal chelators,” *FEBS Lett*, 392, 1996, 40-44.
- [100] Morel, I., Lescoat, G., Cogrel, P., Sergent, O., Padeloup, N., Brissot, P., Cillard, P. and Cilliard, J., “Antioxidant and iron-chelating activities of the flavonoids catechin, quercetin and diosmetin on ironloaded rat hepatocyte cultures,” *Biochemical Pharmacology*, 45, 1993, 13-19.
- [101] Guo, Q., Zhao, B., Li, M., Shen, S., and Xin, W. 1996. Studies on protective mechanisms of four components of green tea polyphenols against lipid peroxidation in synaptosomes,” *Biochimica Biophysica Acta*, 1304, 1996, 210-222.
- [102] Yu F.L., Gapor A. and Bender W., “Evidence for the preventive effect of the polyunsaturated phytol side chain in tocotrienols on 17 beta-estradiol epoxidation,” *Cancer Detection and Prevention*, 29(4), 2005, 383-388.
- [103] Senthilkumar S.R., Sivakumar T., Arulmozhi K.T. and Mythil N., “Gas Chromatography-mass spectroscopy Evaluation of bioactive phytochemicals of commercial green teas (*Camellia sinensis*) of India,” *Asian Journal of Pharmaceutical and Clinical Research*, 8(3), 2015, 278-282.

- [104] Kumar R.N., Reddy J.S., Gopikrishna G. and Anand S.K., “GC-MS Determination of Bioactive Constituents of Cycasbeddomei Cones,” International Journal of Pharma and Bio Sciences, 3, 2012, 344-350.



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved