


CONTENTS

	Page
Acknowledgement	d
Abstract in Thai	e
Abstract in English	g
List of tables	k
List of figures	m
Chapter 1 Introduction	1
1.1 Principles and rationales	1
1.2 Purposes of the study	3
1.3 Research hypothesis	3
1.4 Research designs and procedures	3
1.5 Expecting benefit and goal	3
Chapter 2 Literature review	4
2.1 Anterior open bite	4
2.2 Posterior tooth intrusion	7
2.3 Miniscrew implant placement sites, suitable for maxillary posterior Tooth intrusion	8
2.4 Quantitative and qualitative evaluations of palate for palatal miniscrew implant stability	9
2.5 Palatal bone thickness in different skeletal patterns	13
2.6 Cone beam computed tomography (CBCT)	14
Chapter 3 Materials and methods	16
Chapter 4 Results	25
4.1 Palatal bone thickness	25
4.2 Palatal cortical bone thickness	33

	Page
Chapter 5 Discussion	42
Chapter 6 Conclusions	52
References	53
Appendix	61
Appendix A Information of subjects	62
Appendix B Certificate of ethical clearance	64
Appendix C Informed consent	66
Curriculum Vitae	70




ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved

LIST OF TABLES

		Page
Table 3.1	Age (years) distribution by gender and number of subjects (N)	16
Table 3.2	General information on subjects	18
Table 3.3	Cephalometric measurements in each group	19
Table 4.1	Palatal bone thicknesses (mm) at each anteroposterior (AP) site along The mediolateral 0 (ML0) section in the Class I normal bite and Class I open bite groups	26
Table 4.2	Palatal bone thicknesses (mm) at each anteroposterior (AP) site along The mediolateral 3 (ML3) section in the Class I normal bite and Class I open bite groups	27
Table 4.3	Palatal bone thicknesses (mm) at each anteroposterior (AP) site along The mediolateral 6 (ML6) section in the Class I normal bite and Class I open bite groups	28
Table 4.4	Palatal bone thicknesses (mm) at each anteroposterior (AP) site along The mediolateral 9 (ML9) section in the Class I normal bite and Class I open bite groups	29
Table 4.5	Palatal bone thicknesses (mm) at each anteroposterior (AP) site along The mediolateral 12 (ML12) section in the Class I normal bite and Class I open bite groups	30
Table 4.6	Palatal cortical bone thicknesses (mm) at each anteroposterior (AP) site along the mediolateral 0 (ML0) section in the Class I normal bite and Class I open bite groups	34
Table 4.7	Palatal cortical bone thicknesses (mm) at each anteroposterior (AP) site along the mediolateral 3 (ML3) section in the Class I normal bite and Class I open bite groups	35

	Page
Table 4.8 Palatal cortical bone thicknesses (mm) at each anteroposterior (AP) site along the mediolateral 6 (ML6) section in the Class I normal bite and Class I open bite groups	36
Table 4.9 Palatal cortical bone thicknesses (mm) at each anteroposterior (AP) site along the mediolateral 9 (ML9) section in the Class I normal bite and Class I open bite groups	38
Table 4.10 Palatal cortical bone thicknesses (mm) at each anteroposterior (AP) site along the mediolateral 12 (ML12) section in the Class I normal bite and Class I open bite groups	39



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved

LIST OF FIGURES

		Page
Figure 3.1	CBCT images were oriented, using the Romexis viewer program.	20
Figure 3.2	The three-dimensional image shows that a line joining the middle of the distal bony margin of the incisive foramen and PNS is oriented to the midsagittal plane.	20
Figure 3.3	In the axial view, a line joining the middle of the distal bony margin of the incisive foramen and PNS is oriented to the midsagittal plane.	21
Figure 3.4	In the midsagittal view, the horizontal reference plane is oriented to pass through the middle of those distal bony margins of the incisive foramen and PNS.	21
Figure 3.5	In each sagittal view, the total palatal bone and cortical bone thicknesses were measured perpendicular to the horizontal reference plane at 3.0-mm intervals posteriorly from the middle of the distal bony margin of the incisive foramen to PNS.	22
Figure 3.6	In each frontal view, the total palatal bone and cortical bone thicknesses were measured at 3.0-mm intervals laterally from the midsagittal reference plane on both right and left sides.	23
Figure 3.7	Measurement sites in a grid pattern at 3.0-mm anteroposterior intervals posteriorly from the middle of the distal bony margin of the incisive foramen (AP 3, 6, 9, 12, 15, 18, 21, 24 mm-sections), and at 3.0-mm mediolateral intervals laterally from the midsagittal plane on both right and left sides (ML 0, 3, 6, 9, and 12 mm-sections). The measurement site marked “x” was named Left AP6/ML3.	23
Figure 3.8	Paracoronal image showing measurement of the total palatal bone thickness (yellow line) and the palatal cortical bone thickness (green line)	24

	Page
Figure 4.1	32
<p>Combined plots of average palatal bone thickness at various sites in the Class I normal bite group. The white, translucent plane is inserted at the 5.0 mm level of palatal bone thickness, revealing the sites where the average palatal bone thickness was equal to or greater than 5.0 mm.</p>	
Figure 4.2	33
<p>Combined plots of average palatal bone thickness at various sites in the Class I open bite group. The white, translucent plane is inserted at the 5.0 mm level of palatal bone thickness, revealing the sites where the average palatal bone thickness was equal to or greater than 5.0 mm.</p>	
Figure 4.3	40
<p>Combined plots of average palatal cortical bone thickness at various sites in the Class I normal bite group. The red dash lines are inserted at the 1.0 mm level of palatal cortical bone thickness, revealing the sites where the average palatal cortical bone thickness was equal to or greater than 1.0 mm.</p>	
Figure 4.4	41
<p>Combined plots of average palatal cortical bone thickness at various sites in the Class I open bite group. The red dash lines are inserted at the 1.0 mm level of palatal cortical bone thickness, revealing the sites where the average palatal cortical bone thickness was equal to or greater than 1.0 mm.</p>	
Figure 5.1	45
<p>The palatal map shows the pattern of the bone thickness of the palate at various sites in the Class I normal bite and the Class I open bite groups. The heads of the arrows represent the direction of the increase in bone thickness. The green arrows show the pattern of the palatal bone thickness along the ML0, ML3, AP3 and AP6 sections. The orange arrows show the pattern of the palatal bone thickness in the other remaining areas.</p>	

	Page
Figure 5.2	46
<p>The palatal map shows the pattern of the cortical bone thickness of the palate at various sites in the Class I normal bite and the Class I open bite groups. The heads of the arrows represent the direction of the increase in bone thickness. The green arrows show the pattern of the palatal bone thickness along the ML0, ML3, AP3 and AP6 sections. The orange arrows show the pattern of the palatal bone thickness in the other remaining areas.</p>	
Figure 5.3	49
<p>The palatal map shows the average palatal bone thickness measurements at various sites in the Class I normal bite (Left side) and the Class I open bite groups (Right side). The green dots indicate the sites where the average palatal bone thicknesses were equal to or greater than 8 mm. The yellow dots indicate the sites where the average palatal bone thicknesses were equal to or greater than 5 mm. The red dots indicate the sites where the average palatal bone thicknesses were less than 5 mm.</p>	
Figure 5.4	50
<p>The palatal map shows the average palatal cortical bone thickness measurement at various sites in the Class I normal bite (Left side) and the Class I open bite groups (Right side). The green dots indicate the sites where the average cortical bone thicknesses were equal to or greater than 8 mm. The yellow dots indicate the sites where the average palatal cortical bone thicknesses were equal to or greater than 5 mm. The red dots indicate the sites where the average palatal cortical bone thicknesses were less than 5 mm.</p>	