

REFERENCES

- Abbott WS. A method of computing the effectiveness of an insecticide. J Econ Entomol 1925;18:265-6.
- Afolabi OJ, Simon-Oke IA, Elufisan OO, Oniya MO. Adulticidal and repellent activities of some botanical oils against malaria mosquito: *Anopheles gambiae* (Diptera: Culicidae). BJBAS 2018;7:135-8.
- Ahmad I, Astari S, Tan M. Resistance of *Aedes aegypti* (Diptera: Culicidae) in 2006 to pyrethroid insecticides in Indonesia and its association with oxidase and esterase levels. Pak J Biol Sci 2007;10:3688-92.
- Ahmad R, Yu KX, Wong CL, Jantan I. Larvicidal and adulticidal activities of malaysian seaweeds against *Aedes aegypti* (L.) and *Aedes albopictus* skuse (diptera: Culicidae). Southeast Asian J Trop Med Public Health 2016;47:719-30.
- Akıncı A, Eşrefoglu M, Taşlıdere E, Ateş B. *Petroselinum crispum* is effective in reducing stress-induced gastric oxidative damage. Balkan Med J 2017;53-9.
- Al-Howiniry T, Al-Sohaibani M, El-Tahir K, Rafatullah S. Prevention of experimentally-induced gastric ulcers in rats by an ethanolic extract of "parsley", *Petroselinum crispum*. Am J Chim Med 2003;31:699-711.
- Ali MYS, Ravikumar S, Beula JM. Mosquito larvicidal activity of seaweeds extracts against *Anopheles stephensi*, *Aedes aegypti*, and *Culex quinquefasciatus*. Asian Pac J Trop Dis 2013;3:196-201.
- Amer A, Mehlhorn H. Larvicidal effects of various essential oils against *Aedes*, *Anopheles*, and *Culex* larvae (Diptera: Culicidae). Parasitol Res 2006;99:466-72.
- Amin AM. Preliminary investigation of the mechanisms of DDT and pyrethroid resistance in *Culex quinquefasciatus* Say. (Diptera: Culicidae) from Saudi Arabia. Bull Entomol Res 1989;79:361-6

Asakura K. Phosphatase activity in the larva of the euryhalinemosquito, *Aedes togoi* Theobold, with special reference to sea water adaptation. J Exp Mar Biol Ecol 1978;31:325-37.

Assar AA, Abo El-Mahasen MM, Khalil ME, Mahmoud SH. Biochemical effects of some insect growth regulators on the house fly, *Musca domestica* (diptera: muscidae). Egypt Acad J Biol Sci 2010;2:33-44.

Babu R, Murugan K, Vanithakumari G. Interference of azadirachtin on the food utilization efficiency and gut enzymatic profiles of *Helicoverpa armigera* Hubner (Lepidoptera: Noctuidae). Indian J Environ Toxicol 1996;6:81-4.

Badolo A, Ilboudo-Sanogo E, Ouédraogo AP, Costantini C. Evaluation of the sensitivity of *Aedes aegypti* and *Anopheles gambiae* complex mosquitoes to two insect repellents: DEET and KBR 3023. Trop Med Int Hlth 2004;9:330-4.

Bassal TT, Ismail IE. Acid and phosphatases of normal and juvenilized *Spodoptera littoralis* (Noctuidae: Lepidoptera) during metamorphosis. Proc Zool Soc AR Egypt 1985;9:256-9.

Beaty BJ, Marquardt WC. The Biology of Disease Vectors. Colorado: Am J Trop Med 1996;56:482.

Ben-Dov E, Saxena D, Wang Q, Manasherob R, Boussiba S, Zaritsky A. Ingested particles reduce susceptibility of insect larvae to *Bacillus thuringiensis*. J Appl Entomol 2003;127:146-52.

BPIA. Research shows biopesticides expanding rapidly. Biopesticide Industry Alliance [Internet]. 2015 [cited 20 Aug 2015]. Available from: <https://www.bbia.org/2015/07/research-shows-biopesticides-expanding-rapidly/>

Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. Analyt Biochem 1976;72:248-54.

Bream AS. Effect of azadirachtin on phosphatases and transaminases activities in pupae of the red palm weevil *Rhynchophorus ferrugineus* (Coleoptera: Curculionidae). Proc Inter Conf Date palm, King Saud Univ: Saudi Arabia; 2003.

Brengues C, Hawkes NJ, Chandre F, McCarroll L, Duchon S, Guillet P, et al. Pyrethroid and DDT cross-resistance in *Aedes aegypti* correlated with novel mutations in the voltage-gated sodium channel gene. Med Vet Entomol 2003;17:87-94.

Brogdon WG, McAllister JC, Vulule J. Heme peroxidase activity measured in single mosquitoes identifies individuals expressing an elevated oxidase for insecticide resistance. J Am Mosq Control Assoc 1997;13:233-7.

Buglio FM, Wilkins RM. Influence of malathion resistance status on survival and growth of *Tribolium castaneum* (Coleoptera: Tenebrionidae), when fed on four from insect-resistant and susceptible grain rice cultivars. J Stored Prod Res 2004;40:65-75.

Callaghan A. Insecticide resistance: mechanism and detection methods. Sci Progress 1991;75:423-38.

Carvalho AFU, Melo VMM, Craveiro AA, Machado MIL, Bantim MB, Rabelo EF. Larvicidal activity of the essential oil from *Lippia sidoides* Cham against *Aedes aegypti* L. Mem Inst Oswaldo Cruz 2003;98:569-71.

Chaithong U, Choochote W, Kansuk K, Jitpakdi A, Tippawangkosol P, Chaiyasit D, et al. Larvicidal effect of pepper plants on *Aedes aegypti* (L.) (Diptera: Culicidae). J Vector Ecol 2006;31:138-44.

Champakaew D, Junkum A, Chaithong U, Jitpakdi A, Riyong D, Sanghong R, et al. *Angelica sinensis* (Umbelliferae) with proven repellent properties against *Aedes aegypti*, the primary dengue fever vector in Thailand. Parasitol Res 2015;114:2187-98.

Chareonviriyaphap T, Bangs MJ, Suwonkerd W, Kongmee M, Corbel V, Ngoen-Klan R. Review of insecticide resistance and behavioral avoidance of vectors of human diseases in Thailand. Parasit Vectors 2013;6:280.

Chaubey MK. Fumigant toxicity of essential oils against rice weevil *Sitophilus oryzae* L. (Coleoptera: Curculionidae). J Biol Sci 2011;11:411-6.

Choi WS, Park BS, SK K, Lee SE. Repellent activities of essential oils and monoterpenes against *Culex pipiens pallens*. J Am Mosq Control Assoc 2002;18:348-51.

Choochote W, Tuetun B, Kanjanapothi D, Rattanachanpichai E, Chaithong U, Chaiwong P, et al. Potential of crude seed extract of celery, *Apium graveolens* L., against the mosquito *Aedes aegypti* (L.) (Diptera: Culicidae). J Vector Ecol 2004;29:340-6.

Choochote W, Chaiyasit D, Kanjanapothi D, Rattanachanpichai E, Jitpakdi A, Tuetun B, et al. Chemical composition and antimosquito potential of rhizome extract and volatile oil derived from *Curcuma aromaticata* against *Aedes aegypti* (Diptera: Culicidae). J Vector Ecol 2005;30:302-9.

Choochote W, Chaithong U, Kamsuk K, Rattanachanpichai E, Jitpakdi A, Tippawangkosol P, et al. Adulticidal activity against *Stegomyia aegypti* (Diptera: Culicidae) of three *Piper* spp. Rev Inst Med Trop S Paulo 2006;48:33-7.

Choovattanapakorn N, Yanola J, Lumjuan N, Saengamsook J, Somboon P. Characterization of metabolic detoxifying enzymes in an insecticide resistant strain of *Aedes aegypti* harboring homozygous S989P and V1016G *kdr* mutations. Med Entomol Zool 2017;68:19-26.

Clements AN. The biology of mosquitoes. Vol. 1: Development, nutrition and reproduction. Chapman and Hall: London; 1992.

Dauterman WC. Insect metabolism: Extramicrosomal. In: Kerkut GA, and Gilbert LI. (eds.), Comprehensive Insect Physiology, Biochemistry, and Pharmacology, Pergamon Press; Oxford 1985;12:713-30.

David H. Medical herbalism: The science and practice of herbal medicine. Healing Arts Press 2010:571.

Day MF. The distribution of alkaline phosphatase in insects. Australian J Sci Res 1948;2:31-42.

Dharmagadda VSS, Naik SN, Mittal PK, Vasudevan P. Larvicidal activity of *Tagetes patula* essential oil against three mosquito species. Biores Technol 2005;96:1235-40.

Dua VK, Pandey AC, Dash AP. Adulticidal activity of essential oil of *Lantana camara* leaves against mosquitoes. Indian J Med Res 2010;131:434-9.

Durbin AP. A dengue vaccine. Cell 2016;166:1.

El-Akhal F, Lalami AEO, Zoubi YE, Greche H, Guemmouh R. Chemical composition and larvicidal activity of essential oil of *Origanum majorana* (Lamiaceae) cultivated in Morocco against *Culex pipiens* (Diptera: Culicidae). Asian Pac J Trop Biomed 2014;4:746-50.

El-Bassal MM. Combined effects of insect growth regulators and gamma radiation on *Culex pipiens* [M.Sc. Thesis]. Fac Sci: Cairo Univ; 1993.

El-Kady GA, Kamel NH, Mosleh YY, Bahght IM. Comparative toxicity of two bioinsecticides (spinotoram and vertemic) compared with methomyl against *Culex pipiens* and *Anopheles multicolor*. World J Agric Sci 2008;4:198-205.

Ellman GL, Courtney K, Andres V, Featherstone RM. A new and rapid colorimetric determination of acetylcholinesterase activity. Biochem. Pharmacol 1961;7:88-95.

El-Sheikh TA. Effects of application of selected insect growth regulators and plant extracts on some physiological aspects of the black cutworm, *Agrotis ipsilon* (Huf) [Ph. D. Thesis]. Fac Sci: Ain Shams Univ; 2002.

Emtithal AES, Thanaa AEB. Efficacy of some insecticides on field populations of *Culex pipiens* (Linnaeus) from Egypt. J Basic Appl Zoo 2012;65:62-73.

Farzaei MH, Abbasabadi Z, Ardekani MRS, Rahimi R, Farzaei F. Parsley: a review of ethnopharmacology, phytochemistry and biological activities. J Trad Chin Med 2013;33:815-26.

Fetrow CW, Avila JR. Professional's handbook of complementary and alternative medicine. Spring house corporation: Pennsylvania; 1999.

Feyereisen R. Insect P450 enzymes. Annu Rev Entomol 1999;44:507-33.

Finney DJ. Probit analysis, 3rd edn. Cambridge University Press: New York; 1971.

Fradin MS, Day JF. Comparative efficacy of insect repellents against mosquito bites. N Engl J Med 2002;347:13-8.

Gamil WE, Mariy FM, Youssef LA, Abdel Halim SM. Effect of Indoxacarb on some biological and biochemical aspects of *Spodoptera littoralis* (Boisd.) larvae. Ann Agric Sci 2011;56:121-6.

Garrett M, Bradley TJ. The pattern of osmotic regulation in larvae of the mosquito *Culiseta inornata*. J Exp Biol 1984;113:133-41.

Ghoneim K, Amer M, Al-Daly A, Mohammad A, Khadrawy F, Mahmoud MA. Disturbed acid and alkaline phosphatase activities in desert locust *Schistocerca gregaria* (Forskal) (Orthoptera: Acrididae) by extracts from the khella plant *Ammi visnaga* L. (Apiaceae). Int J Adv Res 2014;2:584-96.

Ghosh A, Chowdhury N, Chandra G. Plant extracts as potential mosquito larvicides. Indian J Med Res 2012;135:581-98.

Govindarajan M, Sivakumar R. Mosquito adulticidal and repellent activities of botanical extracts against malarial vector, *Anopheles stephensi* Liston (Diptera:Culicidae). Asian Pac J Trop Med 2011;4:941-7.

Govindarajan M, Sivakumar R. Adulticidal properties of *Cardiospermum halicacabum* plant extracts against three important vector mosquitoes. Eur Rev Med Pharmacol Sci 2012;16:95-104.

Govindarajan M, Sivakumar R. Larvicidal, ovicidal, and adulticidal efficacy of *Erythrina indica* (Lam.) (Family: Fabaceae) against *Anopheles stephensi*, *Aedes aegypti*, and *Culex quinquefasciatus* (Diptera: Culicidae). Parasitol Res 2014;113:777-91.

Grant DF. Evolution of glutathione S-transferase subunits in Culicidae and related Nematocera: Electrophoretic and immunological evidence for conserved enzyme structure and expression. Insect Biochem 1991;21:435-45.

Grant DF, Dietze EC, Hammock BD. Glutathione S-transferase isozymes in *Aedes aegypti*: purification, characterization, and isozyme specific regulation. *Insect. Biochem* 1991;4:421-33.

Grieco JP, Achee NL, Chareonviriyaphap T. A new classification system for the actions of IRS chemicals traditionally used for malaria control. *PLoS One* 2007;2:e718.doi:10.1371/journal.pone.0000716.

Grundy DL, Still CC. Inhibition of acetylcholinesterases by pulegone-1,2-epoxide. *Pest Biochem Physio* 1985;23:383-8.

Gubler DJ. Epidemic dengue/dengue hemorrhagic fever as a public health, social and economic problem in the 21st century. *Trends Microbiol* 2002;10:100-3.

Habig WH, Pabst MJ, Jakoby WB. Glutathione S-transferases: The first enzymatic step in mercapturic acid formation. *J Biol Chem* 1974;249:7130-9.

Hafeez F, Akram W, Suhail A, Khan MA. Adulticidal action of ten citrus oils against *Aedes albopictus* (Diptera: Culicidae). *Pak J Agri Sci* 2010;47:241-4.

Harel M, Kryger G, Rosenberry TL, Mallender WD, Lewis T, Fletcher RJ, et al. Three-dimensional structures of *Drosophila melanogaster* acetylcholinesterase and its complexes with two potent inhibitors. *Prot Sci* 2000;9:1063-72.

Hassett CC, Jenkins DW. The uptake and effect of radiophosphorus in mosquitoes. *Physiol Zool* 1951;24:257-66.

Hayes JD, Wolf CR. Role of glutathione transferase in drug resistance. In: Sies H, Ketterer B (eds) Glutathione conjugation: Mechanisms and Biological Significance. Academic Press: London; 1988;316-56.

Helmy N, Bakr RFA, Nawwar GA, Ibrahim SE, Helmy OM. Biochemical effects of some agricultural waste extracts against *Culex pipiens* (Diptera: Culicidae). *Egypt Acad J biol Sci* 2010;2:75-81.

Hemingway J, Karunaratne SHPP. Mosquito carboxylesterases: a review of the molecular biology and biochemistry of a major insecticide resistance mechanism. *Med Vet Entomol* 1998;12:1-12.

Hemingway J, Ranson H. Insecticide resistance in insect vectors of human disease. Ann Rev Entomol 2000;45:369-89.

Hopp MJ, Foley J. Global-scale relationships between climate and the dengue fever vector *Aedes aegypti*. Clim Change 2001;48:441-63.

Huron EN, Abbas AA, El-Hamid NAA, Nada MS, Amin TR. Toxicity and acute macromolecular abnormalities induced by some plant extracts against the Cowpea aphid; *Aphis carricivora* Koch. J Plant Prot and Path 2016;7:445-9.

Imam H, Zarnigar, Sofi G. Mosquito larvicidal efficacy of *Acorus calamus* extracts against *Aedes aegypti* L. larvae. Asian Pac J Trop Dis 2014;4:181-5.

Insun D, Choochote W, Jitpakdi A, Chaithong U, Tippawangkosol P, Pitasawat B. Possible site of action of *Kaempferia galanga* in killing *Culex quinquefasciatus* larvae. Southeast Asian J Trop Med Pub Hlth 1999;30:195-9.

Intrirach J, Junkum A, Tuetun B, Choochote W, Chaithong U, Jitpakdi A, et al. Chemical constituents and combined larvicidal effects of selected essential oils against *Anopheles cracens* (Diptera: Culicidae). Psyche 2012;Article ID 591616.

Intrirach J, Junkum, A, Lumjuan N, Chaithong U, Jitpakdi A, Riyong D, et al. Antimosquito property of *Petroselinum crispum* (Umbellifereae) against the pyrethroid resistant and susceptible strains of *Aedes aegypti* (Diptera: Culicidae). Environ Sci Pollut Res Int 2016;23:23994-4008.

Isman MB. Plant essential oils for pest and disease management. Crop Prot 2000;19:603-8.

Jagadesshwaran U, Vijayan V. Biochemical characterization of deltamethrin resistance in a laboratory-selected strain of *Aedes aegypti*. Parasitol Res 2009;104:1431-8.

Jankowska M, Rogalska J, Wyszkowska J, Stankiewicz M. Molecular targets for components of essential oils in the insect nervous system-A review. Molecules 2018;23:34.

Jayaraman M, Senthilkumar A, Venkatesalu V. Evaluation of some aromatic plant extracts for mosquito larvicidal potential against *Culex quinquefasciatus*, *Aedes aegypti*, and *Anopheles stephensi*. Parasitol Res 2015;114:1511-18.

Jeeshna MV, Mallikadevi T, Paulsamy S. Screening of the weed plant species, *Croton bonplandianum* Baill. for larvicidal activity of *Aedes aegypti*. J Biopest 2010;3:192-4.

Joffe T, Gunning RV, Allen GR, Kristensen M, Alptekin S, Field LM, et al. Investigating the potential of selected natural compounds to increase the potency of pyrethrum against houseflies *Musca domestica* (Diptera:Muscidae). Pest Manag Sci 2011;68:178-84.

Jones SC, Bryant JL, Harrison SA. Behavioral responses of the bed bug to permethrin-impregnated ActiveGuardTM fabric. Insects 2013;4:230-40.

Kalimuthu K, Murugan K, Panneerselvam C, Hwang JS. Mosquito larvicidal activity of *Cadaba indica* lam leaf extracts against the dengue vector, *Aedes aegypti*. Asian J Plant Sci Res 2012;2:633-7.

Kamgang B, Marcombe S, Chandre F. Insecticide susceptibility of *Aedes aegypti* and *Aedes albopictus* in Central Africa. Parasit Vectors 2011;4:79.

Kamita SG, Samra AI, Liu JY, Cornel AJ, Hammock BD. Juvenile hormone (JH) esterase of the mosquito *Culex quinquefasciatus* is not a target of the JH analog insecticide methoprene. PLoS One 2011;6:e28392.doi:10.1371/journal.pone.0028392.

Kjanijou M, Jiraungkoorskul K, Kosai P, Jiraungkoorskul W. Effect of *Murraya paniculata* leaf extract against *Culex quinquefasciatus* larva. Asian J Biol Sci 2012;5:201-8.

Klowden MJ. Physiological systems in insects. Oxford: Elsevier Inc; 2007.

Knight KL, Stone A. A catalog of the mosquitoes of the world (Diptera: Culicidae), 2nd ed. Entomol Soc Am; 1977.

Knio KM, Usta J, Dagher S, Zournajian H, Kreydiyyeh S. Larvicidal activity of essential oils extracted from commonly used herbs in Lebanon against the seaside mosquito, *Ochlerotatus caspius*. Bioresour Technol 2008;99:763-8.

Knudsen AB, Sloof R. Vector-borne disease problems in rapid urbanization: new approaches to vector control, Bull. WHO 1992;70:1-6.

Koch HJ. The absorption of chloride ions by the anal papillae of Diptera larvae. J Exp Biol 1938;15:152-60.

Koodalingam A, Mullainadhan P, Arumugam M. Effects of extract of soapnut *Sapindus emarginatus* on esterases and phosphatases of the vector mosquito, *Aedes aegypti* (Diptera: Culicidae). Acta Trop 2011;118:27-36.

Koodalingam A, Mullainadhan P, Rajalakshmi A, Deepalakshmi R, Ammu M. Effect of a Bt-based product (Vectobar) on esterases and phosphatases from larvae of the mosquito *Aedes aegypti*. Pest Biochem Physiol 2012;104:267-72.

Kovendan K, Murugan K, Kumar PM, Thiagarajan P, William SJ. Ovicultural, repellent, adulticidal and field evaluations of plant extract against dengue, malaria and filarial vectors. Parasitol Res 2013;112:1205-19.

Kweka E, Nyindo M, Mosha F, Silva A. Insecticidal activity of the essential oil from fruits and seeds of *Schinus terebinthifolia* Raddi against African malaria vectors. Parasit Vectors 2011;4:129.

Laranja AT, Manzatto AJ, Bicudo HEM de C. Effects of caffeine and used coffee grounds on biological features of *Aedes aegypti* (Diptera, Culicidae) and their possible use in alternative control. Genet Mol Biol 2003;26:419-29.

Li X, Schuler MA, Berenbaum MR. Molecular mechanisms of metabolic resistance to synthetic and natural xenobiotics. Annu Rev Entomol 2007;52:231-53.

Limsuwan S, Rongsriyam Y, Kerdpibule V, Apiwathnasorn C, Chiang GL, Cheong WH. Rearing techniques for mosquitoes. In: Supavej S, ed. Entomology malaria and filariasis practical entomology malaria and filariasis. Bangkok: Museum and Reference Center, Faculty of Tropical Medicine, Mahidol University; 1987.

Linde GA, Gazim ZC, Cardoso BK, Jorge LF, Tešević V, Glamoćlija J, Soković M, Colauto NB. Antifungal and antibacterial activities of *Petroselinum crispum* essential oil. Genet Mol Res 2016;15.doi: 10.4238/gmr.15038538.

Lumjuan N, Wicheer J, Leelapat P, Choochote W, Somboon P. Identification and characterisation of *Aedes aegypti* aldehyde dehydrogenases involved in pyrethroid metabolism. PLoS One 2014;9:e102746.

Lupi E, Hatz C, Schlagenhauf P. The efficacy of repellents against *Aedes*, *Anopheles*, *Culex* and *Ixodes* spp. A literature review. Travel Med Infect Dis 2013;11:374-411.

Macoris MLG, Andrigatti L, Takaku L, Glasser CM, Garbeloto VV, Bracco JE. Resistance of *Aedes aegypti* from the state of São Paulo, Brazil, to organophosphate insecticides. Mem Inst Oswaldo Cruz 2003;98:703-8.

Majerus P, Kissleva M, Norris FA. The role of phosphatases in inositol signaling reactions. J Biol Chem 1999;274:10669-72.

Manimaran A, Cruz MMJJ, Muthu C, Vincent S, Ignacimuthu S. Larvicidal and growth inhibitory activities of different plant volatile oils formulation against *Anopheles stephensi* (Liston), *Culex quinquefasciatus* Say and *Aedes aegypti* (L.). Int J Phytother Res 2013;3:38-48.

Mann RS, Kaufman PE. Natural product pesticides: their development, delivery and use against insect vectors. Mini Rev Org Chem 2012;9:185-202.

Mannervik B, Danielson UH. Glutathione transferases-structure and catalytic activity. CRC Crit Rev Biochem 1988;23:283-337.

Mansour SA, Bakr RFA, Hamouda LS, Mohamed RI. Adulicidal activity of some botanical extracts, commercial insecticides and their binary mixtures against the housefly, *Musca domestica* L. Egypt Acad J Biolog Sci 2012;5:151-67.

McLaughlin LA, Niazi U, Bibby J, David JP, Vontas J, Hemingway, et al. Characterization of inhibitors and substrates of *Anopheles gambiae* CYP6Z2. Insect Mol Biol 2008;17:125-35.

Mehta RM. Pharmaceutics -I. Vallabh Prakashan: Delhi; 2002.

Mirhaghparast SK, Zibaee A. Effects of pyriproxyfen on intermediary metabolism of rice striped stem borer, *Chilo suppressalis* walker (Lepidoptera: Crambidae). Proc Natl Acad Sci India Sect B Biol Sci 2016;86:187-97.

Modise SA, Ashafa AOT. Larvicidal, pupicidal and insecticidal activities of *Cosmos bipinnatus*, *Foeniculum vulgare*, and *Tagetes minuta* against *Culex quinquefasciatus* mosquitoes. Trop J Pharmaceu Res 2016;15:965-72.

Monnerat R, Dumas V, Ramos F, Pimentel L, Nunes A, Sujiet E, et al. Evaluation of different larvicides for the control of *Aedes aegypti* (Linnaeus) (Diptera: Culicidae) under simulated field conditions. Bioassay 2012;7:3.

Nardini L, Blanford S, Coetzee M, Koekemoera LL. Effect of *Beauveria bassiana* infection on detoxification enzyme transcription in pyrethroid resistant *Anopheles arabiensis*: a preliminary study. Trans R Soc Trop Med Hyg 2014;108:221-7.

Nawaz R, Rathor HR, Bilal H, Hassan SA, Khan IA. Adulticidal activity of *Olea vera*, *Linum usitatissimum*. and *Piper nigra* against *Anopheles stephensi* and *Aedes aegypti* under laboratory conditions. Iran J Arthropod Borne Dis 2011;5:2

Pandey SK, Upadhyay S, Tripathi AK. Insecticidal and repellent activities of thymol from the essential oil of *Trachyspermum ammi* (Linn) Sprague seeds against *Anopheles stephensi*. Parasitol Res 2009;105:507-12.

Panneerselvam C, Murugan K, Kovendan K, Kumar PM. Mosquito larvicidal, pupicidal, adulticidal, and repellent activity of *Artemisia nilagirica* (Family: Compositae) against *Anopheles stephensi* and *Aedes aegypti*. Parasitol Res 2012;111:2241-51.

Park BS, Choi WS, Kim JH, Kim KH, Lee SE. Monoterpenes from thyme (*Thymus vulgaris*) as potential mosquito repellents. J Am Mosq Control Assoc 2005;21:80-3.

Patil SV, Patil CD, Salunkhe RB, Salunke BK. Larvicidal activities of six plants extracts against two mosquito species, *Aedes aegypti* and *Anopheles stephensi*. Trop Biomed 2010;27:360-5.

Perumalsamy H, Kim JR, Oh SM, Jung JW, Ahn YJ, Kwon HW. Novel histopathological and molecular effects of natural compound pellitorine on larval midgut epithelium and anal gills of *Aedes aegypti*. PLoS One 2013;8:e80226. doi:10.1371/journal.pone.0080226.

Pitasawat B, Champakaew D, Choochote W, Jitpakdi A, Chaithong U, Kanjanapothi D, et al. Aromatic plant-derived essential oil:an alternative larvicide for mosquito control. Fitoterapia 2007;78:205-10.

Plernsub S, Saengamsook J, Yanola J, Lumjuan N, Tippawangkosol P, Sukontason K, et al. Additive effect of knockdown resistance mutations, S989P, V1016G and F1534C, in a heterozygous genotype conferring pyrethroid resistance in *Aedes aegypti* in Thailand. Parasit Vectors 2016;9:417.

Pickett CB, Lu AY. Glutathione S-transferases: gene structure, regulation, and biological function. Annu Rev Biochem 1989;58:743-64.

Pimsamarn S, Sornpengb W, Akksilp S, Paeporn P, Limpawitthayakul M. Detection of insecticide resistance in *Aedes aegypti* to organophosphate and synthetic pyrethroid compounds in the north-east of Thailand. Dengue Bull 2009;33:194-202.

Prapanthadara L, Koottathep S, Promtet N, Hemingway J, Ketterman AJ. Purification and characterization of a major glutathione S-transferase from the mosquito *Anopheles dirus* (Species B). Insect Biochem Mol Biol 1996;26:277-85.

Prapanthadara L, Promtet N, Koottathep S, Somboon P, Suwonkerd W, McCarroll L, et al. Mechanisms of DDT and permethrin resistance in *Aedes aegypti* from Chiang Mai, Thailand. Dengue Bull 2002;26:185-9.

Prapanthadara L, Reunkum W, Leelepat P, Suwan W, Yanola J, Somboon P. Glutathione-S-transferase isoenzymes and the DDTase activity in two DDT-resistant strains of *Aedes aegypti*. Dengue Bull 2005;29:183-91.

Putra RE, Ahmad I, Prasetyo DB, Susanti S, Rahayu R, Hariani N. Detection of insecticide resistance in the larvae of some *Aedes aegypti* (Diptera: Culicidae) strains from Java, Indonesia to Temephos, Malathion, and Permethrin. Int J Mosq Res 2016;3:23-28.

Ramirez GIJ, Logan JG, Loza-Reyes E, Stashenko E, Moores GD. Repellents inhibit P450 enzymes in *Stegomyia (Aedes) aegypti*. PLoS One 2012;7:e48698.

Ramsay JA. Exchanges of sodium and potassium in mosquito larvae. J Exp Biol 1953;30:79-89.

Ranson H, Burhani J, Lumjuan N, Black WC. Insecticide resistance in dengue vectors. TropIKA.net 2010;1:1-12.

Ravikumar S, Ali MS, Beula JM. Mosquito larvicidal efficacy of seaweed extracts against dengue vector of *Aedes aegypti*. Asian Pac J Trop Biomed 2011;143-6.

Ravindran J, Samuel T, Alex E, William J. Adulticidal activity of *Ageratum houstonianum* Mill. (Asteraceae) leaf extracts against three vector mosquito species (Diptera: Culicidae). Asian Pac J Trop Med 2012;2:177-9.

Ray A, Rao CGP, Srivevi R, Ramamurty PS. Changes in acid phosphatase activity in *Spodoptera litura* (Lepi: Noctuidae) during the post embryonic and adult development. Entomol 1984;9:161-7.

Redford S, Misch D. Cytological effects of ecdysterone on the midgut cells of the fresh fly, *Sarcophaga bullata*. J Cell Biol 1971;49:702-11.

Ribeiro BM, Guedes RNC, Olivira EE, Santos JP. Insecticidal resistance and synergism in Brasilian populations of *Sitophilus zeamais* (Coleoptera: Curculionidae). J Stored Prod Res 2003;39:21-31.

Roy A, Saraf S. Limonoids: overview of significant bioactive triterpenes distributed in plants kingdom. Boil Pham Bull 2006;29:191-201.

Sakthivadivel M, Thilagavathy D. Larvicidal and chemosterilant activity of the acetone fraction of petroleum ether extract from *Argemone mexicana* L. seed. *Biores Technol* 2003;89:213-6.

Sarita N, Moharil MP, Ghodki BS, Lande GK, Bisane KD, Thakare AS, et al. Biochemical analysis and synergistic suppression of indoxacarb resistance in *Plutella xylostella* L. *J Asia-Pacific Entomol* 2010;13:91-5.

Senthil-Nathan S, Chung PG, Murugan K. Effect of botanicals and bacterial toxin on the gut enzyme of *Cnaphalocrocis medinalis*. *Phytoparasitica* 2004;32:433-43.

Seo SM, Jung CS, Kang J, Lee HR, Kim SW, Hyun J, et al. Larvicidal and acetylcholinesterase inhibitory activities of Apiaceae plant essential oils and their constituents against *Aedes albopictus* and formulation development. *J Agric Food Chem* 2015;63:9977-86.

Service MW. Medical entomology for students. Cambridge: Cambridge University Press; 2008.

Shaalan EA, Canyon DV, Younes MW, Abdel-Wahab H, Mansour AH. A review of botanical phytochemicals with mosquitocidal potential. *Environ Int* 2005;31:1149-66.

Shekari M, Sendi JJ, Etebari K, Zibaei A, Shadparvar A. Effects of *Artemisia annua* L. (Asteracea) on nutritional physiology and enzyme activities of elm leaf beetle, *Xanthogaleruca luteola* Mull. (Coleoptera: Chrysomellidae). *Pest Biochem Physiol* 2008;91:66-74.

Shivakumar MS, Srinivasan R, Natarajan D. Larvicidal potential of some indian medicinal plant extracts against *Aedes aegypti* (L.). *Asian J Pharm Clin Res* 2013;6:77-80.

Slama K. Ecdysteroids: Insect hormones, plant defensive factors, or human medicine?. *Phytoparasit* 1993;21:3-8.

Smirle MJ, Lowery DT, Zurowski CL. Influence of neem oil on detoxication enzyme activity in the obliquebanded leafroller, *Choristoneura rosaceana*. Pest Biochem Physiol 1996;56:220-30.

Somwang P, Yanola J, Suwan W, Walton C, Lumjuan N, Prapanthadara LA, et al. Enzymes-based resistant mechanism in pyrethroid resistant and susceptible *Aedes aegypti* strains from northern Thailand. Parasitol Res 2011;109:531-7.

Subramaniam J, Kovendan K, Kumar PM, Murugan K, Walton W. Mosquito larvicidal activity of *Aloe vera* (Family: Liliaceae) leaf extract and *Bacillus sphaericus*, against Chikungunya vector, *Aedes aegypti*. Saudi J Biol Sci 2012;19:503-9.

Sukumar K, Perich MJ, Boobar LR. Botanical derivatives in mosquito control: a review. J Am Mosq Control Assoc 1991;7:210-36.

Sutherland ORW. Plant chemicals influencing insect behaviour. NZ Entomol 1977;6:222-8.

Sutthanont N, Choochote W, Tuetun B, Junkum A, Jitpakdi A, Chaithong U, et al. Chemical composition and larvicidal activity of edible plant-derived essential oils against the pyrethroid-susceptible and -resistant strains of *Aedes aegypti* (Diptera: Culicidae). J Vector Ecol 2010;35:106-15.

Stenhouse SA, Plernsub S, Yanola J, Lumjuan N, Dantrakool A, Choochote W, et al. Detection of the V1016G mutation in the voltage-gated sodium channel gene of *Aedes aegypti* (Diptera: Culicidae) by allele-specific PCR assay, and its distribution and effect on deltamethrin resistance in Thailand. Parasit Vectors 2013;6:253.

Thanigaivel A, Vasantha-Srinivasan P, Edwin ES, Ponsankar A, Selin-Rani S, Chellappandian M, et al. Development of an eco-friendly mosquitocidal agent from *Alangium salvifolium* against the dengue vector *Aedes aegypti* and its biosafety on the aquatic predator. Environ Sci Pollut Res Int 2017;doi:10.1007/s11356-017-9102-6.

Thanispong K, Sathantriphop S, Chareonviriyaphap T. Insecticide resistance of *Aedes aegypti* and *Culex quinquefasciatus* in Thailand. J Pestic Sci 2008;33:351-6.

Tong F, Bloomquist JR. Plant sssential oils affect the toxicities of carbaryl and permethrin against *Aedes aegypti* (Diptera: Culicidae). J Med Entomol 2013;50:826-32.

Traboulsi AF, Taoubi K, El-Haj S, Bessiere JM, Rammal S. Insecticidal properties of essential oils against the mosquito *Culex pipiens molestus* (Diptera: Culicidae). Pest Manag Sci 2002;58:491-5.

Van Asperen K. A study of housefly esterases by means of a sensitive colorimetric method. J Insect Physiol 1962;8:401-16.

Van Pelt-Verkuil E. Hormone mediate induction of acid phosphatase activity of the fat body of *Calliphora erythrocephala* prior to metamorphosis. J Insect Physiol 1979;55:965-73.

Vasantha-Srinivasana P, Senthil-Nathana S, Ponsankar A, Thanigaivel A, Edwina ES, Selin-Rani S, et al. Comparative analysis of mosquito (Diptera: Culicidae: *Aedes aegypti* Liston) responses to the insecticide Temephos and plant derived essential oil derived from *Piper betle* L. Ecotoxicol Environ Saf 2017;139:439-46.

Vieira JR, Leite RM, Lima IR, Navarro Do A, Bianco EM, Leite SP. Oviposition and embryotoxicity of *Indigofera suffruticosa* on early development of *Aedes aegypti* (Diptera: Culicidae). Evid Based Complement Altern Med 2012;Article ID 741638:5.

Vongsombath C, Pålsson K, Björk L, Borg-Karlson AK, Jaenson TG. Mosquito (Diptera: Culicidae) repellency field tests of essential oils from plants traditionally used in Laos. J Med Entomol 2012;49:1398-404.

Waliwitiya R, Kennedy CJ, Lowenberger CA. Larvicidal and oviposition-altering activity of monoterpenoids, trans-anethole and rosemary oil to the yellow fever mosquito *Aedes aegypti* (Diptera: Culicidae). Pest Manag Sci 2008;65:241-8.

Waliwitiya R, Belton P, Nicholson RA, Lowenberger CA. Effects of the essential oil constituent thymol and other neuroactive chemicals on flight motor activity and wing beat frequency in the blowfly *Phaenicia sericata*. Pest Manag Sci 2009;66:277-89.

Waliwitiya R, Nicholson RA, Kennedy CJ, Lowenberger CA. The synergistic effects of insecticidal essential oils and piperonyl butoxide on biotransformational enzyme activities in *Aedes aegypti* (Diptera: Culicidae). J Med Entomol 2012;49:614-23.

Wandscheer CB, Duque JE, da Silva MA, Fukuyama Y, Wohlke JL, Adelmann J, Fontana JD. Larvicidal action of ethanolic extracts from fruit endocarps of *Melia azedarach* and *Azadirachta indica* against the dengue mosquito *Aedes aegypti*. Toxicon 2004;44:829-35.

Wang JJ, Cheng WX, Ding W, Zhao ZM. The effect of insecticide dichlorvos on esterase activity extracted from the psocids, *Liposcelis bostrychophila* and *L. entomophila*. J Insect Sci 2004;4:1-5.

Wang KY, Yong Z, Yan WH, Ming XX, Xian LT. Influence of three diets on susceptibility of selected insecticides and activities of detoxification esterases of *Helicoverpa assulta* (Lepidoptera: Noctuidae). Pest Biochem Physiol 2010;96,51-5.

Warikoo R, Wahab N, Kumar S. Larvicidal potential of commercially available pine (*Pinus longifolia*) and cinnamon (*Cinnamomum zeylanicum*) oils against dengue fever mosquito, *Aedes aegypti* L. (Diptera: Culicidae). Acta Entomol Sin 2011;57:793-9.

Weaver SC, Costa F, Garcia-Blanco MA, Ko AI, Ribeiro GS, Saade G, Shi PY, et al. Zika virus: History, emergence, biology, and prospects for control. Antiviral Res 2016;130:69-80.

World Health Organization. Instructions for determining the susceptibility or resistance of mosquito larvae to insecticides. WHO/VBC/81.807. Geneva: World Health Organization; 1981.

World Health Organization. Report of the WHO informal consultation on the evaluation and testing of insecticides. CTD/WHOPES/IC/96.1, Control of Tropical Diseases Division. Geneva: World Health Organization; 1996.

World Health Organization. Test procedures for insecticide resistance in monitoring in malaria vectors, bio-efficacy and persistence of insecticide on treated surfaces: Report of the WHO informal consultation. Geneva: World Health Organization, 1998.

World Health Organization. Dengue: Guidelines for diagnosis, treatment, prevention and control. Geneva: World Health Organization; 2009.

World Health Organization. Dengue and severe dengue (Fact sheet N°117). Geneva: World Health Organization; 2016

World Health Organization. Dengue vaccine research. World Health Organization: Geneva, 2016.

World Health Organization. Monitoring and managing insecticide resistance in *Aedes aegypti* populations, interim guidance for entomologist. Geneva, Switzerland [Internet]. 2016 [cited 2016 Mar 3]. Available from: <http://www.who.int/csr/resources/publications/zika/insecticide-resistance/en/>.

Yaicharoen R, Kiatfuengfoo R, Chareonviriyaphap T, Rongnoparut P. Characterization of deltamethrin resistance in field populations of *Aedes aegypti* in Thailand. J Vector Ecol 2005;30:144-50.

Yang Y, Cheng J-Z, Singhal SS, Saini M, Pandya U, Awasthi S. Role of glutathione S-transferases in protection against lipid peroxidation. J Biol Chem 2001;276:19220-30.

Yanola J, Somboon P, Walton C, Nachaiwieng W, Prapanthadara L. A novel F1552/C1552 point mutation in the *Aedes aegypti* voltage-gated sodium channel gene associated with permethrin resistance. Pest Biochem Physiol 2010;96:127-31.

Yanola J, Somboon P, Walton C, Nachaiwieng W, Somwang P, Prapanthadara L. High-throughput assays for detection of the F1534C mutation in voltage-gated sodium channel in permethrin-resistant *Aedes aegypti* and the distribution of this mutation throughout Thailand. Tropical Med Int Health 2011;16:501-9.

Zaim M, Aitio A, Nakashima N: Safety of pyrethroid-treated mosquito nets. Med Vet Entomol 2000;14:1-5.

Zibaee A. Botanical insecticides and their effects on insect biochemistry and immunity. In: Stoytcheva M, editor. Pesticides in the Modern World - Pests Control and Pesticides Exposure and Toxicity Assessment. InTech; 2011. p. 55-68.


Zoubiri S, Baaliouamer A. Potentiality of plants as source of insecticide principles. J Saudi. Chem Soc 2014;18:925-38.



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved

LIST OF PUBLICATIONS

- 1) **Intirach J**, Junkum A, Lumjuan N, Chaithong U, Jitpakdi A, Riyong D, Wannasan A, Champakaew D, Muangmoon R, Chansang A, Pitasawat B. Antimosquito property of *Petroselinum crispum* (Umbelliferae) against the pyrethroid resistant and susceptible strains of *Aedes aegypti* (Diptera: Culicidae). Environ Sci Pollut Res Int 2016;23:23994-4008.



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved