

เอกสารอ้างอิง

1. Shapiro, B. and Troncone, L. : “ The metabolically directed radiotherapy of hyperthyroidism with radioiodine.” in *Thyroid disease : Clinical fundamentals and therapy*. CRC Press. Inc., Boca Raton, pp. 383-389, 1993.
2. Sobel, S.H. and Bramlet, R. : “ Iodine-131 treatment of hyperthyroidism.” in *Thyroid disease : Endocrinology, Surgery, Nuclear medicine, and Radiotherapy*, 2d ed. Lippincott-Raven Publishers., Philadelphia, pp. 297-311, 1997.
3. Brunn J., Block, U., Ruf G., Bos I., Kunze W.P., Scriba P.C. Volumetric analysis of thyroid lobes by real-time ultrasound. *Dtsch Med Wochenschr.* 1981; 106(41) : 1338-1340.
4. Jarløv A. E., Gjørup T., Hegedüs L. Accuracy of the clinical assessment of thyroid size. *Dan Med Bull.* 1991; 38 : 87-89.
5. Szebeni Á., Belezay É. New simple method for thyroid volume determination by ultrasonography. *J Clin Ultrasound.* 1992; 20 : 329-337.
6. Knudsen N., Bols Birgitte., Bülow I., Jørgensen T., Perrild H., Ovesen L., Laurberg P. Validation of ultrasonography of the thyroid gland for epidemiological purpose. *Thyroid.* 1999; 9(11) : 1069-1074.
7. Lucas K.J. (No date). Use of thyroid ultrasound volume in calculating radioactive iodine dose in hyperthyroidism [Online]. Available : <http://www.beachdoctor.com> [2003, October 14].

8. Kung A.W.C., Choi P., Lam K.S.L., Pun K.K., Wang C., Yeung R.T.T. Discriminant factors affecting early outcome of radioiodine treatment for Graves' disease. *Clinical Radiology*. 1990; 42 : 52-54.
9. Bruin T.W.A. de., Croon C.D.L., Klerk J.M.H. de., Isselt J.W. van. Standardized radioiodine therapy in Graves' disease : the persistent effect of thyroid weight and radioiodine uptake on outcome. *J Int Med*. 1994; 236 : 507-513.
10. Peters H., Fischer C., Bogner U., Reiners C., Schleusener H. Radioiodine therapy of Graves' hyperthyroidism : Standard vs. calculated ¹³¹I activity. Results from a prospective, randomized, multicentre study. *Eur J Clin Invest*. 1995; 25(3) : 186-193.
11. Berg G.E.B., Michanek A.M.K., Holmberg E.C.V., Fink M. Iodine-131 treatment of hyperthyroidism : Significance of effective half-life measurements. *J Nucl Med*. 1996; 37(2) : 228-232.
12. Catargi B., Leprat F., Guyot M., Vallu N., Ducassou D., Tabarin A. Optimized radioiodine therapy of Graves' disease : analysis of the delivered dose and of other possible factors affecting outcome. *Eur J Endocrinology*. 1999; 141 : 117-121.
13. Allahabadia A., Daykin J., Sheppard M.C., Gough S.C.L., Franklyn J.A. Radioiodine treatment of hyperthyroidism – prognostic factors for outcome. [Online]. Available : <http://jcem.endojournals.org> [2004, February 26].
14. Howarth D., Epstein M., Lan L., Tan P., Booker J. Determination of the optimal minimum radioiodine dose in patients with Graves' disease : a clinical outcome study. *Eur J Nucl Med*. 2001; 28 : 1489-1495.

15. Alexander E.K., Larsen P.R. High dose ^{131}I therapy for the treatment of hyperthyroidism caused by Graves' disease. [Online]. Available : <http://jcem.endojournals.org> [2003, October 14].
16. Water M.A., Christ-Crain M., Eckard B., Schindler C., Nitzsche E.U., Müller-Brand J., Müller B. Radioiodine therapy in hyperthyroidism : inverse correlation of pretherapeutic iodine uptake level and post-therapeutic outcome. *Eur J Clin Invest.* 2004; 34(5) : 365-370.
17. Erem C., Kandemir N., Hacıhasanoglu A., Ersoz H.O., Ukinc K., Kocak M. Radioiodine treatment of hyperthyroidism : prognostic factors affecting outcome. *Endocrine.* 2004; 25(1) : 55-60.
18. Cherry S.R., Sorenson J.A. and Phelps M.E. : "Internal radiation dosimetry." in *Physics in nuclear medicine*, 3rd ed. Saunders., Philadelphia, pp. 405-425, 2003.
19. Koutras D.A., Matovinovic J. and Vought R. : "The ecology of iodine." in *Endemic goitre and cretinism, iodine nutrition in health and disease*. Wiley Eastern Limited., New York, pp. 185-195, 1985.
20. สายพิน โชติวิเชียร, ปิยนิตย์ ธรรมาภรณ์พิลาส และ แสงโสม สีนะวัฒน์. "การได้รับไอโอดีนจากแหล่งต่าง ๆ ของประชากรไทย" (2545). [ระบบออนไลน์]. แหล่งที่มา <http://advisor.anamai.moph.go.th/243/24308.html> [2005, May 15].
21. Weetman A.P. Graves' disease. *N Engl J Med.* 2000; 343(17) : 1236-1248.
22. Lind P. Strategies of radioiodine therapy for Graves' disease. *Eur J Nucl Med.* 2002; 29(suppl. 2) : s453-s457.

23. Soveid M., Shaabani A., Ghaedi GH.H., Jafari S.M., Omrani GH. Prognostic factors in the relapse of Graves' disease following treatment with antithyroid drugs. *Iran J Med Sci.* 2003; 28(3) : 106-110.
24. Mayo Foundation for Medical Education and Research. Graves' disease. [Online]. Available : www.mayoclinic.com [2004, July 6].
25. Kalinyak J.E., McDougall I.R. How should the dose of iodine-131 be determined in the treatment of Graves' hyperthyroidism ?. *J Endocrinol Metab.* 2003; 88 : 975-977.
26. DeGroot L.J. (2005). Diagnosis and treatment of Graves' disease. [Online]. Available : www.thyroidmanager.org/Chapter11/11-text.htm [2005, February 27].
27. Marinelli L.D., Quinby E.H., Hine G.J. Dosage determination with radioactive isotopes. *Am J Roentgenol.* 1948; 59 : 260-281.
28. Vemulakonda U.S., Atkins F.B., Ziessman H.A. Therapy dose calculation in Graves' disease using early I-123 uptake measurements. *Clin Nucl Med.* 1996; 21 : 102-105.
29. Isselt J. W. van., Tetteroo E., Gils A. P. G. van., Rijk P. P. van., Klerk J. M. H. de. Computer tomography is not suited for thyroid volume measurements in patients with Graves' disease before radioiodine therapy. [Online]. Available : <http://www.library.uu.nl/digiarchief/dip/diss/1941365/c7.pdf> [2004, February 4].
30. Isselt J.W. van., Beekman F.J., Kamphuis C., Klerk J.H.H., Polman L.J., Gils A.P.G., Vincken K. and Rijk P.P. A comparison of methods for thyroid volume determination in patients with Graves' disease. *Eur J Nucl Med.* 2003; 30(4) : 525-531.
31. Lisboa H.R., Gross J.L., Orsolin A., Fuchs S. Clinical examination is not an accurate method of defining the presence of goiter in schoolchildren. *Clin Endo.* 1996; 45 : 471.

32. Wacharasin R., Rajatanavin R., Chailurkit L. and Mahachoklertwattana P., 1990, "Thyroid volume measurement by ultrasonography and urinary iodine level in Bangkok schoolchildren", Abstracts Annual Meeting of Endocrine Society of Thailand. Nov. 1990.
33. Langer P. Normal thyroid size versus goiter-postmortem thyroid weight and ultrasonographic volumetry versus physical examination. *Endo Exp.* 1988; 23 : 67.
34. Imseis R.E., Vanmiddlesworth L., Massie J.D., Bush A.J., Vanmiddlesworth N.R. Pretreatment with propylthiouracil but not methimazole reduces the therapeutic efficacy of iodine-131 in hyperthyroidism. *J Clin Endocrinol Metab.* 1998; 83 : 685-687.
35. Andrade V.A., Gross J.L., Maia A.L. The effect of methimazole pretreatment on the efficacy of radioactive iodine therapy in Graves' hyperthyroidism : one-year follow-up of a prospective, randomized study. *J Clin Endocrinol Metab.* 2001; 86 : 3488-3493.
36. Isselt J.W. van. Radioiodine therapy : from medical tradition to clinical science. [Online]. Available : <http://www.library.uu.nl/digiarchief/dip/diss/1941365/c1.pdf> [2004, February 4].
37. Dunning G.E., Schwartz G. Variability in human thyroid characteristic and estimates of dose from ingested iodine-131. *Health Phys.* 1981; 40 : 661-675.
38. Leslie W.D., Ward L., Solomon E.A., Ludwig S., Rowe R.C., Cowden E.A. A randomized comparison of radioiodine dose in Graves' hyperthyroidism. *J Clin Endocrinol Metab.* 2003; 88(3) : 978-983.
39. Coatney R.W. Ultrasound imaging : Principles and applications in rodent research. *ILAR Journal.* 2001; 42(3) : 233-241.

40. Støylen A. Basic ultrasound for clinicians. (2005). [Online]. Available :

<http://www.ntnu.no/~stoylen/strainrate/ultrasound/> (2005, May 26).



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright © by Chiang Mai University
All rights reserved