

เอกสารอ้างอิง

- ตระกูล พรหมจักร. (2552). *การสกัดเจลาตินจากหนังปลาแพะ* (วิทยานิพนธ์). เชียงใหม่: บัณฑิตวิทยาลัย มหาวิทยาลัยเชียงใหม่.
- สำนักวิจัยและพัฒนาประมงน้ำจืด กรมประมง กระทรวงเกษตรและสหกรณ์. (2552). *การเพาะและอนุบาลปลานิล*. กรมประมง.
- สำนักงานมาตรฐานผลิตภัณฑ์อุตสาหกรรม. (2548). *เจลาติน*. มอก. 799-2548.
- สำนักโภชนาการ กระทรวงสาธารณสุข. (2559). สารอาหารของปลานิลสด. สืบค้น 7 พฤศจิกายน 2559, จาก <http://www.thailovehealth.com/food/food-16072.html>
- วิลาศ รัตนานุกูล. (2555). *คอลลาเจน*. สืบค้น 14 มกราคม 2559, จาก <http://biology.ipst.ac.th/?p=949>
- Ali, A. M. M., Benjakul, S., Prodpran, T., & Kishimura, H. (2017). Extraction and Characterisation of Collagen from the Skin of Golden Carp (*Probarbus jullieni*), a Processing By-Product. *Waste and Biomass Valorization*, 1-9.
- Almeida, P. F., & Almeida, A. J. (2004). Cross-linked alginate–gelatine beads: a new matrix for controlled release of pindolol. *Journal of Controlled Release*, 97(3), 431-439.
- Anissa, H., Sabrine, S., Raouia, G., Oscar, M. A., Moncef, N., & Ali B. (2012). Functional, antioxidant and film-forming properties of tuna-skin gelatin with a brown algae extract. *International Journal of Biological Macromolecules*, 51, 477–483.
- AOAC. (2000). *Official methods of analysis of AOAC* (17th ed.). Washington DC: Association of Official Analytical Chemists.
- Atkins, E. D. T., Nieduszynski, I. A., Mackie, W., Parker, K. D., & Smolko, E. E. (1973a). Structural components of alginic acid. I. The crystalline structure of poly-beta-D-mannuronic acid: Results of x-ray diffraction and polarized infrared studies. *Biopolymers*, 12, 1865–1878

- Badii, F., & Howell, N. K. (2006). Fish gelatin: structure, gelling properties and interaction with egg albumen proteins. *Food Hydrocolloids*, 20, 630-640
- Bailey, A. J., & Light, N. D. (1989). Connective tissue in meat and meat products. London & New York: Elsevier Applied Science. pp. 238-242.
- Bartova, V., & Barta, J. (2009). Chemical composition and nutritional value of protein concentrates isolated from potato (*Solanum tuberosum* L.) fruit juice by precipitation with ethanol or ferric chloride. *Journal of Agricultural and Food Chemistry*, 57(19), 9028-9034.
- Bourtoom, T., Chinnan, M. S., Jantawat, P., & Sanguandeeikul, R. (2009). Recovery and characterization of protein precipitated from surimi wash-water. *LWT - Food Science and Technology*, 42, 599-605.
- BSI (British Standards Institution). (1975). *Methods for sampling and testing gelatin (physical and chemical methods)*. London: BSI.
- Chhabra, N. (2013). *Structure of collagen*. Retrieved January 14, 2017, from <http://www.namrata.co/structure-of-collagen-solution-to-problem-based-on-osteogenesis-imperfecta/>
- Cho, S. H., Jahncke, M. L., Chin, K. B., & Eun, J. B. (2006). The effect of processing conditions on the properties of gelatin from skate (*Raja Kenojei*) skins. *Food Hydrocolloids*, 20, 810-816.
- Choi, I., Choi, S. J., Chun, J. K., & Moon, T. W. (2006). Extraction yield of soluble protein and microstructure of soybean affected by microwave heating. *Journal of Food Processing and Preservation*, 30, 407-419.
- Cole, C. G. B. (2000). Gelatin. In J. F. Frederic (Ed.), *Encyclopedia of Food Science and Technology* (2nd). New York: John Wiley & Sons.
- Crum, L. A. (1995). Comments on the evolving field of sonochemistry by a cavitation physicist. *Ultrasonics Sonochemistry*, 2, S147-S152.
- D'Cruz, N. M., & Bell, L. N. (2005). Thermal unfolding of gelatin in solids as affected by the glass transition. *Journal of food science*, 70(2), E64-E68.

- Djabourov, M., Lechaire, J., & Gaill, F. (1993). Structure and rheology of gelatin and collagen gels. *Biorheology*, 30, 191–205.
- Duconseille, A., Astruc, T., Quintana, N., Meersman, F., & Sante-Lhoutellier, V. (2015). Gelatin structure and composition linked to hard capsule dissolution: a review. *Food Hydrocolloids*, 43, 360–376.
- Eastoe, J. E., & Leach, A. A. (1977). Chemical constitution of gelatin. In A. G. Ward, & A. Courts (Eds.), *The science and technology of gelatin* (pp. 73–107). New York: Academic Press.
- Eysturskarð, J., Haug, I. J., Elharfaoui, N., Djabourov, M., & Draget, K. I. (2009). Structural and mechanical properties of fish gelatin as a function of extraction conditions. *Food hydrocolloids*, 23(7), 1702–1711.
- Fang, Y., Li, L., Vreeker, R., Yao, X., Wang, J., Ma, Q., ... & Phillips, G. O. (2011). Rehydration of dried alginate gel beads: Effect of the presence of gelatin and gum arabic. *Carbohydrate polymers*, 86(3), 1145–1150.
- Fellows, P. J. (2000). Dielectric, ohmic and infrared heating. In P. J. Fellows (Ed.), *Food processing technology: principles and practice*. (2nd.) (p. 365–384). Cambridge: Woodhead Publishing Limited.
- Foegeding, E. A., Lanier, T. C., & Hultin, H.O. (1996). Characteristics of Edible Muscle Tissue. In Fennema, O.R. (Ed.), *Food Chemistry*, 3rd Ed. New York: Marcel Dekker.
- Gimenez, B., Turnay, J., Lizarbe, M. A., Montero, M. P., & Gomez-Guillen, M. C. (2005). Use of lactic acid for extraction of fish skin gelatin. *Food Hydrocolloids*, 19, 941–950.
- Gómez-Guillén, M. C. & Montero, P. (2001). Extraction of gelatin from megrim (*Lepidorhombus boscii*) skins with several organic acids. *Journal of Food Science*, 66(2), 213–216.
- Gómez-Guillén, M. C., Turnay, J., Fernandez-Diaz, M. D., Ulmo, N., Lizarbe, M. A., & Montero, P. (2002). Structural and physical properties of gelatin extracted from different marine species: a comparative study. *Food Hydrocolloids*, 16, 25–34.

- Gómez-Guillén, M. C., Pérez-Mateos, M., Gómez-Estaca, J., López-Caballero, E., Giménez, B., & Montero, P. (2009). Fish gelatin: a renewable material for developing active biodegradable films. *Trends in Food Science and Technology*, 20, 3-16.
- Gómez-Guillén, M. C., Giménez, B., López-Caballero, M. E., & Montero, M. P. (2011). Functional and bioactive properties of collagen and gelatin from alternative sources: a review. *Food Hydrocolloids*, 25, 1813-1827.
- Grant, G. T., Morris, E. R., Rees, D. A., Smith, P. J. C., & Thom, D. (1973). Biological interactions between polysaccharides and divalent cations. *FEBS Letters*, 32, 195-198.
- Guo, L., Colby, R. H., Lusignan, C. P., & Whitesides, T. H. (2003). Kinetics of triple helix formation in semidilute gelatin solutions. *Macromolecules*, 36, 9999-10008.
- Haihua, C., & Jun, L. (2009). Application of Response Surface Methodology for Microwave-assisted Extraction Optimization of Alaska Pollock Fishskin Gelatin. *Food and Fermentation Industries*, 8, 168-174.
- Haug, I. J., & Draget, K. I. (2011). Gelatin. In G. O. Phillips, & P. A. Williams (Eds.), *Handbook of Food Proteins* (pp. 98-115). Norway: Norwegian University of Science and Technology (NTNU).
- Haug, I. J., Draget, K. I., & Smidsrød, O. (2004). Physical and rheological properties of fish gelatin compared to mammalian gelatin. *Food Hydrocolloids*, 18, 203-213.
- Jambrak, A. R., Mason, T. J., Lelas, V., Herceg, Z., & Herceg, I. L. (2008). Effect of ultrasound treatment on solubility and foaming properties of whey protein suspensions. *Journal of Food Engineering*, 86(2), 281-287.
- Johnston-Banks, F. A. (1990). Gelatin. In P. Harris (Ed.), *Food gels* (pp. 233-289). London: Elsevier Applied Science Publishers.
- Jongjareonrak, A., Rawdkuen, S., Chaijan, M., Benjakul, S., Osako, K., & Tanaka, M. (2010). Chemical compositions and characterization of skin gelatin from farmed giant catfish (*Pangasianodon gigas*). *LWT - Food Science and Technology*, 43, 161-165.

- Kappe, C.O. (2008). Microwave dielectric heating in synthetic organic chemistry. *The Royal Society of Chemistry*, 37, 1127-1139.
- Karim, A. A., & Bhat, R. (2009). Fish gelatin: properties, challenges, and prospects as an alternative to mammalian gelatins. *Food hydrocolloids*, 23(3), 563-576.
- Kim, H. K., Kim, Y. H., Park, H. J., & Lee, N. H. (2013). Application of ultrasonic treatment of extraction of collagen from the skin of sea bass *Lateolabrax japonicus*. *Fish Science*, 79, 849-856.
- Kittiphattanabawon, P., Benjakul, S., Sinthusamran, S., & Kishimura, H. (2016). Gelatin from clown featherback skin: Extraction conditions. *LWT-Food Science and Technology*, 66, 186-192.
- Koli, J. M., Basu, S., Kannuchamy, N., & Gudipati, V. (2013). Effect of pH and ionic strength on functional properties of fish gelatin in comparison to mammalian gelatin. *Fishery Technology*, 50(2), 126-132.
- Kolodziejska, I., Skierka, E., Sadowska, M., Kolodziejski, W., & Niecikowska, C. (2008). Effect of extracting time and temperature on yield of gelatin from different fish offal. *Food Chemistry*, 107, 700-706.
- Krešić, G., Lelas, V., Jambrak, A. R., Herceg, Z., & Brnčić, S. R. (2008). Influence of novel food processing technologies on the rheological and thermophysical properties of whey proteins. *Journal of Food Engineering*, 87, 64-73.
- Laemmli, U. K. (1970). Cleavage of structural proteins during the assembly of head of bacteriophage T4. *Nature*, 227, 680-685.
- Lee, B. B., Ravindra, P., & Chan, E. S. (2013). Size and shape of calcium alginate beads produced by extrusion dripping. *Chemical Engineering & Technology*, 36(10), 1627-1642.
- Lee, K. Y., & Mooney, D. J. (2012). Alginate: properties and biomedical applications. *Progress in polymer science*, 37(1), 106-126.
- Li, X. W., & Li, H. J. (2012). Optimization of Microwave-assisted Extraction Process for Collagen from Pigskin. *Food Science*, 33(6), 11-14.

- Lupo, B., Maestro, A., Gutiérrez, J. M., & González, C. (2015). Characterization of alginate beads with encapsulated cocoa extract to prepare functional food: comparison of two gelation mechanisms. *Food Hydrocolloids*, 49, 25-34.
- Mackie, W., Perez, S., Rizzo, R., Taravel, F., & Vignon, M. (1983). Aspects of the conformation of polyguluronate in the solid-state and solution. *International Journal of Biological Macromolecules*, 5, 329-341.
- Morris, E. R., Rees, D. A., Thom, D., & Boyd, J. (1978). Chiroptical and stoichiometric evidence of a specific, primary, dimerization process in alginate gelation. *Carbohydrate Research*, 66, 145-154.
- Murado, M. A., Montemayor, M. I., Cabo, M. L., Vázquez, J. A., & González, M. P. (2012). Optimization of extraction and purification process of hyaluronic acid from fish eyeball. *Food and Bioproducts Processing*, 90(3), 491-498.
- Mutilangi, W. A. M., Panyam, D., & Kilara, A. (1996). Functional properties of hydrolysates from proteolysis of heat-denatured whey protein isolate. *Journal of Food Science*, 61, 270-274, 303.
- Muyonga, J. H., Cole, C. G. B., & Duodu, K. G. (2004). Extraction and physico-chemical characterisation of Nile perch (*Lates niloticus*) skin and bone gelatin. *Food Hydrocolloids*, 18, 581-592.
- Nguyen, E., Jones, O., Kim, Y. H. B., San Martin-Gonzalez, F., & Liceaga, A. M. (2017). Impact of microwave-assisted enzymatic hydrolysis on functional and antioxidant properties of rainbow trout *Oncorhynchus mykiss* by-products. *Fisheries Science*, 83(2), 317-331.
- Niu, L., Zhou, X., Yuan, C., Bai, Y., Lai, K., Yang, F., & Huang, Y. (2013). Characterization of tilapia (*Oreochromis niloticus*) skin gelatin extracted with alkaline and different acid pretreatments. *Food Hydrocolloids*, 33, 336-341.
- Norland, R. E. (1990). Fish gelatin. In M. N. Voight & J. K. Botta (Eds.), *Advances in fisheries technology and biotechnology for increased profitability* (pp. 325-333). Lancaster: Technomic Publishing Co.
- Ockerman, H. W., & Hansen, C. L. (1988). Glue and gelatin. In H. W. Ockerman, & C. L. Hansen (Eds.), *Animal by-product processing* (pp. 133-157). England: Ellis Horwood Ltd.

- Piculell, I., Bergfeldt, K., & Nilsson, S. (1995). Factors determining phase behavior of multi-component polymer system. In S. E. Harding, S. E. Hill, & J. R. Mitchell (Eds.), *Biopolymer mixtures* (pp. 13–35). UK: Nottingham University Press.
- Poppe, J. (1997). Gelatin. In A. Imeson (Ed.), *Thickening and gelling agents for food* (pp. 98–123). Glasgow, UK: Blackie Academic and Professional.
- Rahman, M. S., Al-Saidi, G. S., & Guizani, N. (2008). Thermal characterisation of gelatin extracted from yellowfin tuna skin and commercial mammalian gelatin. *Food Chemistry*, 108(2), 472–481.
- Ran, X. G. & Wang, L. Y. (2014). Use of ultrasonic and pepsin treatment in tandem for collagen extraction from meat industry by-product. *Journal of the Science of Food and Agriculture*, 94, 585–590.
- Riesz, P., & Kondo, T. (1992). Free radical formation induce by ultrasound and its biological implications. *Free Radical Biological Medicine*, 13, 247–270.
- Robin Corey. (2015). *Why is hydroxyproline important component of collagen?*. Retrieved February 18, 2017, from <https://www.quora.com/Why-is-hydroxyproline-important-component-of-collagen>
- Rowe, R. C., Sheskey, P. J., & Quinn, M. E. (2009) *Handbook of Pharmaceutical Excipients* (6th ed.). Italy: Pharmaceutical Press and American Pharmacists Association.
- Roy, A., Bajpai, A. K., & Bajpai, J. (2009). Designing swellable beads of alginate and gelatin for controlled release of pesticide (cypermethrin). *Journal of Macromolecular Science*□, Part A: *Pure and Applied Chemistry*, 46(9), 847–859.
- Sai-Ut, S., Jongjareonrak, A., & Rawdkuen, S. (2012). Re-extraction, recovery and characteristics of skin gelatin from farmed giant catfish. *Food Bioprocess Technology*, 5, 1197–1205.
- Sarabia, A. I., Gómez-Guillén, M. C., & Montero, P. (2000). The effect of added salts on the viscoelastic properties of fish skin gelatin. *Food Chemistry*, 70, 71–76.
- Schrieber, R., & Gareis, H. (2007). *Gelatine handbook: theory and industrial practice*. John Wiley & Sons.
- Scopes, R. K. (1994). Separation by precipitation. In R. K. Scopes (Ed.), *Protein purification: Principles and practice* (pp. 71–101). New York: Springer-Verlag.

- Shin, S., Yang, H. J., Kim, J., & Kim, J. (2011). Effect of temperature on ultrasound-assisted tryptic protein digestion. *Analytical Biochemistry*, 414, 125-130.
- Shyni, K., Hema, G. S., Ninan, G., Mathew, S., Joshy, C. G., & Lakshmanan, P. T. (2014). Isolation and characterization of gelatin from the skins of skipjack tuna (*Katsuwonus pelamis*), dog shark (*Scoliodon sorrakowah*), and rohu (*Labeo rohita*). *Food Hydrocolloids*, 39, 68-76.
- Shyni, K., Hema, G. S., Ninan, G., Mathew, S., Joshy, C. G., & Lakshmanan, P. T. (2014). Isolation and characterization of gelatin from the skins of skipjack tuna (*Katsuwonus pelamis*), dog shark (*Scoliodon sorrakowah*), and rohu (*Labeo rohita*). *Food Hydrocolloids*, 39, 68-76.
- Sila, A., Martinez-Alvarez, O., Haddar, A., Gómez-Guillén, M. C., Nasri, M., Montero, M. P., & Bougatef, A. (2015). Recovery, viscoelastic and functional properties of Barbel skin gelatine: Investigation of anti-DPP-IV and anti-prolyl endopeptidase activities of generated gelatine polypeptides. *Food Chemistry*, 168, 478-486.
- Singh, R. P., & Heldman, D. R. (2001). *Introduction to food engineering*. Gulf Professional Publishing.
- Singh, S., Rao, K. R., Venugopal, K., & Manikandan, R. (2002). Alteration in dissolution characteristics of gelatin-containing formulations. *Pharmaceutical technology*, 26(4), 36-54.
- Sinthusamran, S., Benjakul, S., & Kishimura, H. (2014). Characteristics and gel properties of gelatin from skin of seabass (*Lates calcarifer*) as influenced by extraction conditions. *Food Chemistry*, 152, 276-284.
- Soria, A. C., & Villamiel, M. (2010). Effect of ultrasound on the technological properties and bioactivity of food: a review. *Trends in Food Science & Technology*, 21, 323-331.
- Sun, L., Hou, H., Li, B., & Zhang, Y. (2017). Characterization of acid-and pepsin-soluble collagen extracted from the skin of Nile tilapia (*Oreochromis niloticus*). *International Journal of Biological Macromolecules*, 99, 8-14.
- te Nijenhuis, K. (1997). Thermoreversible networks: viscoelastic properties and structure of gels. *Advances Polymer Science*, 130, 1-267.
- Technic-farm. (2015). การเพาะพันธุ์ปลานิลในบ่อดิน. สืบค้น 7 พฤศจิกายน 2559, จาก http://technicfarm.blogspot.com/2015/03/blog-post_47.html

- Tu, Z. C., Huang, T., Wang, H., Sha, X. M., Shi, Y., Huang, X. Q., ... & Li, D. J. (2015). Physico-chemical properties of gelatin from bighead carp (*Hypophthalmichthys nobilis*) scales by ultrasound-assisted extraction. *Journal of food science and technology*, 52(4), 2166-2174.
- Uluko, H., Li, H., Cui, W., Zhang, S., Liu, L., Chen, J., ... & Lv, J. (2013). Response surface optimization of angiotensin converting enzyme inhibition of milk protein concentrate hydrolysates in vitro after ultrasound pretreatment. *Innovative Food Science and Emerging Technologies*, 20, 133-139.
- Waglay, A., Karboune, S., & Alli, I. (2014). Potato protein isolates: recovery and characterization of their properties. *Food Chemistry*, 142, 373-382.
- Wang, J., Sun, B., Liu, Y., & Zhang, H. (2014). Optimisation of ultrasound-assisted enzymatic extraction of arabinoxylan from wheat bran. *Food Chemistry*, 150, 482-288.
- Waterborg, J. H. (2002). The lowry method for protein quantitation. In: Walker J.M., editor. *The protein protocols handbook* (p. 7-9). 2nd ed. Totowa, NJ: Humana Press Inc.
- Weng, W., Zheng, H., & Su, W. (2014). Characterization of edible films based on tilapia (*Tilapia zillii*) scale gelatin with different extraction pH. *Food Hydrocolloids*, 41, 19-26.
- Zayas, J. F. (1997). Solubility of proteins. In *Functionality of proteins in food* (pp. 6-75). Berlin/UK: Springer-Verlag/Harwood Academic Publishers
- Zeeb, B., Saberi, A. H., Weiss, J., & McClements, D. J. (2015). Formation and characterization of filled hydrogel beads based on calcium alginate: factors influencing nanoemulsion retention and release. *Food Hydrocolloids*, 50, 27-36.
- Zhang, Z., Zhang, R., Zou, L., & McClement, D. J. (2016). Protein encapsulation in alginate hydrogel beads: effect of pH on microgel stability, protein retention and protein release. *Food Hydrocolloids*, 58, 308-315.
- Zhou, P., Mulvaney, S. J., & Regenstein, J. M. (2006). Properties of Alaska pollock skin gelatin: a comparison with Tilapia and pork skin gelatins. *Journal of Food Science*, 71, C313-C321.
- Zhou, Y., Zheng, J., Gan, R. Y., Zhou, T., Xu, D. P., & Li, H. B. (2017). Optimization of Ultrasound-Assisted Extraction of Antioxidants from the Mung Bean Coat. *Molecules*, 22(4), 638.