

REFERENCES

- Ahn, Y.-J., Kim, Y.-S. Kim, S.-I and Kim, H.-K. 2002. Composition with acaricidal activity derived from plants. Pages 17 pp *in*. (Naturobiotech Co., Ltd., S. Korea). Application: EP
- Ainge Gary, D., Lorimer Stephen, D., Gerard Philippa, J. and Ruf Leslie, D. 2002. Insecticidal activity of huperzine A from the New Zealand clubmoss, *Lycopodium varium*. *Journal of Agricultural and Food Chemistry* **50**:491-494.
- Apisariyakul, A., Wannareumol, P., Watanakitwichai, T, Apisariyakul, S. 1991. A study of some medicinal plants effective against oral *Streptococcus* spp. *Journal of Phamacology*. 13-15
- Arima, K. H., Imanaka, M., Kousaka, A., Fukuda, A., and Tamura, G. 1965. Studies on pyrrolnitrin. K: Isolation and properties of pyrrolnitrin. *Journal of Antibiotics* **18**:210-204.
- Arnason, J. T., Scott, I., Durst, T., and Philogene, B. J. R. 2003. Insectidal natural products from neotropical forests. Abstracts of Papers, 226th ACS National Meeting, New York, NY, United States, September 7-11, 2003:AGRO-017.
- Balza, F., Abramowski, Z., Tower, G., and Wiriyaichitra, P. 1989. Identification of proanthocyanidin polymer as the piscidal constituents of *Mammea siamensis*, *polygonum* and *Diospyros diepenhorstii*. *Phytochemistry* **28**:1827-1830.

- Bandara, K. A., Kumar, V., Jacobsson, U. and Molleyres, L. P. 2000. Insecticidal piperidine alkaloid from *Microcos paniculata* stem bark. *Phytochemistry* **54**:29-32.
- Begley, M., Crombie, L., Jones, R., and Palmer, C. 1987. Synthesis of Mamea coumarins Part 4. Stereochemical and regiochemical studies, and synthesis of (-)-mameaB/BB. *Journal of Chemistry Society Perkin Trans I*:353-357.
- Bejar, E., Reyes-Chilpa, R. and Jimenez-Estrada, M. 2000. Bioactive compounds from selected plants used in the XVI century Mexican traditional medicine. *Studies in Natural Products Chemistry* **24**:799-844.
- Brem, B., Seger, C., Pacher, T., Hofer, O., Vajrodaya, S. and H. Greger. 2002. Feeding deterrence and contact toxicity of Stemona alkaloids - a source of potent natural insecticides. *Journal of Agricultural and Food Chemistry* **50**:6383-6388.
- Brueggemann, M., McDonald, A. I., Overman, L. E., Rosen, M. D., Schwink, L., and Scott, J. P. 2003. Total Synthesis of (+-)-Didehydrostemofoline (Asparagine A) and (+-)-Isodidehydrostemofoline. *Journal of the American Chemical Society* **125**:15284-15285.
- Carpenter, I., McGarry, E., and Scheinmann, F. 1970. The neoflavonoids and 4-alkylcoumarins from *Mammea africana* G. Don. *Tetrahedron Letters* **46**:3983-3986.
- Carpenter, I., McGarry, E., and Scheinmann, F. 1971. Extractives from Guttiferae. Part XXI. The Isolation and structure of nine coumarins from the bark of *Mammea africana* G. Don. *Journal of Chemistry Society (C)*:3783-3790.

- Chari, M.S., Ramaprasad, G., Sitaramaiah, S. and Murthy, P.S.N. 1993. Bio-efficacy of neem formulations against *Spodoptera litura* (F). in tobacco nurseries. Botanical Pesticide in Integrated Pest Management. 145-153.
- Chiu, S. 1989. Recent advances in research on botanical insecticides in China. ACS Symposium Series 387:69-77.
- Committee on the Standardization of Fungicidal Tests. 1943. The slide-germination method of evaluating protectant fungicides, *Phytopathology* 33: 627-632.
- Crombie, L., Games, D., Haskins, N., and Reed, G. 1970a. Identification of new coumarins: four 7,8-annulated relatives of the mammea coumarins A/AA, A/AB, B/AA, and B/AB, and two of the 6-acyl family, B/AA (isomammein) and B/AB. *Tetrahedron Letters* 46:3979-3982.
- Crombie, L., Games, D., Haskins, N., and Reed, G. 1970b. Identification of nine new 5,6-annulated relatives of mammea coumarins B/BA, B/BB and B/BC. *Tetrahedron Letters* 46:3975-3978.
- Crombie, L., Games, D., Haskins, N., and Reed, G. 1972a. Extractives of *Mammea americana* L. Part III. Identification of new coumarin relatives of mammea B/BA, B/BB, and B/BC having 5,6-annulation and higher oxidation levels. *Journal of Chemistry Society Perkin Trans I*:2241-2247.
- Crombie, L., Games, D., Haskins, N., and Reed, G. 1972b. Extractives of *Mammea americana* l. Part IV. Identification of new 7,8-annulated relatives of the coumarins mammea A/AA, A/AB, B/AA and B/AC. *Journal of Chemistry Society Perkin Trans I*:2248-2253.

- Crombie, L., Games, D., Haskins, N., and Reed, G. 1972c. Extractives of *Mammea americana* L. Part V. The insecticidal compounds. Journal of Chemistry Society Perkin Trans I:2255-2260.
- Crombie, L., Games, D., and McCormick, A. 1966a. Isolation and structure of mammea A/BA, A/AB and A/BB: A group of 4-aryl-coumarin extractives of *Mammea americana* L. Tetrahedron Letters. Tetrahedron Letters 2:145-149.
- Crombie, L., Games, D., and McCormick, A. 1966b. Isolation and structure of mammea B/BA, B/BB, B/BC and C/BB: A group of 4-n-propyl- and 4-n-aryl-coumarin extractives of *Mammea americana* L. Tetrahedron Letters 2:151-156.
- Crombie, L., Games, D., and McCormick, A. 1967a. Extractives of *Mammea americana* L. part I. The 4-n-Alkylcoumarins. Isolation and structure of mammea B/BA, B/BB, B/BC and C/BB. Journal of Chemistry Society (C):2545-2552.
- Crombie, L., Games, D., and McCormick, A. 1967b. Extractives of *Mammea americana* L. Part II. The 4-Phenylcoumarins. Isolation and structure of mammea A/AA, A/A cyclo D, A/BA, A/AB and A/BB. Journal of Chemistry Society (C):2553-2558.
- Crombie, L., Jones, R., and Palmer, C. 1985. Synthesis of the insecticidal 1'-acetoxy-mammeins and surangin B. Tetrahedron Letters 26:2933-2936.
- Crombie, L., Jones, R., and Palmer, C. 1987a. Synthesis of Mammea coumarins. Part 2. Experiments in the mammea E series and synthesis of mammea E/AC. Journal of Chemistry Society Perkin Trans I:333-343.

- Crombie, L., Jones, R., and Palmer, C. 1987b. Synthesis of the *Mammea* coumarins. Part1. The coumarins of the *mammea* A,B and C series. *Journal of Chemistry Society Perkin Trans I*:317-331.
- Chung, H.-s., Hon, P.-m., Lin, G., But, P. P.-h., and Dong, H. 2003. Antitussive activity of *Stemona* alkaloids from *Stemona tuberosa*. *Planta Medica* **69**:914-920.
- Djerassi, C., Eisenbraun, E., Finnegan, R., and Gillbert, B. 1959. Naturally occurring oxygen heterocyclics V. Mammien. *Tetrahedron Letters* **1**:10-14.
- Djerassi, C., Eisenbraun, E., Finnegan, R., and Gillbert, B. 1960. occurring oxygen heterocyclics VII. The structure of mammein. *Journal of Organic Chemistry* **25**:164-169.
- Djerassi, C., Eisenbraun, E., Gillbert, B., Lenin, S., and Morris, M. 1958. Naturally occurring oxygen heterocyclics. II. Characterization of an insecticidal principle from *Mammea americana* L. *Journal of American Chemistry Society* **80**:3686-3691.
- Dalpiazz, M. A., and Dalpiazz, V. R. 2001. Aqueous tobacco extracts as insecticides. Pages 8 pp *in*. (Brazil). Application: BR
- Del Rio, J. A., Arcas, M. C., Botia, A. J. M., Baidez, G., Fuster, M. D. and Ortuno, A. M. 2000. Involvement of phenolic compounds in the antifungal defense mechanisms of *Olea europaea* L. and *Citrus* sp. *Recent Research Developments in Agricultural and Food Chemistry* **4**:331-341.
- Duyfjes, B. E. E. 1993. *Stemonaceae*. *Flora Malesiana Series I* **11**:399-409.

- Dussourd David, E. 2003. Chemical stimulants of leaf-trenching by cabbage loopers: natural products, neurotransmitters, insecticides, and drugs. *Journal of Chemical Ecology* **29**:2023-2047.
- English, L.M. 2001. Organic Gardening-Natural Insecticide. <http://www.luminet.net/~wenona/hydro/pest cont.htm>.
- Finnegan, R., Morris, M., and Djerassi, C. 1961. Naturally occurring oxygen heterocyclicsX. 4-Phenyl-5,7-dihydroxy-6-isovaleryl-8-isopentenyl-coumarin. *Journal of Organic Chemistry* **26**:1180-1184.
- Finnegan, R., and Mueller, W. 1965. Constituents of *Mammea americana* L. IV. The structure of mammeigin. *Journal of Organic Chemistry* **30**:2342-2344.
- Fujimoto, Y., Ichinohe, Y and Takii, S. 2000. Sesquiterpene pyridine alkaloids of *Hippocratea excelsa* and their use as insecticides for agricultural and horticultural use. Pages 9 pp *in*. (Nissan Chemical Industries, Ltd., Japan). Application: JP
- Games, D. 1972. Identification of 4-phenyl and 4-alkylcoumarins in *Mammea americana* L., *Mammea africana* G. Don and *Calophyllum inophyllum* by gas chromatography-mass spectrometry. *Tetrahedron Letters* **31**:3187-3190.
- Gehmann, K., Nyfeler, R., Leadbeater, A. J., Nevill, D., and Sozzi, D. 1990. CGA173506: a new phenylpyrrole fungicide for broad spectrum disease control. Pages 399-406 *in* Proceedings of the Brighton Crop Protection Conference-Pests and Diseases 2.
- Ghisalberti, E.L. 1993. Detection and isolation of bioactive natural products in S.M. Colegate and R.J. Molyneux (eds.), *Bioactive natural products: detection, isolation and structural determination*. CRC Press. Boca Raton. 9-57.

- Glasby, J. S. 1991. Dictionary of Plants Containing Secondary Metabolites. Taylor & Francis Inc., London.
- Godfrey, C. R. A. 1995. Agrochemicals from Natural products. Marcel Dekker, Inc, New York.
- Gonzalez-Coloma, A., Reina, M., Gutierrez, C., and Fraga, B. M. 2002. Natural insecticides: structure diversity, effects and structure-activity relationships. A case study. *Studies in Natural Products Chemistry* 26:849-879.
- Hamilton-Kemp, T. R., Archbold, D. D., Loughrin, J. H., Andersen, R. A., McCracken, Jr., Collins, R. W. and Fallik, E. 2000. Stimulation and inhibition of fungal pathogens of plants by natural volatile phytochemicals and their analogs. *Current Topics in Phytochemistry* 4:95-104.
- Hanawa, F., Tahara, S. and Towers, G. H. N. 2000. Antifungal nitro compounds from Skunk cabbage (*Lysichitum americanum*) leaves treated with cupric chloride. *Phytochemistry* 53:55-58.
- He, W., Van Puyvelde, L., De Kimpe, N., Verbruggen, L., Anthonissen, K., Van der Flaas, M., Bosselaers, J., Mathenge, S. G., and Mudida, F. P. 2002. Chemical constituents and biological activities of *Zanthoxylum usambarense*. *Phytotherapy Research* 16:66-70.
- Hinman, M. M., and Heathcock, C. H. 2001. A synthetic approach to the Stemona alkaloids. *Journal of Organic Chemistry* 66:7751-7756.
- Hostettmann, K., Marston, A. and Wolfender, J.L. 1995. in Hostettmann, K., Marston, A., Maillard M. and Hamberger, M. (eds). Proceeding of the Phytochemical society of Europe. *Phytochemistry of Plants used in Traditional Medicine*. Clarendon Press. Oxford. 17-45.

- Inoue, T., and Inoue, T. 2000. Agrochemical fungicides containing chitosan, salicylic acid, and magnesium for hybrid plants. Pages 5 pp *in*. (Okayama Oyo Kagaku Y. K., Japan). Application: JP
- Jacobson, M., Crosby, D. G. and Editors. 1971. Naturally Occurring Insecticides.
- Jakobi, H., Cramp, S. M., Dickhaut, J., Lindell, S., Tiebes, J., Asuncion, C. M., Jans, D., Hempel, W., Royalty, R. N., McComb, S. M., Thoenessen, M.-T., Waibel, J. M., and Salgado, V. L. 2003. Preparation of propargylamines as pesticides for the protection of plants and animals. Pages 68 pp *in*. (Bayer CropScience GmbH, Germany). Application: DE
- Jacobi, P. A., and Lee, K. 2000. Total Syntheses of (+)- and (-)-Stemoamide. *Journal of the American Chemical Society* **122**:4295-4303.
- Jiang, R.-W., Hon, P.-M., But, P. P.-H., Chung, H.-S., Lin, G., Ye, W.-C., and Mak, T. C. W. 2002. Isolation and stereochemistry of two new alkaloids from *Stemona tuberosa*. *Tetrahedron* **58**:6705-6712.
- Jiwajinda, S., Hirai, N., Watanabe, K., Santisopasri, V., Chuengsamarnyart, N., Koshimizu, K. and Ohigashi, H. 2001. Occurrence of the insecticidal 16,17-didehydro-16(E)-stemofoline in *Stemona collinsae*. *Phytochemistry* **56**:693-695.
- Jiyavoranan, T. 2001. Insecticides from *Acorus calamus* Linn. and *Stemona tuberosa* Lour. and their residues after application. Chiang Mai University, Chiang Mai.
- Joshi, B., Kamat, V., Govindachari, T., and Ganguly, A. 1969. Isolation and structure of surangin A and surangin B, two new coumarins from *Mammea longifolia* (weight) Planch and Triana. *Tetrahedron Letters* **25**:1453-1458.

- Kaklyugin, V. Y., Ismailov, V. Y., Ivanova, T. S., Litvinova, V. I., and Vyalykh, A. K. 2003. Search for biologically active natural substances as raw materials for pesticide production. *Agrokhimiya*:48-54.
- Kakuta, D., Hitotsuyanagi, Y., Matsuura, N., Fukaya, H., and Takeya, K. 2003. Structures of new alkaloids sessilifoliamides A-D from *Stemona sessilifolia*. *Tetrahedron* **59**:7779-7786.
- Kaltenegger, E., Brem, B., Mereiter, K., Kalchhauser, H., Kahlig, H., Hofer, O., Vajrodaya, S., and Greger, H. 2003. Insecticidal pyrido[1,2-a]azepine alkaloids and related derivatives from *Stemona* species. *Phytochemistry (Elsevier)* **63**:803-816.
- Kaweetipob, W., Mahidol, C., Prawat, H., and Ruchirawat, S. 2000. Chemical investigation of *Mammea siamensis*. *Pharmaceutical Biology (Netherlands)*. **38**:55-57.
- Ke, C. Q., He, Z. S., Yang, Y. P., and Ye, Y. 2003. A novel alkaloid from *Stemona parviflora*. *Chinese Chemical Letters* **14**:173-175.
- Kostecki, K., Engelmeier, D., Pacher, T., Hofer, O., Vajrodaya, S., Greger, H. 2004. Dihydrophenanthrenes and other antifungal stilbenoids from *Stemona* cf. *pierrei*. *Phytochemistry* **65**: 99-106.
- Kuo, C., and Chu, T.-T. 1978. A study of *Stemona* alkaloids. I. *Huaxue Xuebao* **36**:291-296.
- Kyung, S.-H., Lee, Y.-K. and Song, C. 2002. Insecticidal activity of some monoterpenoids. *Han'guk Nonghwa Hakhoechi* **45**:37-41.

- Liu, M., and Yang, G. 2001. Bioinsecticide prepared from abamectin and alkaloid extracted from plant and its preparing process. Pages 8 pp *in*. (Peop. Rep. China). Application: CN
- Maxwell, J. F. 1991. Botanical notes on the vascular fora of Chiang Mai. Natural History Siam Society **39**:71-83.
- Mahandru, M., and Ravindran, V. 1986. Surangin C, a coumarin from *Mammea longifolia*. *Phytochemistry* **25**:555-556.
- Mahidol, C., Kaweetipob, W., Prawat, H., and Ruchirawat, S. 2002. *Mammea* coumarins from the flowers of *Mammea siamensis*. *Journal of Natural Products* **65**:757-760.
- Marija, J. 1992. SPSS for Windows Advanced Statistics Release 5. Norusis/SPSS Inc., New York.
- Matolcsy, G., Nadasy, M., and Andrisk, V. 1988. *Pesticide Chemistry*. Elsevier Science Publisher, Budapest. 239 pp.
- Mayer, B.N., Ferrigni, N.R., Putmann, J.E., Jacobsin, L.B., Nichols, D.E. and McLaughlin, J.L. 1982. Brine shrimp: a convenient general bioassay for active plant constituents. *Planta Medica*. **45**, 31-34.
- McLaughlin, J.C. 1991. Crown gall tumours on potato discs and brine shrimp lethality: two simple bioassays for higher plant screening. In: Hostettmann, K. (ed.), *Methods in Biochemistry*, **6**, Assays for Bioactivity. London, Academic Press, 1-32.
- McMakin, P. 1993. *A Field guide to the flowering plants of Thailand*. White Lotus, Bangkok.

- Miyazawa, M., Fujioka, J. and Ishikawa, Y. 2002. Insecticidal compounds from *Evodia rutaecarpa* against *Drosophila melanogaster*. Journal of the Science of Food and Agriculture **82**:1574-1578.
- Morimoto, Y., Iwahashi, M., Kinoshita, T., and Nishida, K. 2001. Stereocontrolled total synthesis of the *Stemona* alkaloid (-)-stenine. Chemistry--A European Journal **7**:4107-4116.
- Morris, M., and Pagan, C. 1953. The isolation of the toxic principle of mammey. Journal of American Chemistry Society **75**:1489.
- Munehiro, N., Samir A.M., A., Junichi, K., Hiroaki, O., Tetsuo, I., and Matsumi, D. 2001. Antifeedant rings B and D opened limonoids from *Khaya senegalensis*. Journal of Natural Products **64**:1262-1265.
- Mungkornasawakul, P. 2001. Fungicide from *Acorus calamus* Linn., *Eugenia caryophyllus* Bullock Et Harrison and *Mammea siamensis* Kosterm. and their residues after application. Chiang Mai, Chiang Mai.
- Nakanishi, K., Goto, T., Ito, S., Natori, S., and Nozoe, S. 1975. Natural products chemistry. KODANSHA LTD. and ACADEMIC PRESS, INC, Tokyo and New York.
- Nevill, D., Nyfeler, R., and Sozzi, D. 1998. CGA 142705, a novel fungicide for seed treatment. Pages 65-71 in Proceeding of the Brington Crop Protection Conference 1988, British Crop protection Council, Farnham.
- Nigg, N. H., and Seigler, D. 1992. Phytochemical resources for medicine and agriculture. Plenum Press, New York.
- Norman, R. F., and Nuntavan, B. 1992. Thai medicinal plants. Prachachon Co. Ltd., Bangkok.

- Nunez Marvin, J., A. Guadano, A. Jimenez Ignacio, G. Ravelo Angel, A. Gonzalez-Coloma, and L. Bazzocchi Isabel. 2004. Insecticidal sesquiterpene pyridine alkaloids from *Maytenus chiapensis*. *Journal of Natural Products* **67**:14-18.
- Pacher, T., Seger, C., Engelmeier, D., Vajrodaya, S., Hofer, O., and Greger, H. Antifungal stilbenoids from *Stemona collinsae*. 2002. *Journal of Natural Products* **65**:820-827.
- Perry, A. S., Yamamoto, I., Ishaaya, I. and Perry, R. 1998. Insecticides in agriculture and environment. Narosa Publishing House, London.
- Pham, H.-D., Yu, B.-W., Chau, V.-M., Ye, Y., and Qin, G.-W. 2002. Alkaloids from *Stemona collinsae*. *Journal of Asian Natural Products Research* **4**:81-85.
- Pilli, R. A., and Ferreira de Oliveira, M. d. C. 2000. Recent progress in the chemistry of the *Stemona* alkaloids. *Natural Product Reports* **17**:117-127.
- Poobraseart, O., Constant, L., Beecher, C., Farnsworth, N., Kinghorn, A., Pezzuto, J., Cordell, G., Santisuk, T., and Reutrakul, V. 1998. Xanthones from the twigs of *Mammea siamensis*. *Phytochemistry* **47**:1661-1663.
- Pottoriff, L.P. 2001. Friendly Pesticides for Home Gardens. <http://www.ext.colostate.edu/pub/garden/02945.html>.
- Prachyawarakorn, V., Mahidol, C., and Ruchirawat, S. 2000. NMR study of seven coumarins from *Mammea siamensis*. *Pharmaceutical Biology* (Netherlands). **38**:58-62.
- Prain, D. 1905. note on the Roxburghiaceae. *Journal of the Asiatic Society of Bengal* **73**:39-44.
- Prucksunand, C., Khunawat, P., Wimolwattanapun, S., and Prucksunand, P. 1985. The effect of "non-tai-yak" (*Stemona curtisii*) on the action potential of

- isolated frog sciatic nerve. A preliminary report. Journal of the Medical Association of Thailand = Chotmai het thangphaet **68**:66-71.
- Ragsdale, N. and Seiber, J. 1999. Pesticides : Managing Risks and Optimizing Benefits. American Chemical Society, Washington, DC. 363 pp.
- Rahuman, A.A., Gopalakrishnan, G., Ghouse, B.S., Arumugam, S., Himalayan, B. 2000. Effect of *Feronia limonia* on mosquito larvae. Fitoterapia **71**: 553-555.
- Rates, S. M. K. 2001. Plant as Source of Drugs. Toxicon **39**:39603-39613.
- Reshmi, B. and Vijiyalakahmi, M. 1998. Effect of aqueous extracts of neem based pesticides on the mortality of second-stage juveniles of *Meloidogyne inconita*. Annals of Plant Protection Sciences. 6(1), 55.
- Reshmi, B. and Vijiyalakahmi, M. 1999. Seed coating of chickpea with neem based pesticidal formulations for the management of *Meloidogyne incognita*. Indian Journal of Nematology. 29(1), 28-32.
- Rodcharoen, J. and Wongsiri, S. 1977. Biopesticide: Toxicity, Safety, Development and Proper Use. Proceeding First International Symposium on Biopesticide. Chulalongkorn University Press. Bangkok.
- Rungrojsakul, M. 2001. Morphological Characteristic, Chromosome Number and Propagation of *Stemona* spp. 2001. Master's thesis. Chiang Mai University, Chiang Mai.
- Sanches, F.F. and Ohsawa, K. 1994. Natural Bio-Active Substances in Tropical Plants. NODAI Center for International Programs, Tokyo University of Agriculture, JSPS-DOST Program.
- Sakata, K., Aoki, K., Chang, C.-F., Sakurai, A., Tamura, S. and Murakoshi, S. 1978. Stemospironine, a new insecticidal alkaloid of *Stemona japonica* Miq.,

isolation, structural determination and activity. *Agricultural and Biological Chemistry* **42**:457-463.

Saralamp, P., Chuakul, W., Temsiririrkkul, R., and Clayton, T. 1980. *Medicinal Plants in Thailand*. Amarin Printing and publishing Public Co. Ltd., Bangkok.

Schmutterer, H. 1995. *The Neem Tree*. VCH, Weinheim, 696 pp.

Seger, C., Mereiter, K., Kaltenegger, E., Pacher, T., Greger, H., and Hofer, O. 2004. Two pyrrolo[1,2-a]azepine type alkaloids from *Stemona collinsae* CRAIB: structure elucidations, relationship to asparagine A, and a new biogenetic concept of their formation. *Chemistry & Biodiversity* **1**:265-279.

Siddiqui, B. S., T. Gulzar, S. Begum, F. Afshan, and F. A. Sattar. 2004. Two new insecticidal amide dimers from fruits of *Piper nigrum* LINN. *Helvetica Chimica Acta* **87**:660-666.

Souza Brito, R. M. A. 1996. How to Study the Pharmacology of Medicinal Plants in undeveloped Countries. *Journal of Ethnopharmacology* **54**:131-138.

Sparks, T. C., Crouse, G. D., and Durst, G. 2001. Natural products as insecticides: the biology, biochemistry and quantitative structure-activity relationships of spinosyns and spinosoids. *Pest Management Science* **57**:896-905.

Srion, V. 2001. The Natural Pesticides. *Journal of Natural Agriculture* **8**:23.

Strange, N. R. 1993. *Plant Disease Control*. Chapman&Hall, London.

Subramanian, T., and Sibi, M. P. 2003. Approaches to the total synthesis of stemona alkaloids. Abstracts of Papers, 225th ACS National Meeting, New Orleans, LA, United States, March 23-27, 2003:ORGN-428.

- Suzuki, K. 1934. Alkaloids of *Stemona* VI. An alkaloid, tuberostemonine, from the roots of *Stemona tuberosa* Loureiro. *Yakugaku Zasshi* **54**:573-579, Abstracts 596-579.
- Tan, R. X., Meng, J. C. and Hostettmann, K.. 2000. Phytochemical investigation of some traditional Chinese medicines and endophyte cultures. *Pharmaceutical Biology (Netherlands)* **38**:25-32.
- Tang, W., and Eisenbrand, G. 1992. Chinese Drugs of Plants origin. Chemistry, Pharmacology, and use in Traditional and Modern Medicine. Springer, Uerlag Berlin Heidelberg.
- Terry, L. A., Joyce, D. C., Adikaram, N. K. B. and Khambay, B. P. S. 2004. Preformed antifungal compounds in strawberry fruit and flower tissues. *Postharvest Biology and Technology* **31**:201-212.
- The university of Arizona. <http://ag.arizona.edu/pubs/garden/mg/pesticides/formulation.html>.
- Tringali, C., Spatafora, C., Cali, V. and Simmonds, M. S. 2001. Antifeedant constituents from *Fagara macrophylla*. *Fitoterapia* **72**:538-543.
- Ulubelen, A., Mericli, A. H., Mericli, F., Kilincer, N., Ferizli, A. G., Emekci, M. and Pelletier, S. W. 2001. Insect repellent activity of diterpenoid alkaloids. *Phytotherapy research* : PTR **15**:170-171.
- Velazquez, F., and Olivo, H. F. 2002. Synthesis of Bicyclic g-Ylidenetetrates. *Organic Letters* **4**:3175-3178.
- Vijiyalakshimi, M. and Rajagopal, R. 1999. Nematicidal efficiency of Neema-S1, an experimental formulation for neem seed treatments against *Heterodera cajani*

and *Meloidogyne incognita* in coepea and chickpea respectively. *Azadirachta*

A. Juss.. 217-222.

Vogel, H.G. and Vogel, W.H. 1997. Drug discovery and evaluation: pharmacological assays. Springer Verlag. Berlin. 1-755.

Wipf, P., Rector, S. R., and Takahashi, H. 2002. Total Synthesis of (-)-Tuberostemonine. *Journal of the American Chemical Society* 124:14848-14849.

Yadava, R. L. 1971. Use of essential oil from Mongolian *Acorus calamus* L. rhizomes. *journal of essential oil reserch* 7:191-193.

Yamamura, S., Lguchi, M., Nishiyama, A., and Niwa, M. 1971. Sesquiterpenes from *Acorus calamus* L. *Tetrahedron* 27:5419-5431.

Yang, Y.-C., Lee, S.-G., Lee, H.-K., Kim, M.-K., Lee, S.-H. and Lee, H.-S. 2002. A piperidine amide extracted from *Piper longum* L. fruit shows activity against *Aedes aegypti* mosquito larvae. *Journal of Agricultural and Food Chemistry* 50:3765-3767.

Zhao, L., and Zhao, Y. 1991. Manufacture of nontoxic insecticides from medicinal plants for agricultural use. Pages 8 pp *in*. (Peop. Rep. China). Application: CN

Zou, C., Hongzheng, F., Li, J., Lei, H., Fu, H., and Lin, W. 1999. New alkaloids from the root of *Stemona japonica* Miq. *Journal of Chinese Pharmaceutical Sciences* 8:185-190.